

References

- Acharyya, S. K., Ray, K. K. and Roy, D. K., 1989, Tectono-stratigraphy and emplacement history of the ophiolite assemblage from the Naga Hills and Andaman island-arc, India. *Jour. Geol. Soc. India*, 33, 4-18.
- Adachi, M. and Kojima, S., 1983, Geology of the Mt. Hikagedaira area, east of Takayama, Gifu Prefecture, central Japan. *Jour. Earth Sci., Nagoya Univ.*, 31, 37-67.
- Aitchison, J. C., Davis, A. M., Badengzhu, Luo, H., 2002, New constraints on the India-Asia collision: the Lower Miocene Gangrinboche conglomerates, Yarlung Tsangpo suture zone, SE Tibet. *Jour. Asian Earth Sci.*, 21, 251-263.
- Akabane, F., 1998MS, Sandstones in the Hota Group and Serpentine Sandstones in the Mineoka Belt, Boso Peninsula, central Japan. *Undergraduate Thesis*, University of Tsukuba, 68p. (in Japanese with English abstract)
- Akimoto, K., Oda, M., Okamura, M. and Koyama, A., 1990, Geological age and paleobathymetry of the Koma Group, Yamanashi Prefecture, central Japan. *Bull. Earthq. Res. Inst. Univ. Tokyo*, 65, 531-539. (in Japanese with English abstract)
- Alam, M., 1989, Geology and depositional history of Cenozoic sediments of the Bengal Basin of Bangladesh. *Palaeogeogr., Palaeoclim., Palaeoecol.*, 69, 125-139.
- Alam, M. K., Hasan, A.K.M.S., Khan, M.R. and Whitney, J.W., 1990, Geological map of Bangladesh. *Geological Survey of Bangladesh, Dhaka*, Scale 1: 1,000,000.
- Amano, K., 1986, South Fossa Magna. Multiple collision zone. *Chikyū (the Earth Monthly)*, 18, 581-585. (in Japanese)
- Amano, K., 1991, Multiple collision tectonics of the South Fossa Magna in Central Japan. *Modern Geol.*, 15, 315-329.
- Amano, K., Martin, A. J., Tanakadate, H., Kanaguri, S., Yoda, N. and Aizu, T., 1999, Tectonics and basin formation in the arc-arc collision zone: The example of the South Fossa Magna in central Japan. *Japan. Jour. Struc. Geol.*, 43, 11-20.
- Amano, K. and Taira, A., 1992, Two-phase uplift of Higher Himalayas since 17 Ma. *Geology*, 20, 391-394.
- Aoike, K., 1998, Collisional accretionary block of the Eo-Oligocene paleo-Izu-Bonin Arc in the Koma massif, the Izu Collision Zone (preliminary report). *Abs. 105th Ann. Meet. Geol. Soc. Japan*, p. 21. (in Japanese)
- Aoike, K., 1999, Tectonic evolution of the Izu Collision Zone. *Res. Rep. Kanagawa Prefect. Mus. Nat. Hist.*, 9, 111-151. (in Japanese with English abstract)
- Arai, S., 1980, Dunite-hartzburgite-chromitite complexes as refractory residue in the

- Sangun-Yamaguchi zone, western Japan. *Jour. Petrol.*, 21, 141-165.
- Arai, S., 1981, Igneous rocks and ultramafic rocks in the Mineoka Belt, Boso Peninsula. Excursion Guidebook, The 88th Annual Meeting of the Geological Society of Japan, 59-72. (in Japanese)
- Arai, S., 1991, The Circum-Izu Massif peridotite, central Japan, as back-arc mantle fragments of the Izu-Bonin arc system. In: Peters, Tj. *et al.*, eds., *Ophiolite Genesis and Evolution of the Oceanic Lithosphere*, Kluwer Academic Pub., Dordrecht, 801-816.
- Arai, S., 1992a, Chemistry of chromian spinel in volcanic rocks as a potential guide to magma chemistry. *Mineral. Mag.*, 56, 173-184.
- Arai, S., 1992b, Petrological and geological significance of detrital chromian spinel and detrital serpentinite. *Mem. Geol. Soc. Japan*, no. 38, 329-344. (in Japanese with English abstract)
- Arai, S., 1994a, The Circum-Izu Massif serpentine belt. *Geosci. Repts. Shizuoka Univ.*, 20, 175-185. (in Japanese with English abstract)
- Arai, S., 1994b, Characterization of spinel peridotites by olivine-spinel compositional relationships: Review and interpretation. *Chem. Geol.*, 113, 191-204.
- Arai, S. and Ishida, T., 1987, Petrological characteristics of serpentinites in the Kobotoke Group, the Sasago area, Yamanashi Prefecture: a comparison with other Circum-Izu Massif serpentinites. *Jour. Mineral. Petrol. Econ. Geol.*, 82, 336-344. (in Japanese with English abstract)
- Arai, S., Ito, T. and Ozawa, K., 1983, Ultramafic-mafic clastic rocks from the Mineoka Belt, central Japan. *Jour. Geol. Soc. Japan*, 89(5), 287-297. (in Japanese with English abstract)
- Arai, S., Ito, M., Nakayama, N. and Masuda, F., 1990, A suspect serpentinite mass in the Tokyo Bay area: petrology and provenance of serpentinite pebbles in the upper Cenozoic system in the Boso Peninsula, central Japan. *Jour. Geol. Soc. Japan*, 96(3), 171-179. (in Japanese with English abstract)
- Arai, S. and Okada, H., 1991, Petrology of serpentine sandstone as a key to tectonic development of serpentine belts. *Tectonophysics*, 195, 65-81.
- Arai, S., Shimokawa, K. and Takahashi, T., 1978, On the mode of emplacement of ultramafic-mafic rocks in the Setogawa belt, Shizuoka Prefecture. *Jour. Geol. Soc. Japan*, 84(11), 691-693. (in Japanese)
- Arai, S. and Takahashi, N., 1988, Relic plagioclase-bearing harzburgite from the Mineoka belt, Boso Peninsula, central Japan. *Jour. Mineral. Petrol. Econ. Geol.*, 83, 210-214. (in Japanese with English abstract)

- Arai, S. and Uchida, T., 1979, Estimation of the equilibrium condition of the ultramafic rocks in the Setogawa Belt, Shizuoka Prefecture. *Geosci. Repts, Shizuoka Univ.*, 4, 19-24. (in Japanese with English abstract)
- Arita, K., 1983, Origin of the inverted metamorphism of the Lower Himalayas, central Nepal. *Tectonophysics*, 95, 43-60.
- Arita, K., Dallmeyer, R. D. and Takasu, A., 1997, Tectonothermal evolution of the Lesser Himalaya, Nepal: Constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ ages from the Kathmandu Nappe. *Island Arc*, 6, 372-385.
- Asami, M. and Hoshino, M., 1980, Staurolite-bearing schists from the Hongu-san area in the Ryoike Metamorphic Belt, central Japan. *Jour. Geol. Soc. Japan*, 86(9), 581-591.
- Banerji, R. K., 1984, Post-Eocene biofacies, palaeoenvironments and palaeogeography of the Bengal Basin, India. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 45, 49-73.
- Banno, S., Sakai, C. and Otsuki, M., 1984, Thermal history of the Sanbagawa Metamorphic Rocks. *Jour. Geogr., Tokyo Geogr. Soc. Japan.*, 93(7), 515-527. (in Japanese)
- Barnes, S. J. and Roeder, P. L., 2001, The range of spinel compositions in terrestrial mafic and ultramafic rocks. *Jour. Petrol.*, 42(12), 2279-2302.
- Bossart, P. and Ottiger, R., 1989, Rocks of the Muree formation in Northern Pakistan: indicators of a descending foreland basin of late Paleocene to middle Eocene age. *Eclogae. Geol. Helv.*, 82(1), 133-165.
- Bouquillon, A., France-Lanord, C., Michard, A. and Tiercelin, J.-J., 1990, Sedimentology and isotopic chemistry of the Bengal Fan sediments: the denudation of the Himalaya. *Proceedings of the Ocean Drilling Program, Scientific Results*, 116, 43-58.
- Burbank, D. W., Beck, R. A. and Mulder, T., 1996, The Himalayan foreland basin. In: Yin, A. and Harrison, T. M. (eds), *The tectonic evolution of Asia*. Cambridge Univ. Press, New York, 149-188.
- Chitani, K., 1926, The Tertiary in the vicinity of Kakegawa and Sagara, Totomi. *Jour. Geogr.*, 38, 84-89. (in Japanese)
- Chitani, K., 1931, *Geology of the Shizuoka district*. With Geological Sheet Map at 1:75,000, Geol. Surv. Japan, 48p. (in Japanese with English abstract)
- Copeland, P. and Harrison, T. M., 1990, Episodic rapid uplift in the Himalaya revealed by $^{40}\text{Ar}/^{39}\text{Ar}$ analysis of detrital K-feldspar and muscovite, Bengal fan. *Geology*, 18, 354-357.
- Corrigan, J. D. and Crowley, K. D., 1990, Fission-track analyses of detrital apatites from

- Sites 717 and 718, Leg 116, central Indian Ocean. *Proceedings of the Ocean Drilling Program, Scientific Results*, 116, 75-92.
- Danbara, T., Iwano, H., Kasuya, M., Yamashita, T. and Sumii, T., 1992, The non-toxic heavy liquid SPT and its use. *Chisitsu News (Geology Monthly)*, no. 455, 31-36. (in Japanese)
- Dick, H. J. B. and Bullen, T., 1984, Chromian spinel as a petrogenetic indicator in abyssal and alpine-type peridotites and spatially associated lavas. *Contrib. Mineral. Petrol.*, 86, 54-76.
- Dickinson, W. R., Braid, L. S., Brakenridge, G. R., Erjavec, J. L., Ferguson, R. C., Inman, K. F., Kenep, R. A., Lindberg, F. A. and Ryberg, P. T., 1983, Provenance of North American Phanerozoic sandstones in relation to tectonic setting. *Geol. Soc. America Bull.*, 94, 222-235.
- Dickinson, W. R. and Suczek, C. A., 1979, Plate Tectonics and Sandstone Compositions. *America. Assoc. Petrol. Geologists Bull.*, 63, 2164-2182.
- Droop, G. T. R., 1987, A general equation for estimating Fe³⁺ concentrations in ferromagnesian silicates and oxides from microprobe analyses, using stoichiometric criteria. *Mineral. Mag.*, 51, 431-435.
- Dürr, S. B., 1996, Provenance of Xigaze fore-arc basin clastic rocks (Cretaceous, south Tibet). *Geol. Soc. America Bull.*, 108(6), 669-684.
- Eto, T., 1986, Stratigraphy of the Hayama Group in the Miura Peninsula, Japan. *Sci. Rep. Yokohama Nat. Univ. Sec. II*, 33, 67-105. (in Japanese with English abstract)
- Fujioka, K., Tanaka, T. and Aoike, K., 1995, Serpentine Seamount in Izu-Bonin and Mariana Forearcs: Observation by a submersible and its relation to onland serpentinite belt. *Jour. Geogr., Tokyo Geogr. Soc. Japan.*, 104(3), 473-494. (in Japanese with English abstract)
- Fujioka, K. and Saito, S., 1992, Composition of heavy minerals from sands and sandstones of the Izu-Bonin Arc, Leg 126. *Proceedings of the Ocean Drilling Program, Scientific Results*, 126, 155-169.
- Gani, M. R. and Alam, M. M., 2003, Sedimentation and basin-fill history of the Neogene clastic succession exposed in the southeastern fold belt of the Bengal Basin, Bangladesh: a high-resolution sequence stratigraphic approach. *Sed. Geol.*, 155, 227-270.
- Garzanti, E., Baud, A. and Mascle, G., 1987, Sedimentary record of the northward flight of India and its collision with Eurasia (Ladakh Himalaya, India). *Geodinamica Acta*, 1(4/5), 297-312.
- Garzanti, E., Critelli, S. and Ingersoll, R. V., 1996, Paleogeographic and paleotectonic

- evolution of the Himalayan Range as reflected by detrital modes of Tertiary sandstones and modern sands (Indus transect, India and Pakistan). *Geol. Soc. America Bull.*, 108(6), 631-642.
- Hall, R., Fuller, M., Ali, J. R. and Anderson, C. D., 1995, The Philippine Sea Plate: magmatism and reconstructions. In: Taylor, B. and Natland, J. (eds.), *Active margins and marginal basins: synthesis of western Pacific drilling results*. AGU Monograph, no. 88, 371-404.
- Hatai, K. and Koike, K., 1957, On some fossil Mollusca from Chiba Prefecture, Japan. *Japan. Jour. Geol. Geogr.*, 28, 77-90.
- Haughton, P. D. W., Todd, S. P. and Morton, A. C., 1991, Sedimentary provenance studies. In: Morton, A. C., *et al.*, (eds.), *Developments in Sedimentary Provenance Studies*. Geol. Soc. London, Spec. Pub. no. 57, 1-11.
- Hirano, N., Ogawa, Y., Saito, K., Taniguchi, H. and Sato, H., 2001, Multi-stage evolution of the Tertiary Mineoka ophiolite (Boso Peninsula, Japan) at a TTT triple junction in the NW Pacific as revealed by new geochemical and age constraints. *EOS Trans. AGU*, 82, 1265.
- Hirano, N. and Okuzawa, K., 2002, Occurrence of the sandstone included in the alkali-basalt lava flow from the western Mineoka Belt, Boso Peninsula, Japan, and its tectonic significance. *Jour. Geol. Soc. Japan*, 108(11), 691-700. (in Japanese with English abstract)
- Hirata, T. and Nesbitt, R. W., 1995, U-Pb isotope geochronology of zircon: Evaluation of the laser probe-inductively coupled plasma mass spectrometry technique. *Geochim. Cosmochim. Acta*, 59(12), 2491-2500.
- Hisada, K. and Arai, S., 1989, Itsukaichi serpentinites in the Kanto mountains, central Japan: possibility of eastern extension of Kurosegawa Tectonic Zone. *Ann. Rep., Inst. Geosci., Univ. Tsukuba*, 15, 78-82.
- Hisada, K. and Arai, S., 1993, Detrital chrome spinels in the Cretaceous Sanchu sandstone, central Japan: indicator of serpentinite protrusion into a fore-arc region. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 105, 95-109.
- Hisada, K. and Arai, S., 1995, Detrital chromian spinels in the Cretaceous Choshi Group. *Jour. Geol. Soc. Japan*, 101(5), 393-396. (in Japanese with English abstract)
- Hodges, K. V., Parrish, R. R. and Searle, M. P., 1996, Tectonic evolution of the central Annapurna Range, Nepalese Himalayas. *Tectonics*, 15(6), 1264-1291.
- Holtrop, J. F. and Keizer, J., 1970, Some aspects of the stratigraphy and correlation of the Surma Basin wells, East Pakistan. In: United Nations. Economic Commission for Asia and the Far East (eds), *Proceedings of the second*

- seminar on geochemical prospecting methods and techniques*. Mineral Resources Development Series no. 36, New York, United Nations, 143-155.
- Huot, F., Hébert, R., Varfalvy, V., Beaudoin, G., Wang, C., Liu, Z., Cotton, J. and Dostal, J., The Beimarang mélange (southern Tibet) brings additional constraints in assessing the origin, metamorphic evolution and obduction process of the Yarlung Zangbo ophiolite. *Jour. Asian Earth Sci.*, 21, 307-322.
- Ibaraki, M., 1981, Geologic ages of the Neogene complex in the South Fossa Magna based upon planktonic foraminifera. *Jour. Geol. Soc. Japan*, 87(1), 47-49. (in Japanese)
- Ibaraki, M., 1983, Occurrences of Middle Eocene planktonic foraminifers in a mollusca-bearing horizon and limestone of the Takisawa Formation, the Setogawa Group. *Jour. Geol. Soc. Japan*, 89(1), 57-59. (in Japanese with English abstract)
- Ibaraki, M., 1984, Middle-Late Eocene planktonic foraminiferal faunas from limestones of the Setogawa Group, central Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, 135, 401-414.
- Ibaraki, M., 1986a, Planktonic foraminiferal datum levels recognized in the Neogene sequence of the Kakegawa area, and their relationship with the lithostratigraphy. *Jour. Geol. Soc. Japan*, 92(2), 119-134. (in Japanese with English abstract)
- Ibaraki, M., 1986b, Neogene planktonic foraminiferal biostratigraphy of the Kakegawa area on the Pacific coast of central Japan. *Rept. Fac. Sci., Shizuoka Univ.*, 20, 39-173.
- Ibaraki, M., Tsuchi, R. and Takayanagi, T., 1983, Early Neogene planktonic foraminiferal biostratigraphy in the Kakegawa area, the Pacific coast of central Japan. *Repts. Fac. Sci., Shizuoka Univ.*, 17, 102-116.
- Iijima, A., Matsumoto, R. and Watanabe, Y., 1981, Geology and siliceous deposits in the Tertiary Setogawa Terrain of Shizuoka, central Honshu. *Jour. Fac. Sic., Univ. Tokyo, Sec. II*, 20(3), 241-276.
- Iijima, A., Watanabe, Y. and Matsumoto, R., 1984, Geologic age of the Setogawa-Mineoka Belt. In: Saito, T. *et al.* (eds.) *Biostratigraphy and international correlation of the Paleogene system in Japan*. Yamagata Univ., Yamagata, p. 69-73. (in Japanese)
- Ingersoll, R. V., Bullard, T. F., Ford, R. L., Grimm, J. P. and Pickle, J. D., 1984, The effect of grain size on detrital modes: A test of the Gazzi-Dickinson point-counting method. *Jour. Sed. Petrol.*, 54(1), 103-116.
- Ingersoll, R. V. and Sucek, C. A., 1979, Petrology and provenance of Neogene sand from Nicobar and Bengal Fans, DSDP Sites 211 and 218. *Jour. Sed. Petrol.*, 49(4), 1217-1228.

- Ishida, T., Arai, S. and Takahashi, N., 1988, The occurrence of picrite basalt in the Kobotoke group, the Hatsukari area, Yamanashi Prefecture, central Japan. *Jour. Mineral. Petrol. Econ. Geol.* 83, 43-50. (in Japanese with English abstract)
- Ishida, T., Arai, S. and Takahashi, N., 1990, Metamorphosed picrite basalts in the northern part of the Setogawa belt, central Japan. *Jour. Geol. Soc. Japan.*, 96(3), 181-191. (in Japanese with English abstract)
- Ishii, M. and Makino, Y., 1946, On the discovery of *Discocyclusina* in the Setogawa Series. *Jour. Geol. Soc. Japan.*, 52, 1-2. (in Japanese)
- Ishii, T., Robinson, P. T., Maekawa, H. and Fiske, R., 1992, Petrological studies of peridotites from diapiric serpentinite seamounts in the Izu-Ogasawara-Mariana forearc, Leg 125. *Proceedings of the Ocean Drilling Program, Scientific Results*, 125, 445-485.
- Ito, H., Sorkhabi, R. B., Tagami, T. and Nishimura, S., 1989, Tectonic history of granitic bodies in the South Fossa Magna region, central Japan: new evidence from fission-track analysis of zircon. *Tectonophysics*, 166, 331-344.
- Johnson, S. Y. and Nur Alam, A.M., 1991, Sedimentation and tectonics of the Sylhet trough, Bangladesh. *Geol. Soc. America Bull.*, 103, 1513-1527.
- Kanehira, K., Oki, Y., Sanada, S., Yako, M. and Ishikawa, F., 1968, Tectonic blocks of metamorphic rocks at Kamogawa, southern Boso Peninsula. *Jour. Geol. Soc. Japan*, 74(10), 529-534. (in Japanese with English abstract)
- Kaneoka, I., Takigami, Y., Tonouchi, S., Furuta, T., Nakamura, Y. and Hirano, M., 1981, Pre-Neogene volcanism in the central Japan based on K-Ar and Ar-Ar analysis. Abs. IAVCEI Symp. Tokyo and Hakone, p. 166.
- Kano, K. and Matsushima, N., 1988, The Shimanto Belt in the Akaishi Mountains, eastern part of southwest Japan. *Modern Geology*, 12, 97-126.
- Kato, S., Shibata, H. and Inoue, Y., 1991, Fossils from the Mikura Group in the lower course of the Oigawa River, Shizuoka Prefecture, Japan. *Bull. Mizunami Fossil.Mus.* 18, 53-66. (in Japanese with English abstract)
- Kato, S., Inoue, Y. and Ukai, M., 1992, Foraminifera from the Oigawa Group, Shizuoka Prefecture. *Bull. Mizunami Fossil.Mus.* 19, 379-391. (in Japanese with English abstract)
- Kawai, K., 1957, Geology of the Kamogawa District, Chiba Prefecture. *Jour. Japan. Assoc. Petrol. Tech.*, 22(6), 1-8. (in Japanese with English abstract)
- Kawakami, S., 2003, Paleogene radiolarian biostratigraphy in the Boso Peninsula, central Japan. *Abst. 2003 Japan Earth Planet. Sci. Joint Meeting*, G082-013.
- Khan, M. R. and Muminallah, M., 1980, Stratigraphy of Bangladesh. *Petroleum and*

- Mineral Resources of Bangladesh, Seminar and Exhibition*, Dhaka, Government of the People's Republic of Bangladesh, p. 35-40.
- Kimura, K., 1992, Petroprovinces of the Eocene-early Oligocene Shimanto Supergroup. *Mem. Geol. Soc. Japan*, no. 38, 299-309. (in Japanese with English abstract)
- Kimura, T. and Tokuyama, A., 1971, Geosynclinal prisms and tectonics in Japan. *Mem. Geol. Soc. Japan*, no. 6, 9-20.
- Koike, K., 1949, Geology of the Middle Part of the Boso Peninsula (II). *Bull. Physiograph. Sci. Res. Inst., Tokyo Imp. Univ.*, 3, 1-6. (in Japanese with English abstract)
- Kosaka, T. and Tsunoda, F., Geology of the Koma Massif in the western part of Yamanashi Prefecture. *Jour. Geol. Soc. Japan*, 75(3), 127-140. (in Japanese with English abstract)
- Koyama, A., 1991, Collision of the Kushigatayama Block with the Honshu Arc during the Middle Miocene. *Modern Geology*, 15, 331-345.
- Koyama, A., 1993, Collision of the Kushigatayama block during the Middle Miocene, northwestern part of South Fossa-Magna, central Japan. *Mem. Geol. Soc. Japan*, no. 42, 245-254. (in Japanese with English abstract)
- Krishnamurthy, P. and Cox, K. G., 1977, Picrite basalts and related lavas from the Deccan Traps of western India. *Contrib. Mineral. Petrol.*, 62, 53-75.
- Kumar, D., Mamallan, R. and Dwivedy, K. K., 1996, Carbonatite magmatism in northeast India. *Jour. Southeast Asian Earth Sci.*, 13(2), 145-158.
- Kushiro, I., 1960, Si-Al relation in clinopyroxenes from igneous rocks. *America. Jour. Sci.*, 258, 548-554.
- Kumon, F., Kiminami, K., Adachi, M., Bessho, T., Kawabata, K., Kusunoki, T., Nishimura, T., Okada, H., Okami, K., Suzuki, S. and Teraoka, Y., 1992, Modal compositions of representative sandstones from the Japanese Islands and their tectonic implications. *Mem. Geol. Soc. Japan*, no. 38, 385-401. (in Japanese with English abstract)
- Krumbein, W. C., 1939, Preferred orientation of pebbles in sedimentary deposits. *Jour. Geol.*, 47(7), 673-706.
- Krumbein, W. C., 1941, Measurement and geological significance of shape and roundness of sedimentary particles. *Jour. Sed. Petrol.*, 11, 64-72.
- Le Bas, M. J., 1962, The role of aluminum in igneous clinopyroxenes with relation to their parentage. *America. Jour. Sci.*, 260, 267-288.
- Leterrier, J., Maury, R. C., Thonon, P., Girard, D. and Marchal, M., 1982, Clinopyroxene composition as a method of identification of the magmatic affinities of paleo-volcanic series. *Earth Planet. Sci. Lett.*, 59, 139-154.

- Makiyama, J., 1941, Tertiary stratigraphy and structure in the lower Ooigawa area. *Mem. Lec. Commen. Prof. Yabe, M I. A. 60th Birth.*, 1-13. (in Japanese with English abstract)
- Makiyama, J., 1950, *Regional Geology of Central Japan*. Asakura Pub. Co., Tokyo, 233p.
- Makiyama, J. and Sakamoto, T., 1957, *Geological map and explanatory text of the geological map of Japan*, 1: 50,000, Mitsuke and Kakezuka, Geol. Surv. Japan, 43p. (in Japanese with English abstract)
- Mannan, A., 2002, Stratigraphic evolution and geochemistry of the Neogene Surma Group, Surma Basin, Sylhet, Bangladesh. Doctoral Thesis of University of Oulu, Finland, 161p.
- Maruo, Y. and Kizaki, K., 1983, Thermal structure in the nappes of the eastern Nepal Himalayas. In: Shams, F. A. (ed.), *Granites of Himalayas, Karakorum & Hindukush: Lahore*. Inst. Geol. Punjab Univ., 271-286.
- Marsaglia, K. M. and Ingersoll, R. V., 1992, Compositional trends in arc-related, deep-marine sand and sandstone: A reassessment of magmatic-arc provenance. *Geol. Soc. America Bull.*, 104, 1637-1649.
- Matsuda, T., 1989, Examination on the multiple collision hypothesis of the South Fossa Magna. *Chikyū (Earth Monthly)*, 11, 522-525 (in Japanese)
- Matsumoto, E., 1964, The Asagai-Poronai molluscs from Shizuoka Prefecture, central Japan. *Mem. Coll. Sci., Univ. Kyoto, Ser. B*, 31, 95-111.
- Matsumoto, E., 1966, Geology of the Palaeogene and lower Neogene Formations in the lower course of River Ooigawa, Shizuoka Prefecture, central Japan. *Mem. Coll. Sci., Univ. Kyoto, Ser. B*, 33, 115-133.
- Meigs, A. J., Burbank, D. W. and Beck, R. A., 1995, Middle-late Miocene (> 10 Ma) formation of the Main Boundary thrust in the western Himalaya. *Geology*, 23(5), 423-426.
- Miller, C., Thöni, M., Frank, W., Schuster, R., Melcher, F., Meisel, T. and Zanetti, A., 2003, Geochemistry and tectonomagmatic affinity of the Yungbwa ophiolite, SW Tibet. *Lithos*, 66, 155-172.
- Mitsunashi, T., Kikuchi, T., Suzuki, J., Hirayama, J., Nakajima, T., Oka, S., Kodama, K., Horiguchi, M., Katsurajima, S., Miyashita, M., Yazaki, K., Kageyama, K., Nasu, N., Kagami, H., Honza, E., Kimura, M., Nirei, H., Higuchi, S., Hara, Y., Huruno, K., Endo, T., Kawashima, S. and Aoki, S., 1979, *Explanatory text of the geological map of Tokyo Bay and adjacent areas*. Miscellaneous map series (20), scale 1:100,000, Geol. Surv. Japan, 91p. (in Japanese with English abstract)
- Miyashiro, A., 1953, Calcium-poor garnet in relation to metamorphism. *Geochim.*

- Cosmochim. Acta*, 4, 179-208.
- Mizuno, A., 1956, A preliminary note on the magafaunal zones of the Paleogene in northwestern Kyushu, Japan (For correlation of the Paleogene formations in Japan). *Bull. Geol. Surv. Japan*, 7(6), 25-34 (in Japanese with English abstract)
- Mochizuki, K., 1956, Geology of Shizuoka Prefecture. With Geologic Sheet map at 1:200,000. Shizuoka Prefecture, 57p. (in Japanese)
- Mohiuddin, M. M. and Ogawa, Y., 1996, Middle Eocene to early Oligocene planktonic foraminifers from the micritic limestone beds of the Heguri area, Mineoka Belt, Boso Peninsula. *Jour. Geol. Soc. Japan*, 102(7), 611-617.
- Mohiuddin, M. M. and Ogawa, Y., 1998a, Early Miocene pelagic sequences in the Mineoka Belt, Boso Peninsula, Japan. *Jour. Geol. Soc. Japan*, 104(1), 1-12.
- Mohiuddin, M. M. and Ogawa, Y., 1998b, Late Paleocene-middle Miocene pelagic sequences in the Boso Peninsula, Japan: New light on northwest Pacific tectonics. *Island Arc*, 7, 301-314.
- Molnar, P. and Tapponnier, P., 1975, Cenozoic tectonics of Asia: Effects of a continental collision. *Science*, 189, 419-426.
- Morimoto, N., 1988, Nomenclature of pyroxenes. *Mineral. Mag.*, 52, 477-493.
- Morton, A. C., 1991, Geochemical studies of detrital heavy minerals and their application to provenance research. In: Morton, A. C., et al., (eds), *Developments in Sedimentary Provenance Studies*. Geol. Soc. London, Spec. Pub. no. 57, 31-45.
- Naka, J., 1985, Seamount origin greenstones in the Setogawa Belt., *Chikyu (the Earth Monthly)*, 7(11), 663-667. (in Japanese)
- Naka, J., 1988, Volcanostratigraphic sequence of greenstones in the Setogawa Terrane, central Japan. *Modern Geology*, 12, 243-282.
- Nakajima, T., Makimoto, H., Hirajima, J. and Tokuhashi, S., 1981, *Geology of the Kamogawa district*. Quadrangle Series, scale 1:50,000, Geol. Surv. Japan, 107p. (in Japanese with English abstract, 7p.)
- Neumann, E. R., 1976, Two refinements for the calculation of structural formulae for pyroxenes and amphiboles. *Norsk Geologisk Tidsskrift*, 56, 1-6.
- Niida, K., Yamamoto, M., Yasui, M. and Haldar, D., 2003, Outline of geology and petrology of the Andaman ophiolite. *Abs. 110th Ann. Meet. Geol. Soc. Japan*, p. 54. (in Japanese)
- Noda, Y., Saito, S. and Nagata, T., 2002, Occurrence of *Mytilus tichanovitchi* from the Hota Group in the Boso Peninsula and its significance. *Mem. Fukui Pref. Dino. Mus.*, 1, 96-101. (in Japanese with English abstract)
- Ogasawara, K., Hisada, K. and Kitada, N., 1994, Early Miocene *Calyptogena* from the

- Aokiyama Formation, Hota Group, Boso Peninsula, Japan. *Ann. Rep., Inst. Geosci., Univ. Tsukuba*, 20, 33-37.
- Ogawa, Y., 1981, Tertiary tectonics of the Miura and Boso Peninsulas: Ophiolite and the Izu forearc sediments trapped to the Honshu arc. *Chikyu (The Earth Monthly)*, Tokyo, 3(7), 411-420. (in Japanese)
- Ogawa, Y., Horiuchi, K., Taniguchi, H. and Naka, J., 1985a, Collision of the Izu arc with Honshu and the effects of oblique subduction in the Miura-Boso Peninsulas. *Tectonophysics*, 119, 349-379.
- Ogawa, Y., Naka, J. and Taniguchi, H., 1985b, Ophiolite-based forearcs: a particular type of plate boundary. In: N. Nasu, *et al.* (eds.), *Formation of Active Ocean Margins*. Terra Sci. Pub., Tokyo, 719-746.
- Ogawa, Y. and Taniguchi, H., 1987, Ophiolitic mélangé in the forearc areas and the development of the Mineoka belt. *Sci. Rept. Fac. Sci. Kyushu Univ.*, 15, 1-23. (in Japanese with English abstract)
- Ogo, Y. and Hiroi, Y., 1991, Origin of various mineral assemblages of the Mineoka metamorphic rocks from Kamogawa, Boso Peninsula, central Japan: with special reference to the effect of high oxygen fugacity. *Jour. Mineral. Petrol. Econ. Geol.* 86, 226-240. (in Japanese with English abstract)
- Ohara, Y., 2003, Reviews on mantle peridotites from the Philippine Sea backarc spreading systems. *Rep. Hydrogr. Oceanogr. Res.*, 39, 63-83. (in Japanese with English abstract)
- Ohashi, F., 1980, An alkali olivine basalt and its related rocks from the Setogawa Group, Shizuoka Prefecture. *Jour. Geol. Soc. Japan*, 86(12), 799-815.
- Ohashi, F. and Shiraki, K., 1981, High-magnesia and high-silica volcanic rock in the Setogawa Ophiolite. *Jour. Mineral. Petrol. Econ. Geol.* 76, 69-79. (in Japanese with English abstract)
- Okada, H., 1971, Classification of sandstone: analysis and proposal. *Jour. Geol.*, 79(5), 509-525.
- Ono, A., 1981, Geology of the Takato-Kashio area, Ryoike belt, central Japan. *Jour. Geol. Soc. Japan*, 87(4), 249-257.
- Osozawa, S., 1986, Geology of the southern Setogawa Belt: Takakusayama and its western district, In: Nakagawa, H. *et al.* (eds.), *Essays in Geology, Professor N. Kitamura Commemorative Volume*, Professor Nobu Kitamura Taikan Kinenjigyokai, Sendai, 337-350. (in Japanese with English abstract)
- Osozawa, S., 1988, Accretionary process of the Tertiary Setogawa and Mikasa Groups, southwest Japan. *Jour. Geol.*, 96, 199-208.

- Osozawa, S., Sakai, T. and Naito, T., 1990, Miocene subduction of an active mid-ocean ridge and origin of the Setogawa Ophiolite, central Japan. *Jour. Geol.*, 98, 763-771.
- Otofujii, Y., Matsuda, T. and Nohda, S., 1985, Opening mode of the Japan Sea inferred from the paleomagnetism of the Japan arc. *Nature*, 317, 603-604.
- Otuka, Y., 1938, Geological study on the Fujimiyama Scarp in the southwestern part of Yamanashi Prefecture. *Geogr. Rev. Japan*, 15(12), 969-984. (in Japanese)
- Otuka, Y., 1941, Geology of around the Mt. Hoo and Jizo-dake, Yamanashi Prefecture. *Bull. Earthquake Res. Inst., Univ. Tokyo*, 19, 115-143. (in Japanese with English abstract)
- Otuka, Y. and Koike, K., 1948, Geology of the Middle Part of the Boso Peninsula. *Bull. Physiograph. Sci. Res. Inst., Tokyo Imp. Univ.*, 2, 31-32. (in Japanese with English abstract)
- Rahman, M. J. J. and Faupl, P., 2003a, $^{40}\text{Ar}/^{39}\text{Ar}$ multigrain dating of detrital white mica of sandstones of the Surma Group in the Sylhet Trough, Bengal Basin, Bangladesh. *Sed. Geol.*, 155, 383-392.
- Rahman, M. J. J. and Faupl, P., 2003b, The composition of the subsurface Neogene shales of the Surma group from the Sylhet Trough, Bengal Basin, Bangladesh. *Ibid.*, 155, 407-417.
- Reimann, K.-U., 1993, Geology of Bangladesh. *Berlin-Stuttgart, Gebruder Borntraeger*, 160p.
- Renard, V., Hakamura, K., Angelier, J., Azema, J., Bourgeois, J., Deplus, C., Fujioka, K., Hamano, Y., Huchon, P., Kinoshita, H., Labaume, P., Ogawa, Y., Seno, T., Takeuchi, A., Tanahashi, M., Uchiyama, A. and Vigeneresse J. L., 1984, Trench triple junction off central Japan: preliminary results of French-Japanese 1984 Kaiko cruise, Leg 2. *Earth Planet. Sci. Lett.*, 83, 243-256.
- Rickwood, P. C., 1968, On recasting analyses of garnet into end-member molecules. *Contrib. Mineral. Petrol.*, 18, 175-198.
- Saito, S., 1991, Strike-slip movement and strike-slip basin formation in the Mineoka Tectonic Belt in the Boso Peninsula, Central Japan. *Jour. Tect. Res. Grope Japan*, 36, 83-91. (in Japanese with English abstract)
- Saito, S., 1992, Stratigraphy of Cenozoic strata in the southern terminus area of Boso Peninsula, Central Japan. *Contr., Inst. Geol. Pal., Tbhoku University*, 93, 1-37. (in Japanese with English abstract)
- Saito, S. and Fujioka, K., 1992, Oceanic arc evolution estimated from volcanoclastic sand and sandstones: an example from the Izu-Bonin Arc. *Mem. Geol. Soc. Japan*, no.

- 38, 373-384. (in Japanese with English abstract)
- Saito, S. and Isomi, H., 1954, *Geology of the Akihasan district*. With Geological Sheet Map at 1: 50,000, Geol. Surv. Japan, 34p. (in Japanese with English abstract)
- Saito, S., Yoshida, T. and Aoki, K., 1992, Geochemistry and tectonic significance of volcanic and volcanoclastic rocks in the Boso Peninsula, Central Japan. *Res. Rep. Lab. Nucl. Sci., Tohoku Univ.* 25(2), 241-255. (in Japanese)
- Saito, T., 1960, Tertiary stratigraphy of the Kakegawa district, central Japan and its planktonic foraminifera. *Contr. Inst. Geol. Pal. Tohoku Univ.*, 51, 1-45. (in Japanese with English abstract)
- Sakagami, S., Ito, M., Endo, H., Okazaki, H., Kazaoka, O., Kamura, K., Kusuda, T., Kotake, N., Sato, K., Takahashi, N., Tokuhashi, S., Nirei, H., Hara, Y. and Matsushima, Y., 1997, Geology. In: Chiba Historical Material Research Foundation (ed), *Natural history of Chiba Prefecture, Japan., II*, Chiba Prefecture (government), p. 81-350. (in Japanese)
- Sakai, A., 1987, *Geology of the Itsukaichi district*. With Geological Sheet Map at 1: 50,000, Geol. Surv. Japan, 75p. (in Japanese with English abstract 5p.)
- Sakai, H., 1990, The Himalayas and the South Fossa Magna. *Kagaku*, 62(7), 445-450. (in Japanese)
- Sakamoto, T., Ogawa, Y. and Nakada, S., 1993, Origin of the greenstones in the Setogawa accretionary complex and their tectonic significance. *Jour. Geol. Soc. Japan*, 99(1), 9-28. (in Japanese with English abstract)
- Sameshima, T., 1960, Picrite basalt dykes in the Paleogene formation in central Japan. *Rept. Liber. Art. Sci. Fac. Shizuoka Univ. Sec. Nat. Sci.*, 3(1), 77-80.
- Sashida, K. and Ogawa, Y., 2003, Early Cretaceous radiolarians from red chert cropped out at Yohka Beach, Kamogawa City, Boso Peninsula. *Abst. 2003 Ann. Meet. Palaeont. Soc. Japan*, p. 83. (in Japanese)
- Sato, H., Taniguchi, H., Takahashi, N., Mohiuddin, M. M., Hirano, N. and Ogawa, Y., 1999, Origin of the Mineoka Ophiolite. *Jour. Geogr., Tokyo Geogr. Soc. Japan.*, 108(3), 203-215. (in Japanese with English abstract)
- Sato, Y. and Suzuki, T., 1991, The guide of the heavy mineral analysis. *Chishitsu News (Geology Monthly)*, no. 444, 21-28. (in Japanese)
- Searle, M. P., Waters, D. J., Rex, D. C. and Wilson, R. N., 1992, Pressure, temperature and time constraints on Himalayan metamorphism from eastern Kashmir and western Zaskar. *Jour. Geol. Soc., London*, 149, 753-773.
- Seno, T. and Maruyama, S., 1984, Paleogeographic reconstruction and origin of the Philippine Sea. *Tectonophysics*, 102, 53-84.

- Shiki, T., 1966, Studies on sandstones in the Maizuru Zone, southwest Japan II: Graded bedding and mineral composition of sandstones of the Maizuru Group. *Mem. Coll. Sci., Univ. Kyoto, Ser. B*, 27, 293-308.
- Shimazu, M. and Ishimaru, 1987, Neogene volcanic rocks of the eastern part of the Koma Mountain, Yamanashi Prefecture. *Jour. Mineral. Petrol. Econ. Geol.* 82, 382-394. (in Japanese with English abstract)
- Shimokawa, K. and Sugiyama, Y., 1982, Pebbles of ultramafic-mafic rocks in the Lower Miocene Mikasa Group, Shizuoka Prefecture, central Japan. *Jour. Geol. Soc. Japan*, 88(11), 915-918. (in Japanese with English abstract)
- Shimokawa, K. and Sugiyama, Y., 1983, Geology and metamorphism of the Takakusayama Group in the central part of Shizuoka Prefecture. *Bull. Geol. Surv. Japan*, 34(9), 435-440. (in Japanese with English abstract)
- Styles, M. T., Stone, P. and Floyd, J. D., 1989, Arc detritus in the Southern Uplands: mineralogical characterization of a 'missing' terrane. *Jour. Geol. Soc., London*, 146, 397-400.
- Sugimura, A., 1972, Plate boundaries in and around Japan. *Kagaku*, 42, 192-202. (in Japanese)
- Sugiyama, Y., 1980, Submarine sliding deposits and allochthonous blocks in the Tertiary Ooigawa Group, Shizuoka Prefecture, Central Japan. *Jour. Geol. Soc. Japan*, 86(10), 673-685. (in Japanese with English abstract)
- Sugiyama, Y., 1992, The Cenozoic tectonic history of the forearc region of southwest Japan, based mainly of the data obtained from the Shizuoka district. *Bull. Geol. Surv. Japan*, 43, 91-112. (in Japanese with English abstract)
- Sugiyama, Y., 1995, Geology of the northern Setogawa Belt in the Akaishi Mountains and the formation of the Setogawa accretionary complex. *Bull. Geol. Surv. Japan*, 46, 177-214. (in Japanese with English abstract)
- Sugiyama, Y. and Shimokawa, K., 1989, Forming process of the early Miocene accretionary prism (Setogawa Belt) of the Shimanto Terrain in the Akaishi Mountains. *Japan. Jour. Struct. Geol.* 34, 173-188. (in Japanese)
- Sugiyama, Y. and Shimokawa, K., 1990, Geology of the Shimizu District. With Geological Sheet Map at 1: 50,000, Geol. Surv. Japan. 103p. (in Japanese with English abstract, 8p.)
- Sugiyama, Y., Shimokawa, K., Sakamoto, T. and Hata, M., 1982, *Geology of the Shizuoka District*. With Geological Sheet Map at 1: 50,000, Geol. Surv. Japan. 82p. (in Japanese with English abstract, 9p.)
- Suzuki, Y., Akiba, F. and Kamiya, M., 1996, Latest Oligocene siliceous microfossils from

- Hota Group in southern Boso Peninsula, eastern Honshu, Japan. *Jour. Geol. Soc. Japan*, 102(12), 1068-1071. (in Japanese)
- Suzuki, Y., Kodama, K. and Mitsuhashi, T., 1990, *Geology of the Nago district*. With Geological Sheet Map at 1: 50,000, Geol. Surv. Japan, 48p. (in Japanese with English abstract, 3p.)
- Suzuki, Y., Kondo, K. and Saito, T., 1984, Latest Eocene planktonic foraminifers from the Mineoka Group, Boso Peninsula. *Jour. Geol. Soc. Japan*, 90(7), 497-499. (in Japanese)
- Takahashi, N., 1994, "Alkali basalt-clastic sequence" in the western end of the Mineoka Belt, Boso Peninsula, Japan. *Jour. Nat. Hist. Mus. and Inst., Chiba*, 3(1), 1-18. (in Japanese)
- Takasawa, K., 1976, Olivine from the Takayama picrite in the Setogawa Group, Shizuoka Prefecture, central Japan. *Jour. Geol. Soc. Japan*, 82(12), 765-771. (in Japanese with English abstract)
- Takeuchi, M., 1986, Detrital garnet in Paleozoic-Mesozoic sandstone from the central part of the Kii Peninsula. *Jour. Geol. Soc. Japan*, 92(4), 289-306. (in Japanese with English abstract)
- Tamura, J., Kaneko, T. and Niitsuma, N., 1984, Geology of the southern part of the Koma Mountains, western part of Yamanashi Prefecture, central Japan. *Geosci. Repts, Shizuoka Univ.*, 10, 23-53. (in Japanese with English abstract)
- Tanaka, S. and Adachi, M., 1999, Occurrence of gneiss clasts with Mg-rich garnets from the Sawado conglomerate in the northeastern Mino terrane: source of detrital Mg-rich garnets in the Jurassic sandstone of the Mino terrane. *Jour. Geol. Soc. Japan*, 105(3), 193-199. (in Japanese with English abstract)
- Taniguchi, H. and Ogawa, Y., 1990, Occurrence, chemistry and tectonic significance of alkali basaltic rocks in the Miura Peninsula, central Japan. *Jour. Geol. Soc. Japan*, 96(2), 101-116. (in Japanese with English abstract)
- Taniguchi, H., Ogawa, Y. and Horiuchi, K., 1988, Occurrence, chemistry and age of an andesitic dike in the Hayama Group, Miura Peninsula. *Kazan (Jour. Volc. Soc. Japan)*, 33, 59-66. (in Japanese with English abstract)
- Tazaki, K., 1975, Chromian spinels in picrite basalt from Mineoka Tectonic Belt, Boso Peninsula, central Japan. *Jour. Geol. Soc. Japan*, 81(7), 399-406. (in Japanese with English abstract)
- Tazaki, K. and Inomata, M., 1980, Picrite basalts and tholeiitic basalts from Mineoka tectonic belt, Central Japan. *Jour. Geol. Soc. Japan*, 86(10), 653-671. (in Japanese with English abstract)

- Teraoka, Y., Suzuki, M., Hayashi, T. and Kawakami, K., 1997, Detrital Garnets from Paleozoic and Mesozoic Sandstones in the Onogawa Area, East Kyushu, Southwest Japan. *Bulletin of the Faculty of School Education, Hiroshima Univ., Part II*, 19, 87-101. (in Japanese with English abstract)
- Teraoka, Y., Suzuki, M. and Kawakami, K., 1998, Provenance of Cretaceous and Paleogene sediments in the Median Zone of Southwest Japan. *Bull. Geol. Surv. Japan*, 49(8), 395-411. (in Japanese with English abstract)
- Teraoka, Y., Okukmura, K., Suzuki, M. and Kawakami, K., 1999, Clastic sediments of the Shimanto Supergroup in Southwest Japan. *Bull. Geol. Surv. Japan*, 50(9), 559-590. (in Japanese with English abstract)
- Thu, M. K., Tokuyama, H., Murayama, M. and KH 00-5 scientific party, 2001, HINDOO cruise deep-sea channel survey in the Bay of Bengal. *Jour. Geol. Soc. Japan*, 107(11), XIX-XX.
- Tokuoka, T. and Kumon, F., 1979, The Shimanto Terrain in the Akaishi Mountainland and the Kii Peninsula —A consideration on mineral composition of sandstones—. *Mem. Nat. Sci. Mus., Tokyo*, 12, 41-54. (in Japanese with English abstract)
- Tokuyama, A., 1972, Geologic outline of the Ryusozan and Okuzure area, Shizuoka Prefecture, central Japan. *Contrib. Geol. Inst., Shizuoka Univ.*, 3, 7-13. (in Japanese)
- Tsuchi, R., 1986, *1:200,000 Geologic map of Shizuoka Prefecture*. Shizuoka Prefecture.
- Tsuchi, R. and Ibaraki, M., 1978, Late Neogene succession of molluscan fauna on the Pacific coast of southwestern Japan, with Reference to planktonic faunal sequence. *Veliger*, 21(2), 216-224.
- Uchida, T. and Arai, S., 1978, Petrology of ultramafic rocks from the Boso Peninsula and the Miura Peninsula. *Jour. Geol. Soc. Japan*, 84(10), 561-570.
- Uddin, A. and Lundberg, N., 1998a, Unroofing history of the eastern Himalaya and the Indo-Burman ranges: heavy-mineral study of Cenozoic sediments from the Bengal Basin, Bangladesh. *Jour. Sed. Res.*, 68(3), 465-472.
- Uddin, A. and Lundberg, N., 1998b, Cenozoic history of the Himalayan-Bengal system: Sand composition in the Bengal basin, Bangladesh. *Geol. Soc. America Bull.*, 110(4), 497-511.
- Ujiie, H., 1958, Geological structure of the Sagara-Kakegawa sedimentary basin, *Proceedings for the Symposium Concerning with the Japanese Neogene Sponsored by the Geol. Soc. Japan*, 1-7. (in Japanese)
- Ujiie, H., 1962, Geology of Sagara-Kakegawa sedimentary basin in central Japan. *Sci. Rep., Tokyo Kyoiku Daigaku*, 8, 123-188.

- Ujiie, H. and Inoue, Y., 1980, Geology of the Lower Miocene Mikasa Group, Shizuoka Prefecture, central Japan. *Mem. Nat. Sci. Mus., Tokyo*, 13, 9-14. (in Japanese with English abstract)
- Wang, C., Liu, Z. and Hébert, R., 2000, The Yarlung-Zangbo paleo-ophiolite, southern Tibet: implications for the dynamic evolution of the Yarlung-Zangbo Suture Zone. *Jour. Asian Earth Sci.*, 18, 651-661.
- Watanabe, Y., 1988, Geology of the Kurami and Yui districts in central Shizuoka: the southwestern extension of the Setogawa Tarrane. *Jour. Geol. Soc. Japan*, 94(3), 207-219. (in Japanese with English abstract)
- Watanabe, Y. and Iijima, A., 1983, Miocene Takakusayama seamount of the Setogawa Terrain in Shizuoka, central Japan. *Jour. Fac. Sci., Univ. Tokyo, Sec. II*, 20, 425-441.
- Watanabe, Y. and Iijima, A., 1990, Evolution of the Tertiary Setogawa-Kobotoke Mineoka forearc basin in central Japan with emphasis on the lower Miocene terrigenous turbidite fills. *Jour. Fac. Sci., Univ. Tokyo, Sec. II*, 22, 53-88.
- Watanabe, M. and Takahashi, M., 2000, Diatom biostratigraphy of the middle Miocene marine sequence of the Kawadani section in the Kamogawa area, Boso Peninsula, central Japan. *Jour. Geol. Soc. Japan*, 106(7), 489-500. (in Japanese with English abstract)
- Yabe, H., 1918, Itoigawa-Shizuoka Tectonic Line. *The Present Science (Gendai-no Kagaku)*, 6(3), 147-150. (in Japanese)
- Yagi, N., Taketani, Y. and Hisada, K., 1996, Radiolarian fossils from Kuchisakamoto in the Setogawa Belt, central Japan. *Bull. Agric. Forest. Res. Cent., Univ. Tsukuba*, 12, 149-158. (in Japanese with English abstract)
- Yokoyama, K., Amano, K., Taira, A. and Saito, Y., 1990, Mineralogy of silts from the Bengal Fan. *Proceedings of the Ocean Drilling Program, Scientific Results*, 116, 59-73.
- Yoshida, Y., 1974, Discovery of foraminifers in the Mineoka Hills, Chiba Prefecture. *Chishitsu News (Geology Monthly)*, no. 233, 30-36. (in Japanese)