

## ACKNOWLEDGEMENTS

I would like to express my deeply grateful and sincere acknowledgements to Associate Professor Ken-ichiro Hisada, for his invaluable supervisions, suggestions, encouragement and contributions, especially his kind-heartedness to make this research well achieved. Greatly sincere acknowledgements to Professor Kenshiro Ogasawara, Professor Katsuo Sashida and Associate Professor Toshiaki Tsuogae of the University of Tsukuba for their helpful suggestions. Deeply sincere gratitude is extended to Associate Professor Punya Charusiri of the Chulalongkorn University, Professor Shoji Arai of Kanazawa University, and Associate Professor Katsumi Ueno, who made the available time to give me the invaluable helps and suggestions.

I am highly indebted to Dr. Yoshihito Kamata of the Yamaguchi University, Dr. Hidetoshi Hara of the Geological Survey of Japan, Dr. Pol Chaodumrong and Senior Geologist Sirot Salyapongse of the Department of Mineral Resources (Thailand), and Dr. Titima Charoentitirat of the Chulalongkorn University, for their generous help and valuable comments. The help and friendship of Dr. Megumi Ichise, Mr. Koichi Okuzawa, and Mr. Tianpan Ampaiwan of the University of Tsukuba are greatly appreciated. Thanks are also made to Mr. Weerasak Lunwongsa and Mr. Viriya Danphaiboonphon of the Chulalongkorn University for their assistance.

I would specially like to express my sincere thanks to the Institute of the Promotion of Teaching Science and Technology (IPST) of Thailand for the scholarship of the Development and Promotion of Science and Technology Talents (DPST) during my education in the University of Tsukuba.

Above all, I thank my parent and family for their love and greatest support to get through all difficult and hard research. I would like to dedicate this thesis to all members of my family.