

LIST OF FIGURES

Figure		Page
1	Distribution of principal continental terranes and sutures of East and SE Asia (Metcalfe, 1998).	2
2	Geotectonic subdivisions of Northern Thailand (Ueno, 1999).	5
3	Location of the Sa Kaeo-Chanthaburi area, eastern Thailand (inserted block) with the distribution of Permian (orange color) and Triassic (purple color) rocks of marine sedimentary deposits along the western margin of the Khorat Plateau.	6
4	Paleogeography of West Gondwana and East Gondwana (Rogers and Santosh, 2003).	13
5	Geologic outline of the Sa Kaeo-Chanthaburi suture zone showing the Chanthaburi chert-clastic sequence and the Thung Kabin <i>mélange</i> (Hada et al., 1997, 1999).	15
6	Geological map of the eastern Thailand (Chaodumrong, 1992b).	17
7	Distribution of granite belts in Thailand (Charusiri et al., 1993).	20
8	Rock distribution in the Sa Kaeo-Chanthaburi accretionary complex showing characteristics of blocks in <i>mélanges</i> .	22

Figure		Page
9	Geological map of the Sa Kaeo-Chanthaburi accretionary complex.	23
10	Outcrop sketch of <i>mélange</i> in the Khao Hleam units displaying a diverse assemblage of rocks of various sizes with the chaotic structures.	27
11	Outcrop sketch of basalts along the Highway 317 (km 90) in the Khao Hleam unit showing the typical pillow structure. Field note in sketch is a scale.	28
12	Outcrop sketch of dark brown basalts along the Highway 317 (km 90+700) in the Khao Hleam unit showing the characteristics of pillow structures with green inter-pillows. Hammer in sketch is a scale.	29
13	Outcrop sketch of basaltic pillow lavas associated with fault contacts at Wat Khao Hleam in the Khao Hleam unit.	30
14	Outcrop sketch of basaltic pillow lavas intruded by black porphyrite or dolerite dikes at the same location as Fig. 13. Hammer in sketch is a scale.	31
15	Outcrop sketch of tectonically deformed basalt rocks associated with reddish brown chert floats. Small blocks of greenish gray chert and sandstone are found in the deformed basalts.	32
16	Outcrop sketch of volcanoclastic rocks associated with pillow lavas (Figs. 11 and 12) showing clasts of amygdaloidal basalts and porphyritic basalts. Some fusulinids are found in matrix part of volcanoclastic rock.	33

Figure		Page
17	Outcrop sketch of reddish brown radiolarian-bearing bedded cherts along the Highway 3448 (km 6+850) showing large size of a block in <i>mélange</i> of the Khao Hleam unit.	35
18	Outcrop sketch of large blocks of reddish brown chert in shale matrix near the radiolarian location in the Khao Hleam unit. Hammer in sketch is a scale.	36
19	Outcrop sketch of reddish brown chert, basalt and limestone near the Highway 317 (km 91+600) in the Khao Hleam unit showing their relationship among each other. Tree (6 m high) in sketch is a scale.	37
20	Outcrop sketch of <i>mélange</i> at type locality of the Ban Nong Bon unit showing the distributions of sandstone, gray chert, limestone and conglomerate blocks in shale matrix.	38
21	Outcrop sketch of <i>mélange</i> at the same location as Fig. 20 showing alignment of blocks of sandstones and limestones.	39
22	Outcrop sketch of close-up on surface of <i>mélange</i> rock at the same location as Fig. 20 displaying the major foliation and slip sense of chert and sandstone blocks.	40
23	Outcrop sketch of <i>mélange</i> along the Highway 3424 (km 3+800) in the Ban Nong Bon unit showing various sizes of sandstone blocks in shale matrix. Hammer in sketch is a scale.	42

Figure		Page
24	Outcrop sketch of duplex developed in sandstone beds of coherent turbidite sequence in the Ban Nong Bon unit along the Highway 3448. Hammer in sketch is a scale.	43
25	Outcrop sketch of the turbidite sequence in the Ban Nong Bon unit along the Highway 317 (km 74+390) showing alternation of sandstone and shale. It is covered by Quaternary sediments. Hammer in sketch is a scale.	44
26	Outcrop sketch of the conglomerate and sandstone of the coherent succession in the Ban Nong Bon unit. Paleocurrent is measured from the alignment of clast imbrication and is from southeast.	45
27	Outcrop sketch of large “loose” block of conglomerate on the top of the Khao Sa Taeng hill showing various kinds of well-rounded rock clasts in sand matrix. Fine to medium grained granite are also found.	46
28	Outcrop sketch of basalts in the Soi Dao unit showing the typical pillow structure. Arrows indicate top from pillow shape. Hammer in sketch is a scale.	48
29	Outcrop sketch of <i>mélange</i> of the Soi Dao unit displaying <i>mélange</i> foliation and their tectonic blocks of greenish gray cherts, basalts and limestones.	49
30	Outcrop sketch of close-up chert blocks in the same place as Fig. 29, showing various sizes of blocks in matrix.	50
31	Sketch of a “loose” block of basalt accompanied with limestone in the <i>mélange</i> at the same location as Fig. 29.	51
32	Outcrop sketch of <i>mélange</i> in the Soi Dao unit showing blocks of cherts and sandstones in shale matrix.	52

Figure		Page
33	Outcrop sketch of radiolarian-bearing gray chert in the Soi Dao unit along the Highway 3210 (~ km 6) showing a well-developed bedding.	53
34	Outcrop sketch of alternation of sandstone and shale along the Highway 317 (km 29+880) showing the characteristics of turbiditic sequence of the Pong Nam Ron Formation. Hammer in sketch is a scale.	56
35	Outcrop sketch of overturned bed of turbiditic sandstone in the Pong Nam Ron Formation displaying the groove casts on sole of bed. Hammer in sketch is a scale.	57
36	Outcrop sketch of the overturned bed of turbiditic sandstone of the Pong Nam Ron Formation showing the flute casts on sole of bed. Measured paleocurrent from these flute casts is from the south.	58
37	Diagram of sandstone classification (after Folk, 1974).	61
38	Standard triangular diagrams of Dickinson et al. (1983).	63
39	The plots of turbiditic sandstones of the Pong Nam Ron Formation on the triangular QtFL diagram of Dickinson (1985).	67
40	The plots of turbiditic sandstones of the Pong Nam Ron Formation on the triangular QmFLt diagram of Dickinson (1985).	68
41	Roundness and sphericity classes of the AGI (American Geological Institute) Data Sheets (Powers, 1989).	70
42	Clast population in conglomerate block in mélangé of the Ban Nong Bon unit.	72

Figure		Page
43	End member compositions in the spinel group as represented in a spinel prism (Klein and Hurlbut, 1993).	78
44	Relationships between Cr# and TiO ₂ content of detrital chromian spinels from the Pong Nam Ron Formation in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	105
45	Relationships between Cr# and TiO ₂ content of detrital chromian spinels from the sandstone and conglomerate blocks in mélangé in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	106
46	Relationships between Cr# and TiO ₂ content of detrital chromian spinels from the volcanoclastic rocks in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	108
47	Relationships between $Fe^{3+}/(Cr+Al+Fe^{3+})$ and TiO ₂ content of detrital chromian spinels from the Pong Nam Ron Formation in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	109
48	Relationships between $Fe^{3+}/(Cr+Al+Fe^{3+})$ and TiO ₂ content of detrital chromian spinels from the sandstone and conglomerate blocks in mélangé in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	110

Figure		Page
49	Relationships between $Fe^{3+}/(Cr+Al+Fe^{3+})$ and TiO_2 content of detrital chromian spinels from the volcanoclastic rocks in comparison with fields of MORB, intraplate basalt and arc magmas proposed by Arai (1992).	111
50	Tectonic model of the western margin of the Indochina around the Saraburi and Phetchabun areas during the Middle Permian to Late Permian.	116
51	Tectonic model of the western margin of the Indochina around the Sa Kaeo-Chanthaburi area during the Latest Permian to Middle Triassic.	119

LIST OF TABLE

Table		Page
1	Tectono-stratigraphic units in the Sa Kaeo-Chanthaburi accretionary complex, eastern Thailand	24