

List of Tables

2-1	Recovery test of thiophenes by the analytical method.	30
2-2	Thiophenes in Neogene sediments of the Shinjo basin and their major mass numbers (<i>m/z</i>).	34
2-3	Relative abundances of C ₂₅ HBI thiophenes and C ₂₅ HBI alkane, and bulk sulfur contents in Neogene sediments of the Shinjo basin.	36
2-4	Concentrations and relative ones of polynuclear aromatic thiophenes in Neogene sediments of the Shinjo basin.	44
3-1	Thiophenes in the K/T boundary sediments at Kawaruppu and their major mass numbers (<i>m/z</i>).	67
3-2	Relative abundances of C ₂₅ HBI thiophene and C ₂₅ HBI alkane in the K/T boundary sediments at Kawaruppu.	69
3-3	Concentrations and relative ones of polynuclear aromatic thiophenes in the K/T boundary sediments at Kawaruppu.	73
4-1	Concentrations of polynuclear aromatic thiophenes in the Murchison carbonaceous chondrite.	98

List of Figures

1-1	Elemental abundances of carbonaceous chondrites (C1) and the solar photosphere.	7
2-1	Stratigraphic and lithologic sections of the Shinjo sedimentary rocks.	25
2-2	Depth profiles of total organic carbon, extractable organic compounds and stable carbon isotopic compositions in Neogene sediments of the Shinjo basin.	26
2-3	Analytical procedure for thiophenes in Neogene sediments of the Shinjo basin.	29
2-4	Mass fragmentograms of alkyl thiophenes in the Shinjo 83109 sample.	32
2-5	Mass spectra of C ₂₀ isoprenoid (a–c) and C ₂₅ HBI (d–f) thiophenes from the Shinjo 83109 sample.	33
2-6	Depth profiles of C ₂₅ HBI thiophenes, C ₂₅ HBI alkane and bulk sulfur contents in Neogene sediments of the Shinjo basin.	37
2-7	Postulated diagenetic pathways of C ₂₅ HBI carbon skeletons.	39
2-8	Compositions of C ₂₅ HBI thiophene isomers in Neogene sediments of the Shinjo basin.	41
2-9	Mass fragmentograms of polynuclear aromatic thiophenes in the Shinjo 90102 sample.	43
2-10	Depth profiles of polynuclear aromatic thiophenes, total organic carbon (TOC) and total <i>n</i> -alkanes contents in Neogene sediments of the Shinjo basin.	45
2-11	Postulated diagenetic pathways of (a) alkyl to benzothiophenes starting from unsaturated phytanols and of (b) 2-2'-biphenol to dibenzothiophene based on the heating experiment.	47

2-12	Time-course plots of dibenzothiophene produced by the heating experiment using 2-2'-biphenol with (a) FeS and (b) elemental sulfur at 200 °C.	48
3-1	Location of the Kawaruppu section, and the lithology and sampling points of the K/T boundary in the Kawaruppu.	61
3-2	Depth profiles of total organic carbon (TOC) contents, and total <i>n</i> -alkanes and polycyclic aromatic hydrocarbons (PAHs) concentrations in the K/T boundary sediments at Kawaruppu.	62
3-3	Mass fragmentograms of C ₂₅ HBI thiophene recovered from the sediment at the -395 cm horizon of the K/T boundary sediments at Kawaruppu.	65
3-4	Mass spectra of (a) C ₂₅ HBI thiophene and (b) C ₂₅ HBI alkane from the sediment at the -395 cm horizon of the K/T boundary sediments at Kawaruppu.	66
3-5	Depth profiles of C ₂₅ HBI thiophene, C ₂₅ HBI alkane and <i>n</i> -alkanes (C ₂₇ to C ₂₉) in the K/T boundary sediments at Kawaruppu.	70
3-6	Mass fragmentograms of polynuclear aromatic thiophenes recovered from the sediment at the 40 cm horizon of the K/T boundary sediments at Kawaruppu.	72
3-7	Depth profiles of polynuclear aromatic thiophenes in the K/T boundary sediments at Kawaruppu.	76
3-8	Depth profiles of ratios of β- to α-alkylated dibenzothiophene, naphthalene and phenanthrene for the K/T boundary sediments at Kawaruppu.	77
3-9	Depth profiles of dibenzothiophene and total sulfide concentrations in the K/T boundary sediments at Kawaruppu.	78
4-1	Analytical procedure for polynuclear aromatic thiophenes in the Murchison carbonaceous chondrite.	91
4-2	Mass fragmentograms of polynuclear aromatic thiophenes in the Murchison carbonaceous chondrite.	94

4-3	Mass spectra of polynuclear aromatic thiophenes in the Murchison carbonaceous chondrite.	95
4-4	Structures of polynuclear aromatic thiophenes found in the Murchison carbonaceous chondrite.	97
4-5	Abundance patterns of polynuclear aromatic thiophenes in the Murchison carbonaceous chondrite, Neogene and K/T boundary sediments.	100