

REFERENCES

- Allemann, F., 1979: Time of emplacement of the Zhob valley ophiolites and Bela ophiolites, Baluchistan (Preliminary Report). In Farah, A. and Dejong, K. A. (eds.), *Geodynamics of Pakistan*, p. 242-261.
- Arenillas, I. and Molina, E., 1996: Biostratigrafia y evolucion de las asociaciones de foraminiferos planctonicos del transito Paleocene-Eocene en Alamedilla (Cordilleras Beticas). *Rev. Esp. Micropal.*, vol. 18 (in press).
- Aubry, M.-P., Berggren, W. A., Stott, L. and Sinha, A., 1996: The upper Paleocene-lower Eocene stratigraphic record and the Paleocene-Eocene boundary carbon isotope excursion: implications for geochronology. In, Knox, R. W. O. B. Corfield, R. M., and Dunay, R. E., eds., *Correlation of the Early Paleogene in Northwest Europe*. Geological Society Special Publication 101, p. 353-380.
- Bandy, O. L., 1949: Eocene and Oligocene foraminifera from little Stave Creek, Clarke County, Alabama. *Bull. Am. Paleont.*, vol. 32, p. 1-211.
- Banks, C. J. and Warburton, J., 1986: "Passive roof" duplex geometry in the frontal structure of the Kirthar and Sulaiman mountain belts, Pakistan. *Journal of Structural Geology*, vol. 8 (3, 4), p. 229-237.
- Banner, F. T. and Blow, W. H., 1959: The classification and stratigraphical distribution of the *Globigerinacea*. *Paleontology, London*, vol. 2, p. 1-27.

Banner, F. T., 1982: A classification and introduction to the *Globigerinacea*. In: Aspects of Micropaleontology (papers presented to Professor Tom Bernord), Banner, F. T. and Lord, a. R. (eds.) Allen and Unwin, London, p. 142-239.

Bannert, D. and Raza, H. A., 1992: The segmentation of Indo-Pakistan Plate. *Pakistan Journal of Hydrocarbon Research*, vol. 4(2), p. 5-12.

Beckmann, J. P., 1954: Die Foraminiferen der Oceanic Formation (Eocene-Oligocaen) von Barbados, Kl. Antillen. *Eclog. Geol. Helv.*, vol. 46(2), p. 301-412.

Beckmann, J. P., 1957: *Chiloguembelina* Loeblich and Tappan and related foraminifera from the lower Tertiary of the Trinidad, B.W.I. In. *Studies in Foraminifera, part 1-planktonic foraminifera*. U. S. Nat. Mus. Bull., Washington, D. C., vol. 215, p. 91.

Bender, F. K., 1995. Tectonics and structure. In Geology of Pakistan, Bender, F. K and Raza, H. A., (eds.) Gebruder Borntraeger, Berlin, 414 p.

Berggren, W. A. and Aubry, M.-P., 1996: A late Paleocene-early Eocene NW European and North Sea magnetobiochronological correlation network. In, Knox, R. W. O. B., Corfield, R. M. and Dunay, R. E., eds., *Correlation of the Early Paleocene in Northwest Europe*. Geological Society Special Publication 101, p. 309-352.

Berggren, W. A. and Miller, K. G., 1988: Paleogene tropical planktonic foraminiferal biostratigraphy and magnetobiochronology. *Micropaleontology*, vol. 34(4), p. 362-380.

Berggren, W. A. and Olsson, R. K., 1967: Taxonomy and phylogeny: *In*. Origin and development of the foraminiferal genus *Pseudohastigerina* Banner and Blow, 1979, Berggren, W. A., Olsson, R. K. and Rayment, R. A. (eds.). p. 265-281.

Berggren, W. A. and Van Couvering, J. A., 1974: The late Neogene biostratigraphy, geochronology and paleoclimatology of the last 15 million years in marine and continental sequences. *Palaeogeography Palaeoclimatology Palaeoecology*, vol. 16, p. 1-215.

Berggren, W. A., 1960: Some planktonic foraminifera from the lower Eocene (Ypresian) of Denmark and northwestern Germany. *Stockholm Contrib. Geol.*, vol. 5, p. 41-108.

Berggren, W. A., 1969: Rates of evolution in some Cenozoic planktonic Foraminifera. *Micropaleontology*, vol. 15, p. 351-365.

Berggren, W. A., 1992: Paleogene planktonic foraminifera Magnetobiostratigraphy of the Southern Kergulen Plateau (Sites 747-749). *In* S. W. Wise, Jr., R. Schlich et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, College Station, Texas, vol. 120, p. 551-569.

Berggren, W. A., Kent, D. V., Swisher, C. C. and Aubry, M. -P., 1995: A revised Cenozoic geochronology and chronostratigraphy. *In* Berggren, W. A., Kent, D. V., Aubry, M. -P. and Hardenbol, J., eds., *Geochronology, Time Scales and Global Stratigraphic Correlation. SEPM Special Publication*, no. 54, p. 129-212.

Berggren, W.A. and Norris, N.D., 1997: Biostratigraphy, phylogeny and systematics of Paleocene trochospiral planktic foraminifera. *Micropaleontology*, vol. 43, supplement 1, p. 1-116.

Blandford, W. T., 1876: Note on the geology of the neighborhood of Lyngan and Runneekot, northwestern Kotrree in Sind. *Geol. Surv. India, Mem.*, vol. 20, 229-237.

Blandford, W. T., 1883: Geological note on the hills in the neighborhood of Sind and Punjab frontier between Quetta and Dera Ghazi Khan. *Geol. Surv. India, Mem.*, vol. 20, 229-237.

Blow, W. H. and Banner, F. T., 1962: The Mid-Tertiary (Upper Eocene to Aquantanian) Globigerinaceae. In. F. E Eames et al., *Fundamentals of Mid-Tertiary Stratigraphical Correlation*, Cambridge University Press, p. 61-151.

Blow, W.H., 1969: Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. *Proceedings of the First International Conference on Planktonic Microfossils*, Geneva, 1967, 1, p. 199-422.

Blow, W.H., 1979: The Cainozoic Globigerinidea, 3 vols., 1452 p., 254 pls. E.J. Brill, Leiden.

Bolli, H. M. and Krasheninnikov, V. A., 1977: Problem in Paleogene and Neogene correlation based on planktonic foraminifera. *Micropaleontology*, vol. 23, p. 436-452.

Bolli, H. M., 1954: Note on *Globigerina concinna* Reuss 1850. *Contri. Cushman Foundation Foramin. Res.*, vol. 5, p. 1-3.

- Bolli, H. M., 1957a: Planktonic foraminifera from the Eocene Navet and San Fernando Formations of the Trinidad, B.W.I. *Bull. U.S. Nat. Mus., Washington*, vol. 215, p. 155-172.
- Bolli, H. M., 1957b: The genera *Globigerina* and *Globorotalia* in the Paleocene-lower Eocene Lizard Spring Formation of Trinidad, B.W.I. *Bull. U.S. Nat. Mus., Washington*, vol. 215, p. 61-81.
- Bolli, H. M., 1957c: Planktonic foraminifera from the Oligocene-Miocene Cipero and Lengua Formations of the Trinidad, B.W.I. *Bull. U.S. Nat. Mus., Washington*, vol. 215, p. 97-123.
- Bolli, H. M., 1966: Zonation of Cretaceous to Pliocene marine sediments based on planktonic foraminifera. *Boletino Informativo Asociación Venezolana de Geología, Mineraria y Petróleo*, vol. 9, p. 3-33.
- Bolli, H. M., Loeblich, Jr. A. R. and Tappan, H., 1957: Planktonic foraminiferal families Hantkeninidae, Orbulinidae, Globorotaliidae, Globotruncanidae. *Bull. U.S. Nat. Mus., Washington*, vol. 215, p. 3-50.
- Bolli, H.M. and Cita, M.B., 1960: Globigerine e Globorotalie del Paleocene de Paderno d'Ada (Italia). *Revista Italiana Paleontología*, vol. 66(3), p. 361-402.
- Bolli, H.M. and Saunders, J. B., 1985: Oligocene to Holocene low latitude planktic foraminifera. In: *Plankton Stratigraphy*, Bolli, H. M., Saunders, J. B. and Nielsen, K. P., (eds). Cambridge University Press, P. 186-187.

Bronnimann, P. and Bermudez, P. J., 1953: *Truncorotaloides*, a new foraminiferal genus from the Eocene of Trinidad, B.W.I. *Journal of Paleontology*, vol. 27, p. 817-820.

Bronnimann, P., 1952: Trinidad Paleocene and Lower Eocene Globigerinidae. *Bulletins of American Paleontology*, vol. 34(143), p. 1-34.

Brotzen, F. and Pozaryska, K., 1961: Foraminifères du Paleocene et de l'Eocene inférieur en Pologne septentrionale; remarques paleogeographiques. *Revue de Micropaleontologie*, vol. 4, p. 155-166.

Butt, A. A. 1991: *Ranikothalia sindensis* Zone in Late Paleocene biostratigraphy. *Micropaleontology*, vol. 37(1), p. 77-85.

Bykova, 1953: Foraminifery Suzakskogo Yarusa Tadzhikskoy Depresii. *Trudy Vsesoyuznogo Naftyanogo Nauchno-issledovate' skogo Geologorazvedochnogo Instituta (Vnigri) nov-ser.*, vol. 69, p. 5-105 (Microfauna SSSR, Sbornik 6).

Canudo, J. I., Keller, G., Molina, E. and Ortiz, N., 1995: Plantic foraminiferal faunal turnover and ^{13}C isotope across the Paleocene-Eocene transition at Caravaca and Zumaya, Spain. *Paleogeography; Paleoclimatology; Paleoecology*. vol. 114, p. 75-100.

Carpenter, W. B., Parker, W. K. and Jones, T. R., 1862: Introduction to the study of the Foraminifera. *London, Ray Society*.

Carpenter, W. B., Parker, W. K. and Jones, T. R., 1876: Introduction to the study of the Foraminifera. *London, Ray Society*.

Cifelli, R. and Belford, D.J., 1977: The types of several species of Tertiary planktonic foraminifera in the collections of the U.S. National Museum of Natural History. *Journal of Foraminiferal Research*, vol.7(2): 100-105, pl. 1.

Cole, W. S., 1927: A foraminiferal fauna from the Guayabal Formation in Mexico. *Am. Paleontol. Ithaca, New York*, vol. 14, p. 1-46.

Cole, W. S., 1928: A foraminiferal fauna from the Chapapote Formation in Mexico. *Bull. Am. Paleontol.*, vol. 14, p. 3-32.

Colom, G., 1954: Estudio de biozonas con foraminíferos del Terciario de Alicante. *Boletin del Instituto Geológico de España. Y Minero*, vol. 66, p. 1-279.

Cordey, W. G., 1968: Morphology and phylogeny of *Orbulinoides beckmanni* (Saito, 1962). *Paleontology*, vol. 11, p. 371-375.

Corlfield, R. M. and Shackleton, N. J., 1988: Productivity change as control of planktonic foraminiferal evolution after the Cretaceous/Tertiary boundary. *Historical Biology*, vol. 1, p. 323-343.

Coward, M. P., Jan, M. q., Rex, D., Tarney, J., Thirwal, M. and Windley, B. W., 1982: Structural evolution of a crustal section in the western Himalaya. *Nature*, vol. 295, 22-24.

Cushman, J. A. and Renz, H.H., 1942: Eocene, Midway, foraminifera from Sodado Rock, Trinidad. *Contributions from the Cushman Foundation for Foraminiferal Research*, vol. 18, p. 1-20.

- Cushman, J. A. and Renz, H.H., 1942: Eocene, Midway, foraminifera from Sodado Rock, Trinidad. *Contributions from the Cushman Foundation for Foraminiferal Research*, vol. 18, p. 1-20.
- Cushman, J. A. and Bermudez, P. J., 1937: Further new species of foraminifera from the Eocene of Cuba. *Contrib Cushman Lab. Foramin. Res.*, vol. 13, 1-29.
- Cushman, J. A. and Bermudez, P. J., 1949: Some Cuban species of *Globorotalia*. *Contributions from the Cushman for Foundation Foraminiferal Research*, vol. 25, p. 26-45.
- Cushman, J. A. and Hana, G. D., 1927: Foraminifera from the Eocene near Coalinga, California. *Proceedings of the California Academy of Science, Series 4*, vol. 8, p. 205-229.
- Cushman, J. A. and Hana, G. D., 1927: Foraminifera from the Eocene near San Diego, California. *Transactions of the San Diego Society of Natural History*, vol. 5, p. 45-64.
- Cushman, J. A. and Jarvis, P. W., 1929: New foraminifera from Trinidad. *Contrib. Cushman Lab. Foramin. Res.*, vol. 5, p. 6-17.
- Cushman, J. A. and Jarvis, P. W., 1933: Post-Cretaceous occurrence of *Gumbilina* with a description of a new species. *Contri. Cushman Lab. Foram. Res.*, Sharon, Mass., U.S.A., vol. 9, p. 69.
- Cushman, J. A. and Ponton, G. M., 1932: An Eocene foraminiferal fauna of Wilcox age from Alabama. *Contributions from the Cushman Foundation for Foraminiferal Research*, vol. 8, p. 51-72.

- Cushman, J. A., 1925a: A new genus of the Eocene Foraminifera. *Proc. U.S. Nat. Mus. Washington*, vol. 66, P. 1-4.
- Cushman, J. A., 1925b: Some new foraminifera from the Velasco Shale of Mexico. *Contribution Cushman Foundation Foraminiferal Research*, vol. 1, p. 18-23.
- Cushman, J. A., 1925c: New foraminifera from the Upper Eocene of Mexico. *Contrib. Cushman lab. Foramin. Res.*, vol. 1, 18-23.
- Cushman, J. A., 1925d: An Eocene fauna from the Moctezuma River, Mexico. *Bull. Amer. Assoc. Petr. Geol.* vol. 9(2), p. 298-303.
- Cushman, J. A., 1927a: An outline of re-classification of the Foraminifera. *Contrib. Cushman lab. Foramin. Res.*, vol. 3, 1-05.
- Cushman, J. A., 1927b: New and interesting foraminifera from Mexico and Texas. *Contrib. Cushman lab. Foramin. Res.*, vol. 3, 111-119.
- Davidzon, R. M., 1976: Novyy paleogenovyy rod planktonnykh foraminifer {A new planktonic foraminiferal genus}. Trudy, vsesoyuznyi Nauchno-Issledovatel'skii Geologorazvedochnyi Neftyanoi Institut (VNIGNI), Tadzhikskoe Otdelenie, vol. 183, p. 197-198 [In Russian]
- Davies, L. M., 1939: Geographical changes in the northwestern India during late Cretaceous and early Tertiary times. *Proc. Pac. Sci. Conf. 6th*, vol. 6 (2), p. 483-501.
- Davis, D. M., 1994: Changing mechanical response during continental collision: active examples from the foreland thrust belts of Pakistan. *Journal of Structural Geology*, vol. 16(1), p. 21-34.

- Dewey, J. F., Cande, S. and Pitman, W. C. III., 1989: Tectonic evolution of India/Eurasia collision zone. *Eclog. Geol. Helv.*, vol. 82, p. 717-734.
- Dorreen, J. M., 1974: The Western Gaj River section, Pakistan and Cretaceous-Tertiary boundary. *Micropaleontology*, vol. 20(2), p. 178-193.
- Dunham, R. J., 1962: Classification of carbonate rocks according to the depositional texture. In, *American Association of Petroleum Geology, Memoir*, no. 1, p. 108-121.
- Eames, F. E., 1952a: A contribution to the study of the Eocene in the West Pakistan and Western India. Description of faunas of certain standard sections and their bearing on the classification and correlation of the Eocene in Western Pakistan and Western India. *Ibid.*, vol. 107, p. 170-200.
- Eames, F. E., 1952b: A contribution to the study of the Eocene in the West Pakistan and Western India. The geology of the standard sections in the Western Punjab and in Kohat district. *Quaternary Journal of the Geological Society, London*, vol. 107, p. 159 -171.
- El Khoudary, R. H., 1977: *Truncorotaloides libyaensis*, a new planktonic foraminifer from Jabal Al Akhdar (Libya). *Rev. Esp. Micropaleontol.* vol. 9, p. 327-336.
- Farah , A., Abbas, G., Dejong, K. A. and Lawrence, R. D., 1984: Evolution of the lithosphere in Pakistan. *Tectonophysics*, vol. 105, p. 207-227.
- Finlay, H. J., 1939a: New Zealand foraminifera: Key species in stratigraphy. No 2. *Trans. Roy. Soc. New Zealand, Wellington*, vol. 69, p. 89-128.

- Finlay, H. J., 1939b: New Zealand foraminifera: Key species in stratigraphy. No 3. *Trans. Roy. Soc. New Zealand*, Wellington, vol. 69 (3), p.327.
- Folk, R. L., 1962: Spectal subdivision of limestone types. In. Ham, W. E. (eds.) classification of carbonate rocks. *American Association of Petroleum Geology, Memoir*, vol. 1, p. 62-84.
- Gansser, A., 1964: Geology of the Himalayas. *Wiley Interscience, London*, 289 p.
- Gansser, A., 1979: Reconnaissance visit to the ophiolites in Baluchistan and Himalaya. In Farah and K. A. Dejong (eds.), *Geodynamics of Pakistan, Geological Survey of Pakistan, Quetta*, p. 193-213.
- Gee, E. R., 1989. Overview of the geology and structure of the Salt Range, with observations on related areas of northern Pakistan. In Malinconico, L. L. and Lillie, R. J. (eds): Tectonics of the Western Himalayas. *Geological Society of America Special Publication, Tulsa*, 232, p. 95-112.
- Gibson, T. G., 1989: Planktonic benthonic foraminiferal ratios: Modern patterns and Tertiary applicability. *Marine Micropaleontology*, vol. 15, p. 29-52.
- Glaessner, M. F., 1937: Planktonoforaminiferen aus der Kreide und dem Eozaen und ihre stratigraphische Bedeutung. *Studies in Micropaleontology, Publications of the Laboratory of Paleontology, Moscow University*. vol. 1, p. 27-46.
- Gohrbandt, K. H. A., 1967: Some new planktonic foraminiferal species from Austrian Eocene, *Micropaleontology*, v. 13, p. 319-326.

Greisbach, C. L., 1881: Report on geology of the section between Bolan Pass and Baluchistan and Girishk in southern Afghanistan: *India Geol. Surv., Mem.*, vol. 18, p. 1-60.

Grimsdale, T. F. and Van Morkhoven, F. P. C. M., 1955: The ratio between pelagic and benthonic foraminifera as a means of estimating depth of deposition of sedimentary rocks. *Proceedings of the World Petroleum Congress*, 4 (Room) Section 1/D4, p. 473-491.

Haque, A. F. M., 1956: The smaller foraminifera of Ranikot and the Laki of Namal George, Salt Range. *Memoires of the Geological Survey of Pakistan, Paleontologica Pakistanica*, vol.1, p. 1-300.

Haque, A. F. M., 1959a: The smaller foraminifera of the Meting Limestone (Lower Eocene) Meting, Hyderabad Division, West Pakistan. *Memoires of the Geological Survey of Pakistan, Paleontologica, Pakistanica*, 2 (part 1), p. 1-43.

Haque, A. F. M., 1959b: Some Middle to Late Eocene smaller foraminifera of the Sore Range, Quetta District, West Pakistan. *Memoires of the Geological Survey of Pakistan, Paleontologica Pakistanica*, vol. 2 (part 2), p.1-79.

Hemphil, W. R. and Kidwai, A. H., 1973: Stratigraphy of Banu and Dera Ismail Khan areas, Oakistan. *U.S. Geol. Surv. Prof. Pap.*, 716B, 36p.

Holiday, D. W., 1970: The petrography of secondary gypsum rock: a review, *J. Sedim. Petrol.*, vol. 40, p. 734-744.

- Howe, H. V. and Wallace, W. E., 1932: Foraminifera of the Jackson Eocene at Danville Landing on the Ouchita, Catahoula Parish, Louisiana. *Bull. Louisiana Dept. Conservation Geol.*, vol. 2, p. 18-79.
- Huber, B. T., 1991: Paleocene and early Neogene planktonic foraminifer biostratigraphy of Sites 738 and 744, Kerguelen Plateau (Southern Indian Ocean). In J. Barron, B. Larsen et al., *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 114, p. 281-298.
- Humayon, M., Lillie, R. J. and Lawrence, R. D., 1991: Structural interpretations of the eastern Sulaiman fold belt and foredeep, Pakistan. *Tectonis*, vol. 10, no. 2, p. 299-324.
- Hunting Survey Corporation (H. S. C.) 1961: Reconnaissance geology of part of West Pakistan (Colombo plan cooperative project. *Canadian Govt., Toronto*, p. 1-550.
- Ingle, Jr, J. D., Jr. 1980: Cenozoic paleobathymetry and depositional history of selected sequences within the southern California continental borderland. *Cushman Foundation for foraminiferal Research Special Publication*, no. 19, p. 163-195.
- Jadoon, I. A. K, Lillie, R. J., Humayun, M., Lawrence, R. D., Ali, S. M. and Cheema, A., 1989: Mechanism of deformation and the nature of the crust underneath the Himalayan foreland fold-and-thrust belts in Pakistan (abstract), *Eos. Trans. AGU*, 70, 1372-1373.

Jadoon, I. A. K., 1991: The style and evolution of foreland structures: An example from Sulaiman lobe, Pakistan. *Pakistan Journal of Hydrocarbon Resources*, vol. 3, p. 1-17.

Jadoon, I. A. K., 1992: Oceanic/Continental transitional crust underneath the Sulaiman thrust lobe and evolutionary tectonic model for the Indian/Afghan collision zone. *Pakistan Journal of Hydrocarbon Resources*, vol.4 (2), p. 33-44.

Jadoon, I. A. K., Lawrence, R. D. and Khan, S. H., 1993: Mari-Bugti pop-up zone in the central Sulaiman fold belt, Pakistan. *Journal of Structural Geology*, (In press).

Jadoon, I. A. K., Lawrence, R. D. and Lillie, R. J., 1992: Balanced and retrodeformed geological cross-section from the frontal Sulaiman lobe, Pakistan: Duplex development in the thick strata along western margin of the Indian Plate, In. Thrust Tectonics edited by McClay, K., Chapman and Hall, p. 343-356.

Jamil, A., Hussain, T. and Memon, A. R., 1971: Geological map of Sinjidi '35 1/1; Khuzdar district, Baluchistan, Sheet No. 251, Scale 1: 50, 000. *Geol. Surv. Pak. Map Series, Quetta*, vol. II.

Jenkins, D. G., 1971: New Zealand Cenozoic planktonic foraminifera. *Paleontol. Bull. Geol. Surv. New Zealand*, vol. 42, p. 1-278.

Jones, R. W., 1997: Aspect of the Cenozoic stratigraphy of the Northern Sulaiman Ranges, Pakistan. *Journal of Micropaleontology*, vol. 16, p. 51-58.

- Kazmi, A. H. 1995: Sedimentary sequence. In *Geology of Pakistan*, Bender, F. K and Raza, H. A., (eds.) Gebruder Borntraeger, Berlin, 414p.
- Kazmi, A. H. and Rana, R. A., 1982: Tectonic map of Pakistan, scale 1: 2, 000,000. *Geological Survey of Pakistan, Quetta*.
- Kazmi, A. H., 1988: Stratigraphy of the Dunghan Group in Kach-Ziarat area, N. E Baluchistan. *Geol. Bull. Univ. Peshawar*, vol. 21, p. 117-130.
- Kazmin, V. G., 1991: Collision and rifting in the Tethys Ocean: geodynamic implications. *Tectonophysics*, vol. 196, p. 371-384.
- Kennett, J. P. and Stott, L. D., 1990: Proteus and Proto-Oceanus: ancestral Pleogene oceans as revealed from Antarctic stable isotopic results. In, Barker, P. F., kennett, J. P. et. al., *Proceedings of the Ocean Drilling Project Science Results College Station, TX (Ocean Drilling Program)*. vol. 113, p. 865-880.
- Kennett, J. P. and Stott, L. D., 1991: Terminal Paleocene deep -see benthic crisis: sharp deep-see warming and paleoceanographic changes in Antarctica: *Nature*, vol. 353, p. 225-229.
- Khalilov, D. M., 1956: On the pelagic foraminiferal fauna of the Paleogene of Azerbaijan, *Trudy Akad. Nauk. Azerb. S.S.R., Inst. Geol. Baku*, vol. 17, p. 234-261.
- Krasheninikove, V. A. and Hoskin, R. S., 1973: Late Cretaceous, Paleocene and Eocene planktonic foraminifera. In Heezen, B. C., McGregor, I. D., et al., *Initial Reports of the Deep Sea Drilling Project, Washington*, vol. 20, p. 105-203.

Krasheninikove, V. A., 1965a: Zonal stratigraphy of the Paleogene deposits. 21st International Geological Congress, Norden, 1960, Doklade Soviet Geologists, Problem 16, Problems of Cenozoic stratigraphy, Akademi Nauk SSSR Izd., Moscow, p. 37-61 (*in Russian*).

Krasheninikove, V. A., 1965a: Zonal stratigraphy of the Paleogene in the eastern Mediterranean. Akademi Nauk SSSR Geol. Inst. Trudy, vol. 133, P. 1-76 (*in Russian*).

Latif, M. A., 1961: The use of the pelagic foraminifera in the subdivision of the Paleocene-Eocene of the Rakhi Nala, West Pakistan. *Geol. Bull. Punjab Univ., Lahore*, vol. 1, p. 31-45.

Latif, M. A., 1964: Variation in the abundance and morphology of pelagic foraminifera in the Paleocene-Eocene of the Rakhi Nala, West Pakistan. *Geol. Bull. Punjab Univ., Lahore*, vol. 4, p. 29-109.

LeRoy, L. W., 1953: Biostratigraphy of Maqfi section, Egypt. *The Geological Society of America, Memoir* 54, p. 1-73.

Lillie, R. J., Lawrence, R. D., Humayun, M. and Jadoon, I. A. K., 1989: The Sulaiman thrust lobe of Pakistan: early stage underthrusting of the Mesozoic rifted margin of the Indian subcontinent (abstract). *Geol. Soc. Am. Abstr. Programs*, vol. 21, A318.

Loeblich, A. R., Jr. and Tappan, H., 1956: Chiloguembla a new Tertiary genus of the Heterohelicidae (Foraminifera), Journal of the Washington Academy of Sciences, vol. 46, 340p.

Loeblich, A. R., Jr. and Tappan, H., 1957: Planktonic foraminifera of Paleocene and early Eocene age from the Gulf and Atlantic Coastal Plains, U.S. National Museum, Bulletin, vol. 215, p. 173-198.

Loeblich, A. R., Jr. and Tappan, H., 1961: Stratigraphic classification of the Rhizopoda, Journal of Paleontology, vol. 35, p. 245-304.

Lu, G. and Keller, G., 1993: Climatic and oceanographic events across the Paleocene-Eocene transition in the Antarctic Indian Ocean: Inference from planktic foraminifera . Marine Micropaleontology, vol. 21, p. 101-132.

Lu, G. and Keller, G., 1995a: Planktic foraminiferal faunal turnovers in the subtropical Pacific during the late Paleocene to early Eocene. Journal of Foraminiferal Research, vol. 25(2), p. 97-116.

Lu, G. and Keller, G., 1995b: Ecological stasis and saltation: Is specimen richness change in Planktic foraminifera during the late Paleocene to early Eocene, DSDP Site 577. Paleogeogr., Palaeoclimatol., Palaeoecol., vol. 117, p. 211-227.

Lu, G., Keller, G., Adatte, T., Ortiz, N. and Molina, E., 1996: Long-term (10^5) or short-term (10^3) δC^{13} excursion near the Paleocene-Eocene transition event. Isr. Journal. Earth Sci., vol. 44, p. 185-195.

Luterbacher, H. P., 1964: Studies in some *Globorotalia* from the Paleocene and lower Eocene of central Apennines. *Eclogae Geologicae Helveticae*, vol. 57, p. 631-730.

Martin, L. T., 1943: Eocene foraminifera from the type Lodo Formation, Fresno County, California. *Stanford University Publications. Geological Sciences*, vol. 3, p. 93-125.

McGowran, B., 1968: Reclassification of early Tertiary *Globorotalia*. *Micropaleontology*, vol. 14, p. 179-198.

Metcalfe, I., 1995: Gondwana dispersion and Asian accretion. *Journal of Geology, Series B*, p. 223-266.

Miller, K. G., Janecek, T. R., Katz, M. E. and Kell, D. K., 1987: Abyssal circulation and benthic foraminiferal changes near the Paleocene/Eocene boundary. *Paleogeography*, vol. 2, p. 741-764.

Morozova, V. G., 1939: Stratigrafii verkhnego mela i paleogena embenskoio blasti po faune foraminifer [On the stratigraphy of the Upper Cretaceous and Paleogene of Emba region according to the foraminiferal faunas]. *Blulleten Moskovskogo obshchestva ispytatelei prirody. Otdel. geologicheskii*, vol. 17, no. 4-5, p. 59-86.

Nagapa, Y., 1959: Foraminiferal biostratigraphy of the Cretaceous-Eocene succession in the India-Pakistan-Burma region. *Micropaleontology*, vol. 5, p. 145 -179.

Nishi, H., Ashraf, M. T. Elewa and Ishizaki, K., 1994: Planktonic foraminiferal biostratigraphy of upper Paleocene to middle Eocene sequences in the

eastern desert area, Egypt. *Trans. Proc. Paleont. Soc. Japan*, N. S. No. 175, p. 521-552.

Noetling, F., 1903: Übergang Zwischen Kreide und Eocan in Baluchistan: *Centralb. Mineral., Geol., Paleont., Jahrb.*, vol. 4, p. 514- 523.

Nomura, R. and Brohi, I. A., 1995: Benthic foraminiferal fauna during the time of Indian-Asian contact, in southern Baluchistan, Pakistan. *Marine Micropaleontology*, vol. 24, p. 215-238.

Nuttall, W. L. W., 1930: Eocene foraminifera from Mexico. *Journal of Paleontology*, vol. 4, p. 69-101.

Oldham , R. D., 1890: Report on the geology and economic resources of the country adjoining the Sind-Pishin railway between Sharigh and spintagi, and of the country between it and Khattan. *Ibid., Rocs.*, vol. 23, p. 93-109.

Olsson, R. K., Hemleben, C., William A, B. and Huber, B. T., eds. 1999: Atlas of Paleocene planktonic foraminifera. *Smithsonian Contributions to Paleobiology*, vol. 85, p. 1-152

Olsson, R. K., Hemleben, C., William A, B. and Liu, C., 1992: Wall Texture Classification of Planktonic Foraminifera Genera in the Lower Danian. *Journal of Foraminiferal Research*, vol. 22, p. 195-213.

Ortiz, N., 1995: Differential patterns of benthic foraminiferal extinctions near the Paleocene-Eocene boundary in the North Atlantic and western Tethys. *Mar. Micropaleontol.*, vol. 26, p. 341-359.

- Pak, D. and Miller, K., 1992: Paleocene to Eocene benthic foraminiferal isotopes and assemblages: implications for deep water circulation. *Paleoceanography*, vol. 7(4), p. 405-422.
- Pardo, A., Keller, G., Molina, E. and Canudo, J. I., 1997: Planktic foraminiferal turnovers across the Paleocene-Eocene transition at DSDP Site 401, Bay of Biscay, north Atlantic. *Marine Micropaleontology*, vol. 29, p. 129-158.
- Parr, W. J., 1938: Upper Eocene foraminifera from the deep borings in King's Park, Perth, Western Australia. *Journal of Royal Society of Western Australia*, vol. 24, p. 69-101.
- Pearson, P. N., Shackleton, N. J. and Hall, M. A., 1993: The stable isotope paleoecology of middle Eocene planktonic foraminifera and multi-species integrated isotopes stratigraphy, DSDP Site 522, South Atlantic. *Journal of Foraminiferal Research*, vol. 23, p. 123-140.
- Petters, V., 1954: Tertiary and upper Cretaceous foraminifera from Columbia, S. A. *Contrib. Cushman Foundation Foramin. Res.*, vol. 5, 37-41.
- Pijpers, P. J., 1933: Geology and paleontology of Bonaire (Dutch West Indies). *Meded. Geogr. Geol. Physiogr.. Geol. Recks, Utrecht*, vol. 8, p. 1-103.
- Pinfold, E. S., 1939: The Dunghan Limestone and the Cretaceous unconformity in northwest India. *Rec. Geol. Surv. Of India*, vol. 74, p. 189-98.
- Plummer, H. J., 1926: Foraminifera of the Midway Formation of Texas. *University of Texas Bulletin*, Austin, no. 2644, p. 1-206.

- Powell, C. M. A., 1979: A speculative tectonic history of Pakistan: Some constraints from the Indian Ocean. In *Geodynamics of Pakistan* edited by A. Farah and K. A. DeJong, *Geological survey of Pakistan, Quetta*, p. 5-24.
- Primoli Silva, I. and Bolli, H. M., 1973: Late Cretaceous to Eocene planktonic foraminifera and stratigraphy of Leg 15 Sites in the Caribbean Sea. In, Edgar, N. T., Saunders, J. B., et al., *Initial Reports of Deep Sea Drilling Project*, vol. 15, p. 449-547.
- Proto Decima, F., and Bolli, H. M., 1970: Evolution and variability of *Orbulinoides beckmanni* (Saito), *Eclogae Geologicae Helvetiae*, vol. 63, p. 883-905.
- Raju, D. S. N., 1968: Eocene-Oligocene planktonic foraminiferal biostratigraphy of Cauvery Basin, South India. *Mem. Geol. Soc. India, Bangalore*, vol. 2, p. 286-299.
- Raza, H. A., Ahmad, R. and Ali, S. M., 1989. Petroleum zones of Pakistan. *Pakistan Journal of Hydrocarbon Resources, Islamabad*, vol. 2(2), p. 1-20.
- Reiss, Z., 1963: Reclassification of perforate foraminifera, *Bulletin of the Geological Survey of Israel*, vol. 35, p. 1-111.
- Saito, T., 1962: Eocene planktonic foraminifera from Hahajima (Hillsborough Island). *Trans. Proc. Paleontol. Soc. Japan*, new series, vol. 45, p. 209-205.
- Samanta, B. K., 1970: Middle Eocene planktonic foraminifera from the Lakhpat, Cutch, Western India. *Micropaleontology*, vol. 16, p. 185-215.

- Samanta, B. K., 1973: Planktonic foraminifera from the Paleocene-Eocene succession in the Rakhi Nala, Sulaiman Range, Pakistan. *Bulletin of the British Museum (Natural History) Geology*, vol. 22, p. 433-481.
- Sarwar, G. and Dejong, K. A., 1979: Arcs, oroclines, syntaxes: The curvatures of the mountain belts in Pakistan. In Farah, A. and Dejong, K. A. (eds.), *Geodynamics of Pakistan*, 341-349.
- Sarwar, G. and Dejong, K. A., 1984: Composition and origin of the Kanar melange, southern Pakistan. *Geol. Soc. Amer. Special Paper 198, Melange, their nature, origin and significance*, (Ramond, L. ed.) p. 127-137.
- Schmidt, R. and Raju, D. S. N., 1973: *Globorotalia palmerae* Cushman and Bermudez, and closely related species from the Lower Eocene, Cauvery Basin, South India. *Proc. Koninkle. Nederlandse Akad. Wetenschappen, ser. B*, vol. 76, p. 167-178.
- Searle, M. P., Windley, B. F., Coward, M. P., Cooper, D. J., Rex, A. J., Rex, D. C., Li Tingdon, Xiao Xuchang, Jan, M. Q., Thakur, V. C., and Kumar, S., 1987: The closing of Tethys and tectonics of Himalaya. *Geol. Soc. Amer. Bull., Tulsa*, vol. 98, p678-701.
- Shackleton, N. J., 1986. Paleogene stable isotope events. *Paleogeography, Paleoclimatology, Paleoecology*, vol. 57, p. 91-102.
- Shackleton, N. J., Corlfield, R. M. and Hall, M. A., 1985: Stable isotope data and ontogeny of Paleogene planktonic foraminifera. *Journal of Foraminifera, Research*, vol. 21, p. 321-336.

Shah, S. M. I., 1977: Stratigraphy of Pakistan. *Memoires of the Geological Survey of Pakistan*, vol. 12. p. 1-138.

Shah, S. M. I., 1987: Lithostratigraphic units of the Sulaiman and Kirthar Provinces, Lower Indus Basin, Pakistan. *Geol. Surv. Pak. Mem. Quetta*, vol. 17 (I-III)

Shah, S. M. I., 1990: Coal resources of Baluchistan. In Kazmi, A. H. and Siddiqui, R. A. (eds.), Significance of the coal resources of Pakistan. *Geol. Surv. Pak./ U.S. Geol. Surv.*, p. 63-93.

Smith, F. G., 1955: Planktonic foraminifera as indicators of depositional environment. *Micropaleontology*, vol. 1, p. 147-151.

Snyder, S.W. and Waters, V.J., 1985: Cenozoic planktonic foraminiferal biostratigraphy of the Goban Spur region, DSDP, Vol. 80, Washington (U.S. Govt. Printing Office): p. 439-472.

Stainforth, R. M., Lamb, J. L., Luterbacher, H., Beard, J. H. and Jeffords, R. M., 1975: Cenozoic planktonic foraminiferal zonation and characteristics of index forms. *University Kansas of Paleontological Contribution*, vol. 62, p. 1-425.

Subbotina, N. N., 1947: Foraminifera of Danian and Paleogene deposits of the northern Caucasus. *Trudy Neftianogo razvedochnogo Institituta. Novaia seria, Leningrad - Moscow*, p. 39-160.

Subbotina, N. N., 1953: Fossil foraminifera of the U.S.S.R., Globigerinidae, Hantkeninidae and Globorotaliidae. *Trudy Neftianogo razvedochnogo Institituta. Novaia seria, Leningrad - Moscow*, vol. 76, p. 1-296.

- Tahirkheli, R. A. K., Mattauer, M., Proust, R. and Tapponnier, P., 1979: The India Eurasia sutute zone in northern Pakistan: Synthesis and interpretation of recent data at Plate scale. In Farah, A. and Dejong, K. A. (eds.): Geodynamics of Pakistan, *Geological Survey of Pakistan, Quetta*, p. 125-130.
- Tainsh, H. R., Stringer, K. V. and Azad, J., 1959: Major gas fields of West Pakistan. *Amer. Assoc. Petroleum Geol., Bull.*, vol. 43, p. 2675-2700.
- Thomas, E. and Shackleton, N. J., 1996: The Paleocene-Eocene benthic foraminiferal extinction and stable isotope anomalies. In, Knox, R. W. O. B., Corfield, R. M. and Dunnay, R. E., eds., *Correlation of the Early Pleogene in Northwestern Europe*, Geological Society Special Publication 101, p. 401-441.
- Thomson, F. R. S., 1876: In. Preliminary reports to Professor Wyville Thomson, F. R. S., director of the civilian scientific staff, on work done on board the "Challenger," Murray, J., 1876 (eds.). *Proceedings of the Royal Society, London*, vol. 24, p. 471-544.
- Todd, R. and Kniker, H. T., 1952: An Eocene foraminiferal fauna from the Agua Fresca shale of Magallanes province, southernmost Chile. *Cushman Foundation for Foraminiferal Research Special Publication*, no. 1, p. 26.
- Toulmin, L.D., 1941: Eocene smaller foraminifera from the Salt Mountain limestone of Alabama. *Journal of Paleontology*, vol. 15, 567-611.

- Toumarkine, M. and Bolli, H. M., 1970: Evolution de *Globorotalia cerroazulensis* (Cole) dans l'Eocene moyen et supérieur de Possagno (Italie). *Rev. Micropaleontol.*, vol. 13, p. 131-145.
- Toumarkine, M. and Louterbacher, H., 1985: Paleocene and Eocene planktic foraminifera. In, Bolli, H. M., Saunders, J.B. and Perch-Nelson, K., eds., *Plankton Stratigraphy*, p. 87-154. Cambridge University Press, Cambridge.
- Treloar, P. J. and Searle, M. P. (eds.), 1993: Himalayan Tectonics, *Geol. Soc. London, Special Publication*, vol. 74, 630p.
- Van Ejden, A. J. M. and Smit, J., 1991: Eastern Indian Ocean Cretaceous and Paleocene quantitative biostratigraphy. In Weissel, J., Pierce, J., Taylor, E., Alt, J., et al. 1991 (eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 121, p. 77-123.
- Vredenburg, E. W., 1906: A geological sketch of the Baluchistan desert and part of Eastern Persia: *India, Geol. Surv. Mem.* vol. 31, p. 179-302.
- Warraich, M. Y. Ogawara, K. and Nishi, H., 1999: Late Paleocene to early Eocene planktonic foraminiferal biostratigraphy of the Sulaiman Range, Southern Indus Basin, Pakistan (Submitted to *Paleontological Society of Japan*).
- Warraich, M.Y. and Natori, H., 1997: Geology and planktonic foraminiferal biostratigraphy of the Paleocene-Eocene succession of the Zinda Pir section, Sulaiman Range, Southern Indus Basin, Pakistan. *Bulletin of the Geological Survey of Japan*, vol. 48, p. 595 -630.

- Warraich, M.Y., Ali, M., Ahamd, M. A., Siddiqui, R. H. and Hirayama, J., 1995: Geology and structure of calcareous zone in Muslim Baghand Kila Saifullah Area, Pakistan. *Geologica, Geo. Surv. Pakistan, Geoscience Lab, Islamabad*, vol. 1 (1), p. 61-77.
- Warwick, P. D., Johnson, E. A. and Khan, I. H., 1998: Collision-induced tectonism along the northwestern margin of the Indian subcontinent as recorded in the Upper Paleocene to Middle Eocene strata of central Pakistan (Kirthar and Sulaiman Ranges). *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 142, p.201-216.
- Weinzierl, L. L. and Applin, E. R., 1929: The Claiborne Formation on the Coastal Domes. *Journal of Paleontology*, vol. 3, p. 384-410.
- White, M. P., 1928: Some index foraminifera of the Tempico embayment area of Mexico. *Journal of Paleontology*, vol. 2, p. 177-317.
- Williams, M. D., 1959: Stratigraphy of Lower Indus Basin, West Pakistan. *World Petroleum Cong., 5th, New York, Proc., Sec. 1 paper 19*, p. 377-390.
- Yeats, R. S. and Lawrence, R. D., 1984: Tectonics of the Himalayan thrust belt in northern Pakistan. In *Marine Geology and Oceanography of Arabian Sea and Coastal Pakistan*, edited by B. U. Haq and J. D. Milliman, Van Nostrand Reinhold, New York, p. 177-198.
- Zachos, J. C., Lohmann, K. C., Walker, J. D. G. and Wise, S. W., 1993: Abrupt climate change and transient climates during Paleogene: A marine perspective. *Journal of Geology*, vol. 101, p. 191-213.