

List of illustrations

Figures 1~16

- Figure 1.** Regional tectonic setting of Pakistan showing the deposition of Paleogene marine sedimentary deposits distributed in fold and thrust belt of Pakistan formed as a result of the Indo-Pakistan and Eurasian Plate. Here the rectangular box indicates the position of the Sulaiman Range (Modified after Kazmi and Rana, 1982; Hamyon, 1991) -----3
- Figure 2.** The geological map of the Sulaiman Range showing the locations of the Rakhi Nala section and the Zinda Pir Anticline (Modified after Kazmi and Rana 1982; Hamyon et al., 1991) -----7
- Figure 3.** Route geological map of the Rakhi Nala section showing lithostratigraphic units and sample locations, exposed along the Rakhi Nala (river), Sulaiman Range, Southern Indus Basin, Pakistan -----8
- Figure 4.** Route geological map of the Zinda Pir Anticline showing lithostratigraphic units and sample locations, exposed along the Sori Nadi (river), Sulaiman Range, Southern Indus Basin, Pakistan -----9
- Figure 5.** Regional tectonic map of Pakistan and adjoining areas showing main tectonic features such as major tectonic blocks, faults and structure of the fold and thrust belt (Modified after Bannert et al, 1992, Bender, 1995). Here
BWZ: Bela Waziristan Ophiolite Zone, I: Islamabad, KBF: Kirthar Basement Fault, KHB: Khuzdar Block, KOHA: Kohistan Island Arc, KTS: Kirthar Thrust Sheet, MBH: Mari Bugti Hills, MMT: Main Mantle Thrust, ONF: Ornach-Nal

Fault, SBF: Sulaiman Basement Fault, SR: Sulaiman Range, WSTF:
Western Sulaiman Transform Fault -----13

Figure 6. Cenozoic lithostratigraphic correlation of this work with that of previous workers (Eames, 1952b; Shah, 1977, and Kazmi, 1995) in the Sulaiman Range, Southern Indus Basin, Pakistan. -----16

Figure 7. Cenozoic lithostratigraphic subdivisions of the Sulaiman Range, Southern Indus Basin, Pakistan -----17

Figure 8. Correlation of the Paleogene lithostratigraphic units exposed in the Rakhi Nala section and in the Zinda Pir Anticline, Sulaiman Range, Southern Indus Basin, Pakistan. This figure also shows established planktonic foraminiferal biostratigraphic zones and the criteria (datum markers) used for the definition of the bio-zones along with other characteristic planktonic foraminiferal assemblage used for the definition of the bio-zones established in the Sulaiman Range, Southern Indus Basin, Pakistan -----33

Figure 9. Distribution and stratigraphic ranges of the planktonic foraminiferal taxa in the Dunghan, Shaheed Ghat, Baska and Kirthar Formations exposed along the Rakhi Nala (river) in the Sulaiman Range, Southern Indus Basin, Pakistan. Measured columnar shows lithofacies changes and sampling locations. Here small filled circles indicate occurrence of the planktonic foraminifers whereas small horizontal lines against the lithology represent sampling points and numbers indicate samples containing planktonic foraminifera -----37

Figure 10. Distribution and stratigraphic ranges of the planktonic foraminiferal taxa in the Dunghan, Shaheed Ghat, Baska and Kirthar Formations exposed in the Zinda Pir eastern section, Sulaiman Range, Southern Indus Basin, Pakistan. Measured columnar shows lithofacies changes and sampling locations. Here small filled circles indicate occurrence of the planktonic foraminifers whereas small horizontal lines against the lithology represent sampling points and numbers indicate samples containing planktonic foraminifera -----38

Figure 11. Distribution and stratigraphic ranges of the planktonic foraminiferal taxa in the Dunghan, Shaheed Ghat, Baska and Kirthar Formations exposed in the Zinda Pir western section, Sulaiman Range, Southern Indus Basin, Pakistan. Measured columnar shows lithofacies changes and sampling locations. Here small filled circles indicate occurrence of the planktonic foraminifers whereas small horizontal lines against the lithology represent sampling points and numbers indicate samples containing planktonic foraminifera -----39

Figure 12. Correlation planktonic foraminiferal biostratigraphic zones established herein the Sulaiman Range (this work) with internationally recognized zonal schemes of Bolli, 1957; Premoli Silva & Bolli, 1973; Blow, 1979; Toumarkine and Luterbacher, 1985; Berggren and Miller, 1988; Berggren et al., 1995; Berggren and Noriss, 1997, and Olsson et al., 1999. Correlation with biostratigraphic zones of Samanta (1973) and Dorreen (1974) established in Sulaiman and Kirthar Ranges, respectively, is also given -----46

Figure 13. Dominance (the most abundant taxon; *M. acuta*), diversity, P- ratio and relative abundances of the characteristic planktonic foraminiferal species with lithology, biostratigraphy and position in the measured columnar section along the eastern limb of the Zinda Pir Anticline. The numbers indicate the samples containing planktonic foraminifers whereas small horizontal lines against lithology indicate sample positions -----163

Figure 14. Dominance (the most abundant taxon; *M. acuta*), diversity, P- ratio and relative abundances of the characteristic planktonic foraminiferal species with lithology, biostratigraphy and position in the measured columnar section along the western limb of the Zinda Pir Anticline. The numbers indicate the samples containing planktonic foraminifers whereas small horizontal lines against lithology indicate sample positions -----164

Figure 15. Dominance (the most abundant taxon; *M. acuta*), diversity, P-ratio and relative abundances of the characteristic planktonic foraminiferal species with lithology, biostratigraphy and position in the measured columnar section from Rakhi Nala section. The numbers indicate the samples containing planktonic foraminifers whereas small horizontal lines against lithology indicate sample positions -----165

Figure 16. Lithofacies and microfacies correlation showing depositional environments for Paleogene strata of the Sulaiman Range, Southern Indus Basin, Pakistan -----185

Tables 1~6

Table 1. Distribution and relative abundances (%) of the planktonic foraminiferal species in the Dunghan Formation exposed along the Rakhi Nala (river), Sulaiman Range, Southern Indus Basin, Pakistan. Here x= less than 1%--40

Table 2. Distribution and relative abundances (%) of the planktonic foraminiferal species in the Dunghan Formation exposed along eastern and western limbs of the Zinda Pir Anticline, eastern Sulaiman Range, Southern Indus Basin, Pakistan. Here x= less than 1%-----41

Table 3. Distribution of the planktonic foraminiferal species identified from the Shaheed Ghat, Baska, and Kirthar Formations exposed along the Rakhi Nala (river), Sulaiman Range, Southern Indus Basin, Pakistan. Here X represents the occurrence of the species. Here the numbers represent actual relative abundances as found in the representative species count numbers -----42

Table 4. Distribution of the planktonic foraminiferal species identified from the Shaheed Ghat, Baska, and Kirthar Formations exposed along the eastern limb of the Zinda Pir Anticline, Sulaiman Range, Southern Indus Basin, Pakistan. Here X represents the occurrence of the species -----43

Table 5. Distribution of the planktonic foraminiferal species identified from the Shaheed Ghat, Baska, and Kirthar Formations exposed along the western limb of the Zinda Pir Anticline, Sulaiman Range, Southern Indus Basin, Pakistan. Here X represents the occurrence of the species -----44

Table 6. Distribution of the benthic foraminiferal species identified from the Dunghan Formation exposed in the Rakhi Nala section, Sulaiman Range, Southern Indus Basin, Pakistan. Here X represents the occurrence of the species -----171