

References

1. Hall, J.E., and Guyton, A.C. (1990) in *Hypertension: Pathophysiology, Diagnosis and Management* (Laragh, J. H., and Brenner, B.M., eds) pp.1105-1129, Raven Press, New York
2. Timmermans, P. B., Wong, P. C., Chiu, A. T., Herblin, W. F., Benfield, P., Carini, D. J., Lee, R. J., Wexler, R. R., Saye, J. A., and Smith, R. D. (1993) Angiotensin II receptors and angiotensin II receptor antagonists. *Pharmacol Rev* **45**, 205-251
3. Iwai, N., and Inagami, T. (1992) *FEBS Lett.* **298**, 257-260
4. Fukamizu, A., Sugimura, K., Takimoto, E., Sugiyama, F., Seo, M. S., Takahashi, S., Hatae, T., Kajiwara, N., Yagami, K., and Murakami, K. (1993) Chimeric renin-angiotensin system demonstrates sustained increase in blood pressure of transgenic mice carrying both human renin and human angiotensinogen genes. *J Biol Chem* **268**, 11617-11621
5. Tanimoto, K., Sugiyama, F., Goto, Y., Ishida, J., Takimoto, E., Yagami, K., Fukamizu, A., and Murakami, K. (1994) Angiotensinogen-deficient mice with hypotension. *J.Biol.Chem.* **269**, 31334-31337
6. Sasamura, H., Hein, L., Krieger, J.E., Pratt, R.E., Kobilka, B.K., and Dzau, V.J. (1992) *Biochem. Biophys. Res. Comm.* **185**, 253-259
7. Saga, Y., Yagi, T., Ikawa, Y., Sakakura, T., and Aizawa, S.. (1992) *Genes & Development* **6**, 1821-1831
8. Yagi, T., Tokunaga, T., Furuta, Y., Nada, S., Yoshida, M., Tsukada, T., Saga, Y., Takeda, N., Ikawa, Y., and Aizawa, S. (1993) *Anal. Biochem.* **214**, 70-76
9. Imaizumi, K., Tsuda, M., Wanaka, A., Tohyama, M., and Takagi, T. (1994) *Mol. Brain Res.* **26**, 189-196
10. Chomczynski, P., and Sacchi, N. (1987) *Anal. Biochem.* **162**, 156-159
11. Tanimoto, K., Murakami, K., and Fukamizu, A. (1994) *Exp. Cell Res.* **211**, 408-414
12. Printz, M.P., Printz, J.M., and Dworschack, R.T. (1977) *J. Biol. Chem.* **252**,

1654-1662.

13. Johns, D.W., Peach, M.J., Gomez R.A., Inagami, T., and Carey, R.M. (1990) *Am. J. Physiol.* **259**, F882-F887
14. Smithies, O., and Kim, H.-S. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 3612-3615
15. Ito, M., Oliverio, M. I., Mannon, P. J., Best, C. F., Maeda, N., Smithies, O., and Coffman, T. M. (1995) Regulation of blood pressure by the type 1A angiotensin II receptor gene. *Proc Natl Acad Sci U S A* **92**, 3521-3525
16. Blair-West, J.R., Coghlan, J.P., Denton, D.A., Funder, J.W., Scoggins, B.A., and Wright, R.D. (1971) *Am. J. Physiol.* **220**, 1309-1315
17. Vandongen, R., Peart, W.S., and Boyd, G.W. (1974) *Am. J. Physiol.* **226**, 277-282
18. Pinet, F., Mizrahi, J., Laboulandine, I., Menard, J., and Corvol, P. (1987) *J. Clin. Invest.* **80**, 724-731
19. Sugaya, T., Nishimatsu, S., Tanimoto, K., Takimoto, E., Yamagishi, T., Imamura, K., Goto, S., Imaizumi, K., Hisada, Y., Otsuka, A., Uchida, H., Sugiura, M., Fukuta, K., Fukamizu, A., and Murakami, K. (1995) Angiotensin II type 1a receptor-deficient mice with hypotension and hyperreninemia. *J Biol Chem* **270**, 18719-18722
20. Kimura, K., Inokuchi, S., Sugaya, T., Suzuki, N., Yoneda, H., Shirato, I., Mise, N., Oba, S., Miyashita, K., Tojo, A., Hirata, Y., Goto, A., Sakai, T., Murakami, K., and Omata, M. (1997) Location and action of angiotensin II type 1 receptor in the renal microcirculation. *Kidney Int Suppl* **63**, S201-204
21. Anderson Pw, Do YS, Hsueh WA (1993) Angiotensin II causes mesangial cell hypertrophy. *Hypertension* **21**, 29-35
22. Blants RC, Konnen KS, Tucker BJ (1976) Angiotensin II effects upon the glomerular microcirculation and ultrafiltration coefficient of the rat. *J Clin Invest* **57**:419-434
23. Border WA, Okuda S, Languino LR, Ruoslaht IE (1990) Transforming

growth factor-b regulates production of proteoglycans by mesangial cells.
Kidney Int 37:689-695

24. Chen X, Li W, Yoshida H, Tsuchida S, Nishimura H, Takemoto F, Okubo S, Fogo A, Matsusaka T, Ichikawa I (1997) Am J Physiol 272:F299-F304
25. Denton KM, Fensy PA, Alcorn D, Anderson WP (1992) Morphometric analysis of the actions of angiotensin II on renal arterioles and glomeruli. Am J Physiol 262:F367-372
26. Edwards DR, Murphy G, Reynolds JJ, Whitham SE, Docherty JP, Angel P, Heath JK (1987) Transforming growth factor beta modulates the expression of collagenase and metalloproteinase inhibitor. Eur Mol Biol Organ J 6:1899-1904
27. Elger M, Sakai T, Kriz W (1990) Role of mesangial cell contraction in adaptation of the glomerular tuft to changes in extracellular volume. Europ J Physiol 415:598-605
28. Hahn AWA, Resnick TJ, Bernhardt TJ, Feraccin F, Buhler FR (1991) Stimulation of autocrine platelet-derived growth factor AA-homodimer and transforming growth factor b in vascular smooth muscle cells. Biochem Biophys Res Commun 178:1451-1458
29. Hein L, Barsh GS, Pratt RE, Dzau VJ, Kobilka BK (1995) Behavioural and cardiovascular effects of disrupting the angiotensin II type-2 receptor gene in mice. Nature 377:744-747
30. Hirose K, Osterby R, Nakazawa M, Jorgensen H, Gumderson G (1982) Development of glomerular lesions in experimental long-term diabetes in the rat. Kidney Int 21:689-695

- 31 Ichiki T, Labosky PA, Shiota C, Okuyama S, Imagawa Y, Fogo A, Niimura F, Ichikawa I, Hogan BLM, Inagami T (1995) Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type-2 receptor. *Nature* 377:748-750
- 32 Ito M, Oliverio MI, Mannon PJ, Best CF, Maeda N, Smities O, Coffman TM (1995) Regulation of blood pressure by the type 1a angiotensin II receptor gene. *Proc Natl Acad Sci USA* 92:3521-3525
33. Kakinuma Y, Fogo A, Inagami T, Ichikawa I (1993) Intrarenal localization of angiotensin II type 1 receptor mRNA in the rat. *Kidney Int* 43:1229-1235
34. Kimura K, Nagai R, Sakai T, Aikawa M, Kuro-o M, Kobayashi N, Shirato I, Inagami T, Oshi M, Suzuki N, Oba S, Mise N, Tojo A, Hirata Y, Goto A, Yazaki Y (1995) Diversity and variability of smooth muscle phenotypes of renal arterioles as revealed by myosin isoform expression. *Kidney Int* 48:372-382
35. Lee LK, Meyer TW, Pollock AS, Lovett DH (1995) Endothelial cell injury initiates glomerular sclerosis in the rat remnant kidney. *J Clin Invest* 96:953-964
36. Marti HP, Lee L, Kashgarian M, Lovett DH (1994) Transforming growth factor- β 1 stimulates glomerular mesangial cell synthesis of the 72 kDa type IV collagenase. *Am J Pathol* 144:82-94
37. Myers BD, Deen WM, Brenner BM (1975) Effects of norepinephrine and angiotensin II on the determinants of glomerular ultrafiltration and proximal tubule fluid reabsorption in the rat. *Circ Res* 37:101-110
- 38 Niimura F, Labosky PA, Kakuchi J, Okubo S, Yoshida H, Oikawa T, Ichiki T, Naftilan AJ, Fogo A, Inagami T, Hogan BLM, Ichikawa I (1995) Gene

targeting in mice reveals a requirement for angiotensin in the development and maintenance of kidney morphology and growth factor regulation. *J Clin Invest* 96:2947-2954

39. Paxton WG, Runge M, Horaist C, Cohen C, Alexander RW, Bernstein KE (1993) Immunohistochemical localization of rat angiotensin II AT1 receptor. *Am J Physiol* 264:F989-995
40. Ray PE, Aguilera G, Kopp JB, Horikoshi S, Klotman PE (1991) Angiotensin II receptor-mediated proliferation of cultured human fetal mesangial cells. *Kidney Int* 40:764-771
41. Ray PE, Bruggman LA, Horikoshi S, Aguilera G, Klotman PE (1994) AngiotensinII stimulates human fetal mesangial cell proliferation and fibronectin biosynthesis by binding to AT1 receptors. *Kidney Int* 45:177-184
42. Sakai T, Kriz W (1987) The structural relationship between mesangial cells and basement membrane of the renal glomerulus. *Anat Embryol* 176:373-386
43. Shirato I, Sakai T, Fukui M, Tomino Y, Koide H (1993) Widening of capillary neck and alteration of extracellular matrix ultrastructure in diabetic rat glomerulus as revealed by computer morphometry and improved tissue processing. *Virchow Archiv A Pathol Anat* 423:121-129
44. Stouffer GA, Owens GK (1992) Angiotensin II-induced mitogenesis of spontaneously hypertensive rat-derived cultured smooth muscle cells is dependent on autocrine production of transforming growth factor-b. *Circ Res* 70:820-828
45. Kihara M, Umemura S, Sugaya T, Toya Y, Yabana M, Kobayashi S, Tamura K,

- Kadota T, Kishida R, Murakami K, Fukamizu A, Ishii M (1998) Expression of neuronal type nitric oxide synthase and renin in the juxtaglomerular apparatus of angiotensin type-1a receptor gene-knockout mice. *Kidney Int* 53, 1585-1593
46. Weibel ER (1979) *Stereological methods. Practical methods for biological morphometry.* Academic Press, London vol 1
47. Weihprecht H, Lorenz JN, Briggs JP, Schnermann J (1991) Vasoconstrictor effect of angiotensin and vasopressin in isolated rabbit afferent arterioles. *Am J Physiol* 261:F273-282
48. Wolf G, Neilson EG (1993) Angiotensin II as a renal growth factor. *J Am Soc Nephrol* 3:1531-1540
49. Wolf G, Haberstroh U, Neilson EG (1992) Angiotensin II stimulates the proliferation and biosynthesis of type I collagen in cultured murine mesangial cells. *Am J Pathol* 140:95-107
50. Matsusaka, T., and Ichikawa, I. (1997) Biological function of angiotensin and its receptors. *Annu Rev Physiol* 59, 395-412
51. Gomez-Garre D, Ruiz-Ortega M, Ortego M, Larugo R, Lopez-Armada MJ, Plaza JJ Gonzales E, Egido (1996) J: Effects and interaction of endothelin-1 and angiotensin II on matrix protein expression and synthesis and mesangial cell growth. *Hypertension* 27:885-892
52. Edwards RM (1983) Segmental effects of norepinephrine and angiotensin II on isolated renal microvessels. *Am J Physiol* 244:F526-F534
53. Ito S, Johnston CS, Carretero OA (1991) Modulation of angiotensin II-induced vasoconstriction by endothelium-derived relaxing factor in the isolated

- microperfused rabbit afferent arteriole. *J Clin Invest* 87:1656-1663
54. Dworkin LD, Brenner BM: The renal circulation, in *The kidney*, edited by Brenner BM, Rector FC (1991) Philadelphia, Saunders Company, p. 164
 55. Kimura K, Nagai R, Sakai T, Aikawa M, Kuro-o M, Kobayashi N, Shirato I, Inagami T, Oshi M, Suzuki N, Oba S, Mise N, Tojo A, Hirata Y, Goto A, Yazaki Y, Omata M (1995) Diversity and variability of smooth muscle phenotypes of renal arterioles as revealed by myosin isoform expression. *Kidney Int* 48:372-382
 56. Johns DW, Peach MJ, Gomez RA, Inagami T, Carey RM: Angiotensin II regulates renin gene expression.(1990) *Am J Physiol* 259: F882-F887
 57. Steinhausen M, Snoei H, Parekh N, Baker R, Johnson PC: Hydronephrosis (1983) a new method to visualize vas afferens, efferens, and glomerular network. *Kidney Int* 23:794-806
 58. Kimura K, Tojo A, Hirata Y, Hayakawa H, Goto A, Omata M (1994) Effects of a calcium antagonist and an angiotensin II receptor antagonist on rat renal arterioles. *Blood Press Suppl* 5:71-74
 59. Sakai T, Kriz W The structural relationship between mesangial cells and basement membrane of the renal glomerulus. *Anat Embryol (Berl)* 176:373-386, 1987
 60. Inokuchi S, Kimura K, Sugaya T, Yoneda H, Shirato I, Murakami K, Sakai T Angiotensin II maintains the structure and function of glomerular mesangium via type 1a receptor. What we have learned from null mutant mice minus the angiotensin II type 1a receptor gene. *Virchows Arch.* 433, 349-357, 1998
 61. Hackenthal E, Paul M, Ganten D, and Taugner R: Morphology, physiology, molecular biology of renin secretion. *Physiol Rev* 70: 1067-1116, 1990
 62. Lorenz JN, Weihprecht H, Schnermann J, Skøtt O, Briggs JP: Renin release

- from isolated juxtaglomerular apparatus depends on macula densa chloride transport. *Am J Physiol* 260: F486-F493, 1991
63. Persson AEG, Salomonsson M, Westerlund P, Greger R, Schlatter E, Gonzales E: Macula densa cell function. *Kidney Int* 39: S39-S44, 1991
 64. Briggs JP, Schnermann JB: Whys and wherefores of juxtaglomerular apparatus function. *Kidney Int* 49: 1724-1726, 1996
 65. Ito S, Carretero OA, Abe K: Nitric oxide in the juxtaglomerular apparatus. *Kidney Int* 49: S6-S8, 1996
 66. Persson PB, Baumann JE, Ehmke H, Hackenthal E, Kirchheim HR, Nafz B: Endothelium-derived NO stimulates pressure-dependent renin release in conscious dogs. *Am J Physiol* 264: F943-F947, 1993
 67. Beierwaltes WH: Selective neuronal nitric oxide synthase inhibition blocks furosemide-stimulated renin secretion in vivo. *Am J Physiol* 269: F134-F139, 1995
 68. Schricker K, Ivan H, Hamann M, Kaissling B, Kurtz A: Tonic stimulation of renin gene expression by nitric oxide is counteracted by tonic inhibition through angiotensin II. *Proc Natl Acad Sci USA* 92: 8006-8010, 1995
 69. Schricker K, Kurtz A: Blockade of nitric oxide formation inhibits the stimulation of the renin system by a low salt intake. *Pflügers Arch* 432: 187-191, 1996
 70. Thorup C, Persson AEG: Macula densa derived nitric oxide in regulation of glomerular capillary pressure. *Kidney Int* 49: 430-436, 1996
 71. Beierwaltes WH: Macula densa stimulation of renin is reversed by selective inhibition of neuronal nitric oxide synthase. *Am J Physiol* 272: R1359-R1364,

1997

72. Tharaux PL, Dussaule jc, Pauti MD, Vasitch Y, Ardaillou R, Chatziantoniou C: Activation of renin synthesis is dependent on intact nitric oxide production. *Kidney Int* 51: 1780-1787, 1997
73. Tojo A, Gross SS, Ahang L, Tischer CC, Schmidt HHW, Wilcox CS, Madsen KM: Immunocytochemical localization of distinct isoforms of nitric oxide synthase in the juxtaglomerular apparatus of normal rat kidney. *J Am Soc Nephrol* 4: 1438-1447, 1994
74. Mundel P, Bachmann S, Bader M, Fischer A, Kummer W, Mayer B, Kriz W: Expression of nitric oxide synthase in kidney macula densa cells. *Kidney Int* 42: 1017-1019, 1992
75. He X-R, Greenberg SG, Briggs JP, Schnermann JB: Effect of nitric oxide on renin secretion II. Studies in the perfused juxtaglomerular apparatus. *Am J Physiol* 268: F953-F959, 1995
76. Bosse HM, Böhm R, Resch S, and Bachmann S: Parallel regulation of constitutive NO synthase and renin at JGA of rat kidney under various stimuli. *Am J Physiol* 269: F793-F805, 1995
77. Singh I, Grams J, Wang WH, Yang TX, Killen P, Smart A, Schnermann J, Briggs JP: Coordinate regulation of renal expression of nitric oxide synthase, renin, and angiotensinogen mRNA by dietary salt. *Am J Physiol* 270: F1027-F1037, 1996
78. Vidal JM, Romero JC, Vanhoutte PM: Endothelium-derived relaxing factor inhibits renin release. *Eur J Pharmacol* 149: 401-402, 1988

79. Sigmon DH, Carretero OA, Beierwaltes WH: Endothelium-derived relaxing factor regulates renin release in vivo. *Am J Physiol* 263: F256-F261, 1992
80. Ichihara A, Suzuki H, Murakami M, Naitoh M, Matsumoto A, Saruta T: Interactions between angiotensin II and norepinephrine on renin release by juxtaglomerular cells. *Eur J Endocrinol* 133: 569-577, 1995
81. Matsusaka T, Nishimura H, Utsunomiya H, Kakuchi J, Niimura F, Inagami T, Fogo A, Ichikawa I: Chimeric mice carrying regional targeted deletion of the angiotensin type 1a receptor gene. *J Clin Invest* 98: 1867-1877, 1996
82. Tanimoto K, Sugiyama F, Goto Y, Ishida J, Takimoto E, Yagami K, Fukamizu A, Murakami K: Angiotensinogen-deficient mice with hypotension. *J Biol Chem* 269: 31334-31337, 1994
83. Kihara M, Umemura S, Kadota T, Tabana M, Tamura K, Nyuui N, Ogawa N, Murakami K, Fukamizu A, Ishii M: The neuronal isoform of constitutive nitric oxide synthase is up-regulated in the macula densa of angiotensinogen gene-knockout mice. *Lab Invest* 76: 285-294, 1997
84. Kihara M, Umemura S, Yabana M, Sumida Y, Nyui N, Tamura K, Kadota T, Kishida R, Murakami K, Fukamizu A, Ishii M. Dietary salt loading decreases the expression of neuronal type nitric oxide synthase and renin in the juxtaglomerular apparatus of angiotensinogen gene-knockout mice. *J Am Soc Nephrol* (in press)
85. Kihara M, Umemura S, Sumida Y, Yokoyama N, Yabana M, Nyui N, Tamura K, Murakami K, Fukamizu A, Ishii M. Genetic deficiency of angiotensinogen produces an impaired urine concentrating ability in mice. *Kidney Int* (in press)
86. Umemura S, Kihara M, Sumida Y, Yabana M, Ishigami T, Tamura K, Nyui N, Hibi K, Murakami K, Fukamizu A, Ishii M: Endocrinological abnormalities in

angiotensinogen-gene knockout mice: studies on hormonal responses to dietary salt loading. *J Hypertension* (in press)

87. Tojo A, Madsen KM, Wilcox CS: Expression of immunoreactive nitric oxide synthase isoforms in rat kidney. *Jpn Heart J* 36: 389-398, 1995
88. Kihara M, Kubo T: Immunocytochemical localization of GABA containing neurons in the ventrolateral medulla oblongata of the rat. *Histochem* 91:309-314, 1989
89. Chomczynski P, Sacchi N: Single-step method of RNA isolation by acid guanidium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156-159, 1987
90. Shaul P W, North AJ, Brannon TS, Ujiie K, Wells LB, Nisen PA, Lowenstein CJ, Snyder SH, Star RA: Prolonged in vivo hypoxia enhances nitric oxide synthase type I and type III gene expression in adult rat lung. *Am J Respir Cell Mol Biol* 13: 167-174, 1995
91. Dzau VJ, Brody T, Ellison KE, Pratt RE, Ingelfinger JR: Tissue-specific regulation of renin expression in the mouse. *Hypertension* 9(Suppl III): 36-41, 1987
92. Miller CCJ, Samani NJ, Carter AT, Brooks JJ, Brammar WJ: Modulation of mouse renin gene expression by dietary sodium chloride intake in one-gene, two-gene and transgenic animals. *J Hypertension* 7: 861-863, 1989
93. Sharp MGF, Fettes D, Brooker G, Clark AF, Peters J, Fleming S, Mullins JJ: Targeted inactivation of the Ren-2 gene in mice. *Hypertension* 28: 1126-1131, 1996
94. Matsumoto T, Nakane M, Pollock JS, Kuk JE, Förstermann U: A correlation between soluble brain nitric oxide synthase and NADPH-diaphorase activity is only seen after exposure of the tissue to fixative. *Neurosci Lett* 155: 61-63, 1993

95. Schricker K, Della Bruna R, Jamann M, Kurtz A: Endothelium derived relaxing factor is involved in the pressure control of renin gene expression in the kidney. *Pflügers Arch* 428: 261-268, 1994
96. Blaine EH, Davis JO, Prewitt RL: Evidence for a renal vascular receptor in control of renin secretion. *Am J Physiol* 220: 1593-1597, 1970
97. Fray JCS: Stretch receptor model for renin release with evidence from perfused rat kidney. *Am J Physiol* 231: 936-944, 1977
98. Bock HA, Hermle M, Fiallo A, Osgood RW, Fried TA: Measurement of renin secretion in single perfused rabbit glomeruli. *Am J Physiol* 258: F1460-F1465, 1990
99. Salomonsson M, Skott O, Persson AEG: Influence of intraluminalarterial pressure on renin release. *Acta Physiol Scand* 141: 285-286, 1991
100. Lorenz JN, Weihprecht H, He X-R, Skøtt O, Briggs JP, Schnermann J: Effect of adenosine and angiotensin on macula densa-stimulated renin secretion. *Am J Physiol* 1993, 265: F187-F194
101. de Gasparo R, Levens NR: Pharmacology of angiotensin II receptors in the kidney. *Kidney Int.* 1994;46:1486-1491
102. Douglas JG, Hopfer U: Novel aspect of angiotensin receptors and signal transduction in the kidney. *Annu. Rev. Physiol.* 1995;56:649-669
103. Inagami T, Kitami Y: Angiotensin II receptor: molecular cloning, functions, and regulation. *Hypertens Res.* 1994; 17: 87-97
104. Edwards RM, Stack EJ, Weidley EF, Aiyar N, Keehan RM, Hill DT, Weinstock J: Characterization of renal angiotensin II receptors using subtype selective antagonists. *J Pharmacol Exp Ther* 1992;260:933-938

105. Gauquelin G, Garcia R. Characterization of glomerular angiotensin II receptor subtypes: *Receptor* 1992;2:207-212
106. Gibson RE, Thorpe HH, Cartwright ME, Frank JD, Schorn TW, Bunting PB, Siegl PK. Angiotensin II receptor subtypes in renal cortex of rats and rhesus monkeys: *Am J Physiol* 1991;261:F512-518
107. Chang RS, Lotti VJ. Angiotensin II receptor subtypes in rat, rabbit and monkey tissues: Relative distribution and species dependency: *Life Sci.* 1991;49:1485-1490
108. Sechi LA, Grady EF, Griffin CA, Kalinyak JE, Schambelan M. Distribution of angiotensin II receptor subtypes in rat and human kidney: *Am J Physiol.* 1992;262:F236-240
109. Jourdain M, Amiel C, Friedlander G: Modulation of Na-H exchange activity by angiotensin II in opossum kidney cells. *Am J Physiol* 1992;263:C1141-1146
110. Grone HJ, Simon M, Fuchs E. Autoradiographic characterization of angiotensin receptor subtypes in fetal and human kidney. *Am J Physiol* 1992;262:F326-F331
111. Goldfarb DA, Diz DI, Tubbs RR, Ferrario CM, Novick AC. Angiotensin II receptor subtypes in the human renal cortex and renal cell carcinoma. *J Urol* 1994;151:208-213
112. De leon H, Garcia R. Angiotensin II receptor subtypes in rat renal preglomerular vessels. *Receptor* 1992;2:253-260
113. Chansel D, Vandermeersch S, Pham P, Ardaillou R. Characterization of [³H]losartan receptors in isolated rat glomeruli. *Eur J Pharmacol-Molec Pharm.* 1993;247:193-198
114. Terada Y, Tomita K, Nonoguchi H, Marumo F. PCR localization of angiotensin II receptor and angiotensinogen messenger RNAs in rat kidney. *Kidney Int* 1993;43:1251-1259

115. Kakinuma Y, Fogo A, Inagami T, Ichikawa I. Intrarenal localization of angiotensin II type I receptor messenger RNA in the rat. *Kidney Int* 1993;43:1229-1235
116. Meister B, Lippoldt A, Bunnemann B, Inagami T, Ganten D, Fuxe K. Cellular expression of angiotensin type-1 receptor messenger RNA in the kidney. *Kidney Int.* 1993;44:331-336
117. Paxton WG, Runge M, Horaist C, Cohen C, Alexander RW, Bernstein KE. Immunohisto-chemical localization of rat angiotensin II AT1 receptor. *Am J Physiol* 1993;264:F989-F995
118. Matsubara H, Sugaya T, Murasawa S, Nozawa Y, Mori Y, Masaki H, Maruyama K, Tsutumi Y, Shibasaki Y, Moriguchi Y, Tanaka Y, Iwasaka T, Inada M Tissue-specific expression of human angiotensin II AT1 and AT2 receptors and cellular localization of subtype mRNAs in adult human renal cortex using in situ hybridization. *Nephron* 1998; 80:25-34
119. Kambayashi Y, Bardhas S, Takahashi K, Tsuzuki S, Inui H, Hamakubo T, Inagami T. Molecular cloning of a novel angiotensin II receptor isoform involved in phosphotyrosine phosphatase inhibition. *J Biol Chem.* 1993; 268: 24543-24546
120. Herblin WF, Diamond SM, Timmermans PBMWM. Localization of angiotensin II receptor subtypes in the rabbit adrenal and kidney. *Peptides.* 1991;12:581-584
121. Suzuki J, Matsubara H, Inada M: Rat angiotensin II (type 1A) receptor mRNA regulation and Subtype expression in myocardial growth and hypertrophy. *Circ Res* 1993;73: 439-447
122. Matsubara H., Kanasaki M, Murasawa S, Tsukaguchi Y, Nio Y, Inada M. Differential gene expression and regulation of angiotensin II receptor subtypes in rat cardiac fibroblasts and cardiomyocytes in culture. *J. Clin. Invest.* 1994;93: 1592-1601.

123. Nozawa Y, Haruno A, Oda N, Yamazaki Y, Yamada S, Suzuki H: Angiotensin II receptor subtypes in bovine and human ventricular myocardium. *J Pharmacol Exp Ther* 1994;270: 566-571
124. Mori Y., Matsubara H, Murasawa S, Kijima K, Maruyama K, Hamakubo N, Inagami T, Iwasaka T, Inada M. Translational regulation of angiotensin II receptor: Role of the upstream AUG triplets. *Hypertension*. 1996;28:810-817
125. Murasawa S, Matsubara H, Kijima K, Maruyama K, Mori Y, Inada M. Structure of the V1a vasopressin receptor gene and characterization of its promoter region and complete cDNA sequence of the 3'-end. *J Biol Chem*. 1995;270: 20042-20050
126. Kizima K, Matsubara H, Murasawa S, Maruyama K, Ohkubo N, Mori Y, Ohkubo N, Komuro I, Yazaki Y, Iwasaka T, Inada M. Mechanical stretch induces enhanced expression of angiotensin II receptor subtypes in neonatal rat cardiac myocytes. *Circ Res*. 1996;79:887-897
127. Kizima K, Matsubara H, Murasawa S, Ohkubo N, Mori Y, Inada M. Regulation of angiotensin II type 2 receptor gene by the protein kinase C-calcium pathway. *Hypertension* 1996;27:529-534
128. de Gasparo M, Rogg H, Brink M, Wang L, Whitebread S, Bullock G, Erne P. Angiotensin II receptor subtypes and cardiac function. *Eur Heart J*. 1994;15 Supp D: 98-103.
129. Regitz-Zagrosek V, Friedel N, Heymann A, Bauer P, NeuÉl M, Rolfs A, Steffen C, Hildebrandt A, Hetzer R, Fleck E. Regulation, Chamber localization, and subtype distribution of angiotensin II receptors in human hearts. *Circulation*. 1995;91: 1461-1471.

130. Zhou J, Ernsberger P, Douglas JG. Angiotensin II receptor subtypes in rat renal mesangial cells. *FASEB J.* 1991;5:A870
131. Whitebread S, Michele M, Kamber B, de Gasparo, M. Preliminary biochemical characterization of two angiotensin II receptor subtypes. *Biochem Biophys Res Commun.* 1989;163:284-289Diz DI, Tubbs RR, Ferrario CM, Novick AC.