

## Acknowledgments

The author is most grateful to Dr. Penmetcha K. R. Kumar and Professor Kazunari Taira for their many advises and support.

The author thanks Ms. Shiori Koseki and Dr. Jun Ohkawa for construction of pU6 vectors and the stable cell line, HeLa cells carrying the chimeric HIV-1 LTR-Luc and Mr. Satoshi Fujita for helpful suggestions on kinetics. I would like to thank Mr. Shin-ichiro Kobayashi and Prof. Masato Katahira for NMR studies with free RNA<sup>Tat</sup> and RNA<sup>Tat</sup>-CQ complex. I would also like to thank Dr. Masahiro Iwakura and Dr. Yoshiyuki Tanaka for suggestions on CD measurements and analyses and Dr. Naruhisa Ota for useful discussions on the assay with fluorescence.

The author wishes to thank Professor Kazuo Murakami, Professor Tadashi Baba, and Dr. Satoshi Nishikawa for their support.

The author thanks the present and former members of Taira's laboratory and Nishikawa's laboratory for their kindness.

This research was supported by various grants from the Ministry of International Trade and Industry (MITI) of Japan and by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture, Japan. The author is a recipient of research fellowships for young scientists of Japan Society for the Promotion of Science.

Finally, the author is thankful to her family and her friends for their great support and encouragement.

## References

- Arya, S.K., Guo, C., Joseph, S.F. and Wong-Staal, F. (1985) *Trans*-activator gene of human T-lymphotropic virus type III (HTLV-III). *Science*, 229, 69-73.
- Berkhout, B., Silverman, R.H. and Jeang, K.T. (1989) Tat *trans*-activates the human immunodeficiency virus through a nascent RNA target. *Cell*, 59, 273-282.
- Berkhout, B. and Jeang, K.T. (1989) *Trans*-activation of human immunodeficiency virus type 1 is sequence-specific for both the single-stranded bulge and loop of the *trans*-acting-responsive hairpin: a quantitative analysis. *J. Virol.*, 63, 5501-5504.
- Berkhout, B. and Jeang, K.T. (1991) Detailed mutational analysis of TAR RNA: critical spacing between the bulge and loop recognition domains. *Nucleic Acids Res.*, 19, 6169-6176.
- Bohjanen, P.R., Colvin, R.A., Puttaraju, M., Been, M.D. and Garcia-Blanco, M.A. (1996) A small circular TAR RNA decoy specifically inhibits Tat-activated HIV-1 transcription. *Nucleic Acids Res.*, 24, 3733-3738.
- Calnan, B.J., Biancalana, S., Hudson, D. and Frankel, A.D. (1991) Analysis of arginine-rich peptides from the HIV Tat protein reveals unusual features of RNA-protein recognition. *Genes Dev.*, 5, 201-210.
- Chang, Y.-N. and Jeang, K.-T. (1992) The basic RNA-binding domain of HIV-2 Tat contributes to preferential *trans*-activation of a TAR-2-containing LTR. *Nucleic Acids Res.*, 20, 5465-5472.
- Churcher, M.J., Lmont, C., Hamy, F., Dingwell, C., Greem, S.M., Lowe, A.D., Butler, P.D.G., Gait, M.J. and Karn, J. (1993) High-affinity binding of TAR RNA by the human immunodeficiency virus Tat

- protein requires amino acid residues flanking the basic domain and base pairs in the RNA stem. *J. Mol. Biol.*, 230, 90-110.
- Cordingly, M.G., La Femina, R.L., Callahan, P.L., Condra, J.H., Sardana, V.V., Graham, D.J., Nguyen, T.M., Le Grow, K., Gotlib, L., Schlabach, A.J. and Colonno, R.J. (1990) Sequence-specific interaction of Tat protein and Tat peptides with the transactivation-responsive sequence element of human immunodeficiency virus type 1 *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.*, 87, 8985-8989.
- Cullen, B.R. (1986) *Trans*-activation of human immunodeficiency virus occurs via a bimodal mechanism. *Cell*, 46, 973-982.
- Cullen, B.R. (1992) Mechanism of action of regulatory proteins encoded by complex retroviruses. *Microbiol. Rev.*, 56, 375-394.
- Dang, C.V. and Lee, W.M.F. (1989) Nuclear and nucleolar targetting sequences of c-erb-A, c-myb, N-myc, p53, HSP70, and HIV tat proteins. *J. Biol. Chem.*, 264, 18019-18023.
- Dayton, A.I., Sodroski, J.G., Rosen, C.A., Goh, W.C. and Haseltine, W.A. (1986) The *trans*-activator gene of the human T cell lymphotropic virus type III is required for replication. *Cell*, 44, 941-947.
- Dingwell, C., Ernberg, I., Gait, M.J., Green, S.M., Heaphy, S., Karn, J., Lowe, A.D., Singh, M., Skinner, M.A. and Valerio, R. (1989) Human immunodeficiency virus 1 Tat protein binds *trans*-activation-response region (TAR) RNA *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.*, 87, 8985-8989.
- Drolet, D.W., Moon-McDermott, L. and Roming, T.S. (1996) An enzyme-linked oligonucleotide assay. *Nature Biotechnol.*, 14, 1021-1025.

- Elangovan, B., Subramanian, T. and Chinnadurai, G. (1992) Functional comparison of the basic domains of the Tat proteins of human immunodeficiency virus type 1 and 2 in trans activation. *J. Virol.*, 66, 2031-2036.
- Ellington, A.D. and Szostak, J. (1992) Selection *in vitro* of single-stranded DNA molecules that fold into specific ligand-binding structures. *Nature*, 355, 850-852.
- Emerman, M., Guyader, M., Montagnier, L., Baltimore, D. and Muesing, M.A. (1987) The specificity of the human immunodeficiency virus type-2 transactivator is different from that of human immunodeficiency virus type 1. *EMBO J.*, 6, 3755-3760.
- Endo, S.-I., Kubota, S., Siomi, H., Adachi, A., Oroszlan, S., Maki, M. and Hatanaka, M. (1989) A region of basic amino-acid cluster in HIV-1 Tat protein is essential for *trans*-acting activity and nucleolar localization. *Virus Genes*, 3, 99-110.
- Ensoli, B., Barillari, G., Salahuddin, S.Z., Gallo, R.C. and Wong-Staal, F. (1990) Tat protein of HIV-1 stimulates growth of cells derived from Kaposi's sarcoma lesions of AIDS patients. *Nature*, 344, 84-86.
- Ensoli, B., Buonaguro, L., Barillari, G., Fiorelli, V., Gendelman, R., Morgan, R.A., Wingfield, P. and Gallo, R.C. (1993) Release, uptake, and effects of extracellular human immunodeficiency virus type 1 Tat protein on cell growth and viral *trans*-activation. *J. Virol.*, 67, 277-287.
- Feng, S. and Holland, E.C. (1988) HIV-1 Tat *trans*-activation requires the loop sequence within TAR. *Nature*, 334, 165-167.
- Ferbeyre, G., Bourdeau, V. and Cedergren, R. (1997) Does HIV tat protein also regulate genes of other viruses present in HIV infection? *Trends in Biochem. Sci.*, 22, 115-116.

- Fisher A.G., Feinberg, M.B., Josephs, S.F., Harper, M.E., Marselle, L.M., Reyes, G., Gonda, M.A., Aldovini, A., Debouk, C., Gallo, R.C. and Wong-Staal, F. (1986) The *trans*-activator gene of HTLV-III is essential for virus replication. *Nature*, 320, 367-371.
- Frankel, A.D., Brecht, D.S. and Pabo, C.O. (1988) Tat protein from human immunodeficiency virus forms a metal-linked dimer. *Science*, 240, 70-73.
- Garber, M.E., Wei, P., KewalRamani, V.N., Mayall, T.P., Herrmann, C.H., Rice, A.P., Littman, D.R. and Jones, R.A. (1998) The interaction between HIV-1 Tat and human cyclin T1 requires zinc and a critical cysteine residue that is not conserved in the murine CycT1 protein. *Genes Dev.*, 12, 3512-3527.
- Garcia, J.A., Harrich, D., Pearson L., Mitsuyasu, R. and Gaynor, R. (1988) Functional domains required for tat-induced transcriptional activation of the HIV-1 long terminal repeat. *EMBO J.*, 7, 3143-3147.
- Garcia-Martinez, L.F., Gopinath, M., Peters, P., Wu-baer, F. and Gaynor, R.B. (1995) Tat functions to stimulate the elongation properties of transcription complexes paused by the duplicated TAR RNA element on human immunodeficiency virus-2. *J. Mol. Biol.*, 254, 350-363.
- Gatignol, A., Buckler-White, A., Berkhout, B. and Jeang K.-T. (1991) Characterization of a human TAR RNA-binding protein that activates the HIV-1 LTR. *Science*, 251, 1597-1600.
- Gaynor, R. (1992) Cellular transcription factors involved in the regulation of HIV-1 gene expression. *AIDS*, 6, 347-363.
- Gold, L., Polisky, B., Uhlenbeck, O. and Yarus, M. (1995) Diversity of oligonucleotide functions. *Annu. Rev. Biochem.*, 64, 763-797.

- Graham, G.J. and Maio, J.J. (1990) RNA transcription of the human immunodeficiency virus *trans*-activation response element can inhibit action of the viral *trans*-activator. Proc. Natl. Acad. Sci. U.S.A., 87, 5817-5821.
- Guyader, M., Emerman, M., Sonigo, P., Clavel, F., Montagnier, L. and Alizon, M. (1987) Genome organization and trans-activation of the human immuno-deficiency virus type 2. Nature, 326, 662-669.
- Hannibal, M., Markovitz, D.M., Clark, N. and Nabel, G.J. (1993) Differential activation of human immunodeficiency virus type 1 and 2 transcription by specific T-cell activation signals. J. Virol., 67, 5035-5040.
- Harrich, D., Hsu, C., Race, E. and Gaynor, R.B. (1994) Differential growth kinetics are exhibited by human immunodeficiency virus type 1 TAR mutants. J. Virol., 68, 5899-5910.
- Harrich, D., Ulich, C., Garcia-Martinez, L.F. and Gaynor, R.B. (1997) Tat is required for efficient HIV-1 reverse transcription. EMBO J., 16, 1224-1235.
- Harrison, G.S., Long, C.J., Curiel, T.J., Maxwell, F. and Maxwell, H. (1992) Inhibition of human immunodeficiency virus-1 production resulting from transduction with a retrovirus containing an HIV-1-regulated diphtheria toxin A chain gene. Hum. Gen. Ther., 3, 461-469.
- Hauber, J. and Cullen, B.R. (1988) Mutational analysis of the *trans*-activation-responsive region of the human immunodeficiency virus type I long terminal repeat. J. Virol., 62, 673-679.
- Jakobovits, A., Smith, D.H., Jakobovits, E.B. and Capon, D.J. (1988) A discrete element 3' of human immunodeficiency virus 1 (HIV-1) and HIV-2 mRNA initiation sites mediates transcriptional

- activation by an HIV transactivator. *Mol. Cell. Biol.*, 8, 2555-2561.
- Jeang, K.T., Berhout, B. and Dropulic, B. (1993) Effects of integration and replication on transcription of the HIV-1 long terminal repeat. *J. Biol. Chem.*, 268, 24940-24949.
- Jones, K.A. and Peterlin, B.M. (1994) Control of RNA initiation and elongation at the HIV-1 promoter. *Annu. Rev. Biochem.*, 63, 717-743.
- Koseki, S., Ohkawa, J., Yamamoto, R., Takebe, Y. and Taira, K. (1998) A simple assay system for examination of the inhibitory potential *in vivo* of decoy RNAs, ribozymes and other drugs by measuring the Tat-mediated transcription of a fusion gene composed of the long terminal repeat of HIV-1 and a gene for luciferase. *J. Controlled release*, 53, 159-173.
- Kubota, S., Endo, S.-I., Maki, M. and Hatanaka, M. (1988) Role of cysteine-rich region of HIV Tat protein in its *trans*-activation ability. *Virus Genes*, 2, 113-118.
- Lazinski, D., Grzadzielska, E. and Das, A. (1989) Sequence-specific recognition of RNA hairpins by bacteriophage antiterminators requires a conserved arginine-rich motif. *Cell*, 59, 207-218.
- Leung, D.W., Chen, E. and Goeddel, D.V. (1989) A method for random mutagenesis of a defined DNA segments using a modified polymerase chain reaction. *J. Methods Cell and Mol. Biol.*, 1, 11-15.
- Liszewicz, J., Rappaport, J. and Dhar, R. (1991) Tat-regulated production of multimerized TAR RNA inhibits HIV-1 gene expression. *New Biol.*, 3, 82-89.
- Liszewicz, J., Sun D., Smythe, Lusso, P., Lori, F., Louie A., Markham, P., Rossi, J., Reitz, M. and Gallo, R.C. (1993) Inhibition of human

- immunodeficiency virus type I replication by regulated expression of a polymeric Tat activation response RNA decoy as a strategy for gene therapy in AIDS. *Proc. Natl. Acad. Sci. U.S.A.*, 90, 8000-8004.
- Long, K.S. and Crothers, D.M. (1995) Interaction of human immunodeficiency virus type 1 Tat-derived peptides with TAR RNA. *Biochemistry*, 34, 8885-8895.
- Marciniak, R.A., Calnan, B.J., Frankel, A.D. and Sharp, P.A. (1990) HIV-1 Tat protein *trans*-activates transcription *in vitro*. *Cell*, 63, 791-802.
- Osborne, S.E. and Ellington, A.D. (1997) Nucleic acid selection and the challenge of combinatorial chemistry. *Chem. Rev.*, 97, 349-370.
- Peterline, B.M., Luciw, P.A., Barr, P.J. and Walker, M.D. (1986) Elevated levels of mRNA can account for the *trans*-activation of human immunodeficiency virus. *Proc. Natl. Acad. Sci. U.S.A.*, 83, 9734-9738.
- Piotto, M., Saudek, V. and Sklenar, V. (1992) Gradient-tailored excitation for single-quantum NMR spectroscopy of aqueous solution. *J. Biomol. NMR*, 2, 661-665.
- Potyrailo, R.A., Conrad, C.R., Ellington, A.D. and Hieftje, G.M. (1998) Adapting selected nucleic acid ligands (aptamers) to biosensors. *Anal. Chem.*, 70, 3419-3425.
- Puglisi, J.D., Tan, R., Calnan B.J., Frankel, A.D. and Williamson, J.R. (1992) Conformation of the TAR RNA-arginine complex by NMR spectroscopy. *Science*, 257, 76-80.
- Puglisi, J.D., Chen, L., Frankel, A.D. and Williamson, J.R. (1993) The role of RNA structure in arginine recognition of TAR RNA. *Proc. Natl. Acad. Sci. U.S.A.*, 90, 3680-3684.



- Rhim, H. and Rice, A.P. (1993) TAR RNA binding properties and relative transactivation activities of human immunodeficiency virus type 1 and 2 Tat proteins. *J. Virol.*, 67, 1110-1121.
- Rice, A.P. and Mathews, M.B. (1988) Transcriptional but not translational regulation of HIV-1 by the Tat gene product. *Nature*, 332, 551-553.
- Rosen, C.A., Sodroski J.G. and Haseltine W.A. (1985) The location of *cis*-acting regulatory sequences in human T cell lymphotropic virus type III. *Cell*, 41, 813-823.
- Rounseville, M.P. and Kumar, A. (1992) Binding of a host cell nuclear protein to the stem region of human immunodeficiency virus type 1 *trans*-activation-response RNA. *J. Virol.*, 66, 1688-1694.
- Rounseville, M.P., Lin, H.-C., Agbottah E., Shukla, R.R., Rabson, A.B. and Kumar, A. (1996) Inhibition of HIV-1 replication in viral mutants with altered TAR RNA stem structures. *Virol.*, 216, 411-417.
- Sheline, C.T., Milocco, L.H. and Jones, K.A. (1991) Two distinct nuclear transcription factors recognize loop and bulge residues of the HIV-1 TAR RNA hairpin. *Genes Dev.*, 5, 2508-2520.
- Sodroski, J., Patarca, R. Rosen, C., Wong-Staal, F. and Haseltine, W.A. (1985) Location of the *trans*-acting region on the genome of human T-cell lymphotropic virus type III. *Science*, 229, 74-77.
- Sullenger, B.A., Gallardo, H.F., Ungers, G.E. and Gilboa, E. (1990) Overexpression of TAR sequences renders cells resistant to human immunodeficiency virus replication. *Cell*, 63, 601-608.
- Tao, J., Chen, L. and Frankel, A.D. (1997) Dissection of the proposed base triple in human immunodeficiency virus TAR RNA indicates the importance of the Hoogsteen interaction. *Biochemistry*, 36, 3491-3495.

- Tuerk, C. and MacDougall-Waugh, S. (1993) *In vitro* evolution of functional nucleic acids: high-affinity RNA ligands of HIV-1 proteins. *Gene*, 137, 33-39.
- Urvil, P.T., Kakiuchi, N., Zhou, D.-M., Shimotohno, K., Kumar, P.K.R. and Nishikawa, S. (1997) Selection of RNA aptamers that bind specifically to the NS3 protease of hepatitis C virus. *Eur. J. Biochem.*, 248, 130-138.
- Vaishav, Y.N. and Wong-Staal, F. (1991) The biochemistry of AIDS. *Annu. Rev. Biochem.*, 60, 577-630.
- Verhoef, K., Tijms, and Berkhout, B. (1997) Optimal Tat-mediated activation of the HIV-1 LTR promoter requires a full-length TAR RNA hairpin. *Nucleic Acids Res.*, 19, 6169-6176.
- Weeks, K.M., Ampe, C., Schults, S.C., Steitz, T.A. and Crothers, D.M. (1990) Fragments of the HIV-1 Tat protein specifically bind to TAR RNA. *Science*, 249, 1281-1285.
- Weeks, K.M. and Crothers, D.M. (1991) RNA recognition by Tat-derived peptides: interaction in the major groove? *Cell*, 66, 577-588.
- Wei, P., Garber, M.E., Fang, S.-M., Fischer, W.H. and Jones, K.A. (1998) A novel CDK-9-associated C-type cyclin interacts directly with HIV-1 Tat and mediates its high-affinity, loop-specific binding to TAR RNA. *Cell*, 92, 451-462.
- Wu-Baer, F., Lane, W.S. and Gaynor R.B. (1995) The cellular factor TRP-185 regulates RNA polymerase II binding to HIV-1 TAR RNA. *EMBO J.*, 14, 5995-6009.
- Wu-Baer, F., Lane, W.S. and Gaynor, R.B. (1996) Identification of a group of cellular factors that stimulates the binding of RNA polymerase II and TRP-185 to human immunodeficiency virus 1 TAR RNA. *J. Biol. Chem.*, 271, 4201-4208.

- Yamada, O., Kraus, G., Luznik, L., Yu, M. and Wong-Staal, F. (1996) A chimeric human immunodeficiency virus type 1 minimal rev responsive element-ribozyme molecules exhibit dual antiviral function and inhibits cell-cell transmission. *J. Virol.*, 70, 1596-1601.
- Yamamoto, R., Koseki, S., Ohkawa, J., Murakami, K., Nishikawa, S., Taira, K. and Kumar, P.K.R. (1997) Inhibition of transcription by the TAR RNA of HIV-1 in a nuclear extract of HeLa cells. *Nucleic Acids Res.*, 25, 3445-3450.
- Yamamoto, R., Murakami, K., Taira, K. and Kumar, P.K.R. (1998) Isolation and characterization of an RNA that binds with high affinity to Tat protein of HIV-1 from a completely random pool of RNA. *Gene Ther. Mol. Biol.*, 1, 451-466.
- Yu, M., Poeschla, E. and Wong-Staal, F. (1994) Progress towards gene therapy for HIV infection. *Gene Ther.*, 1, 13-26.
- Yuyama, N., Ohkawa, J., Koguma, T., Shirai, M. and Taira, K. (1994) A multifunctional expression vector for an anti-HIV-1 ribozyme that produces a 5'- and 3'- trimmed *trans*-acting ribozyme, targeted against HIV-1 RNA, and *cis*-acting ribozymes that are designed to bind to and thereby sequester *trans*-activator proteins such as Tat and Rev. *Nucleic Acids Res.*, 22, 5060-5067.
- Zhou, Q. and Sharp, P.A. (1996) Tat-SF-1: Cofactor for stimulation of transcriptional elongation by HIV-1 Tat. *Science*, 274, 605-610.
- Zuker, M. (1989) On finding all suboptimal foldings of an RNA molecule. *Science*, 244, 48-52.