

ACKNOWLEDGMENTS

The author wishes to thank the Japanese Ministry of Education (MONBUSHO) for giving her the opportunity to study in University of Tsukuba, Japan.

This work would not have been possible without the warm support, encouragements and invaluable guidance of her research advisers Dr. Hiroshi Mizuno and Prof. Masatoshi Matsumura.

The author deeply indebted to Dr. Zui Fujimoto and to Dr. Kenji Takase for sharing their knowledge and experience, for the constructive suggestions and assistance in her research as well as for the critical revising all papers and related manuscripts of the author.

The author gratefully acknowledge Drs. Noriyoshi Sakabe, Nobuhisa Watanabe, Mamoru Suzuki and Noriyuki Igarashi for the use of the Weissenberg camera detector system at the Photon Factory.

The author would like appreciate cooperative, positive environment and favorable facilities for the research in the laboratory of Protein Engineering of NIAR. She wishes to thank the students from Nagaoka University. Shi is indebted in many aspects to all previous and present members of the laboratory.

The authors gratitude to Dr. Pi-Chao Wang, Mrs. Irene Abe and to all members in the Bioprocess Engineering Laboratory, Institute of Applied Biochemistry, University of Tsukuba for warm support and positive attitude in all aspects.

Finally, to the author's mother, this work is a testimony to her untarnished support and encouragements.