

References

- 1) Alsaadawi, I. S., J. K. Al-Uquili, M. Al-Hadithy and A. I. Alrubeaa 1985. Effect of gamma irradiation on allelopathic potential of *Sorghum bicolor* against weeds and nitrification. J. Chem. Ecol. 11(12) 1735-1745.
- 2) Alsaadawi, I. S., F. A. K. Sakeri, S. M. Al-Dulaimy 1990. Allelopathic inhibition of *Cynodon dactylon* (L.) Pers. and other plant species by *Euphorbia prostrata* L. J. Chem. Ecol. 16(9) : 2746-2754.
- 3) An, M., J. E. Pratley and T. Haig 1997. Phytotoxicity of vulpia residues: I. Investigation of aqueous extracts. J. Chem. Ecol. 23(8) 1979-1995.
- 4) Anaya, A. I., B. E. Hernandez-Bautista, H. R. Pelayo-Benavides, M. Calera and E. Fernandez-Luiselli 1995. Allelopathy in Mexican plants : More recent studies. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 224-241.
- 5) Anderson, I. C. and R.M. Cruse 1995. Tillage and allelopathic aspects of the corn-soybean rotation effect. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 184-192.
- 6) Ayeni, A. O., D. T. Lordbanjou and B. A. Majek 1997a.

Tithonia diversifolia (Mexican sunflower) in south-western Nigeria : occurrence and growth habit. Weed Res. 37 : 443-449.

- 7) Ayeni, A. O., S. O. Agbato and B. A. Majek 1997b. Seed depth influence on Mexican sunflower (*Tithonia diversifolia*) emergence and control. Weed Technology, 11 : 417-427.
- 8) Baruah, N. C., R. P. Sharma, K. P. Madhusudanan and G. Thyagarajan 1979. Sesquiterpene lactones of *Tithonia diversifolia*. Stereochemistry of the tagitininins and related compounds. J. Org. Chem. 44(11) : 1831-1835.
- 9) Baruah, N. C., J. C. Sarma. N. C. Barua, S. Sarma and R. P. Sharma 1994. Germination and growth inhibitory sesquiterpene lactones and a flavone from *Tithonia diversifolia*. Phytochemistry, 36(1) : 29-36.
- 10) Bewick, T. A., D. G. Shilling, J. A. Dusky and D. Williams 1994. Effects of celery (*Apium graveolens*) root residue on growth of various crops and weeds. Weed Technology, 8 : 625-629.
- 11) Blum, U. 1995. The value of model plant-microbe-soil systems for understanding processes associated with allelopathic interaction : One example. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 127-131.
- 12) Blum, U. 1998. Effects of microbial utilization of phenolic

acids and their phenolic acid breakdown products on allelopathic interactions. J. Chem. Ecol. 24(4) : 685-708.

- 13) Blum, U., T. M. Gerig, A. D. Worsham and L. D. King 1993. Modification of allelopathic effects of *p*-coumaric acid on morning glory seedling biomass by glucose, methionine and nitrate. J. Chem. Ecol. 19(12) : 2791-2811.
- 14) Bordoloi, M., N. C. Barua and A. C. Ghosh 1996. An artemisinic acid analogue from *Tithonia diversifolia*. Phytochemistry, 41(2) : 557-559.
- 15) Chase, W. R., M. G. Nair and S. K. Mishra 1991. 2,2-oxo-1,1-azobenzene microbial transformation of rye (*Secale cereale* L.) allelochemical in field soils by *Acinetobacter calcoaceticus* : III. J. Chem. Ecol. 17(8) : 1575-1584.
- 16) Chaves, N., J. C. Escudero and C. Gutierrez-Merino 1997. Role of ecological variables in the seasonal variation of flavonoid content of *Cistus ladaifer* exudate. J. Chem. Ecol. 23(3) : 579-603.
- 17) Cheam, A., J. Peirce, L. Martin, S. Lee and B. Rayner 1995. Allelopathy in crop and pasture management with special reference to goosefoot (*Chenopodium pumilio*). In proceedings I (A) 15th Asian-Pacific Weed Science Society Conference, Tsukuba, Japan, pp. 305-310.
- 18) Cheng, H. H. 1995. Characterization of mechanisms of allelopathy : Modeling and experimental approaches. In "Allelopathy : Organisms, Processes, and Applications" ed. by

- Inderjit, K. M. M., Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 132-141.
- 19) Choesin, D. N. and R. E. J. Boerner 1991. Allyl isothiocyanate and the allelopathic potential of *Brassica napus* (Brassicaceae). Amer. J. Bot. 78(8) : 1083-1090.
- 20) Chowdhury, P. K., N. C. Barua, R. P. Sharma and G. Thyagarajan 1980. Structure of deacetyltagitinin (Tagitin E). J. Org. Chem. 45 : 535-536.
- 21) Chowdhury, P. K., N. C. Barua, R. P. Sharma, J. N. Barua, W. Herz and K. Watanabe 1983. Cyclotagitin C and its transformation. J. Org. Chem. 48 : 732-738.
- 22) Chou, C. H. 1987. Allelopathy in subtropical vegetation and soils in Taiwan. In "Allelochemicals : Role in agriculture and forestry" ed. by G. R. Waller, American Chemical Society, Washington, D.C., pp. 102-117.
- 23) Chou, C. H. 1995a. Allelopathy and sustainable agriculture. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 211-223.
- 24) Chou, C. H. 1995b. Allelopathic compounds as naturally occurring herbicides. In proceedings I (A) 15th Asian-Pacific Weed Science Society Conference, Tsukuba, Japan, pp. 154-159.
- 25) Chung, I. M. and D. A. Miller 1995a. Differences in autotoxicity among seven alfalfa cultivars. Agron. J. 87 : 596-

600.

- 26) Chung, I. M. and D. A. Miller 1995b. Effect of alfalfa plant and soil extracts on germination and growth of alfalfa. *Agron. J.* 87 : 762-767.
- 27) Chung, I. M. and D. A. Miller 1995c. Natural herbicide potential of alfalfa residue on selected weed species. *Agron. J.* 87 : 920-925.
- 28) Cox, L., A. Walker, M. C. Hermosin and J. Cornejo 1996. Measurement and stimulation of movement of thaizafluron, cropyralid and metamitron in soil columns. *Weed Res.* 36(5) : 419-429.
- 29) Dalton, B. R., U. Blum and S. B. Weed 1983. Allelopathic substances in ecosystems : Effectiveness of sterile soil components in altering recovery of ferulic acid. *J. Chem. Ecol.* 9(8) : 1185-1201.
- 30) Dietz, H, T. Steinlein, P. Winterhalter and I. Ullmann 1996. Role of allelopathy as a possible factor associated with the rising dominance of *Bunias orientalis* L. (Brassicaceae) in some native plant assemblages. *J. Chem. Ecol.* 22(10) : 1797-1811.
- 31) Dolling, A., O. Zackrisson and M. C. Nilsson 1994. Seasonal variation in phytotoxicity of bracken (*Pteridium aquilinum* L. Kuhn). *J. Chem. Ecol.* 20(12) : 3163-3172.
- 32) Duke, S. O. and H. K. Abbas 1995. Natural products with potential use as herbicides. In "Allelopathy : Organisms,

Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 348-362.

- 33) Dutta, P., R. P. Chaudhuri and R. P. Sharma 1993. Insect feeding deterrents from *Tithonia diversifolia* (Hemsl.) Gray. J. Environ. Biol. 14(1) : 27-33.
- 34) Einhellig, F. A. 1987. Interactions among allelochemicals and other stress factors of the plant environment. In "Allelochemicals : Role in agriculture and forestry" ed. by G. R. Waller, American Chemical Society, Washington, D.C., pp. 343-370.
- 35) Einhellig, F. A. 1995a. Allelopathy : Current status and future goals. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American chemical Society, Washington, D.C., pp. 1-24.
- 36) Einhellig, F. A. 1995b. Mechanism of action of allelochemicals in allelopathy. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American chemical Society, Washington, D.C., pp. 96-116.
- 37) Einhellig, F. A. 1996. Interactions involving allelopathy in cropping systems. Agron. J. 88 : 886-893.
- 38) Einhellig, F. A., M. S. Muth and M. K. Schon 1985. Effect of allelochemicals on plant-water relationships. In "The

- chemistry of allelopathy biochemical interactions among plants" ed. by A. C. Thompson, American Chemical Society, Washington, D.C., pp. 179-196.
- 39) Elakovich, S. D. and J. W. Wooten 1995. Allelopathic woody plants, Part I : *Abies alba* through *Lyonia lucida*. *Allelopathy J.* 2(2) : 117-146.
 - 40) Elakovich, S. D. and J. W. Wooten 1996. Allelopathic woody plants, Part II : *Mabea* through *Zelkova*. *Allelopathy J.* 3(1) : 9-32.
 - 41) Eskelsen, S. R. and G. D. Crabtree 1995. The allelopathy in buckwheat (*Fagopyrum sagittatum*) inhibition of Canada thistle (*Cirsium arvense*). *Weed Sci.* 43 : 70-74.
 - 42) Harada, J. 1994. Utilization of weeds for weed control. *Farming Japan*, 28(3) : 10-16.
 - 43) Haugland, E. and L. O. Brandsaeter 1996. Experiments on bioassay sensitivity in the study of allelopathy. *J. Chem. Ecol.* 22(10) : 1845-1859.
 - 44) Hegde, R. S. and D. A. Miller 1990. Allelopathy and autotoxicity in alfalfa : Characterization and effects of preceding crops and residue incorporation. *Crop Sci.* 30(6) : 1255-1259.
 - 45) Heissey, R. M. 1990. Evidence for allelopathy by tree-of-heaven (*Ailanthus altissima*). *J. Chem. Ecol.* 16(6) : 2039-2055.
 - 46) Hoffman, M. L., L. A. Weston, J. C. Snyder and E. E. Regnier 1996a. Separating the effects of sorghum (*Sorghum bicolor*)

- and rye (*Secale cereale*) root and shoot residues on weed development. Weed Sci. 44(2) : 402-407.
- 47) Hoffman, M. L., L. A. Weston, J. C. Snyder and E. E. Regnier 1996b. Allelopathic influence of germinating seeds and seedlings of cover crops on weed species. Weed Sci. 44(3) : 579-584.
- 48) Horie, H., M. Nemoto and N. Nishimura 1989. Dynamics of allelochemicals from broom (*Cytisus scoparius*) and their accumulation in soil. Weed Res., Japan, 34(1) : 47-56. (in Japanese).
- 49) Inderjit 1996a. Allelopathic potential of *Pluchea lanceolata* : Comparative studies of cultivated fields. Weed Sci. 44 : 393-396.
- 50) Inderjit 1996b. Plant phenolics in allelopathy. Bot. Rev. 62(2) : 186-202.
- 51) Inderjit and K. M. M. Dakshini 1994a. Allelopathic effect of *Pluchea lanceolata* (Asteraceae) on characteristics of four soils and tomato and mustard growth. Am. J. Bot. 81(7) : 799-804.
- 52) Inderjit and K. M. M. Dakshini 1994b. Allelopathic potential of the phenolics from the roots of *Pluchea lanceolata*. Physiol. Plant. 92 : 571-576.
- 53) Inderjit and K. M. M. Dakshini 1995a. Allelopathic potential of an annual weed, *Polypogon monspeliensis*, in crops in India. Plant and Soil 173 : 251-257.
- 54) Inderjit and K. M. M. Dakshini 1995b. On laboratory bioassay

- in allelopathy. Bot. Rev. 61(1) : 28-44.
- 55) Inderjit and K. M. M. Dakshini 1996. Allelopathic potential of well water from *Pluchea lanceolata* infested cultivated fields. J. Chem. Ecol. 22(6) : 1123-1131.
- 56) Inderjit and A. U. Mallik 1996. The nature of interference potential of *Kalmia angustifolia*. Can. J. For. Res. 26 : 1899-1904.
- 57) Inderjit, S. Kaur and K. M. M. Dakshini 1996. Determination of allelopathic potential of a weed *Pluchea lanceolata* through a multifaceted approach. Can. J. Bot. 74 : 1445-1450.
- 58) Inderjit and R. D. Moral 1997. Is separating resource competition from allelopathy realistic?. Bot. Rev. 63(3) : 221-230.
- 59) Irons, S. M. and O. C. Brunside 1982. Competitive and allelopathic effects of sunflowers (*Helianthus annuus*). Weed Sci. 30 : 372-377.
- 60) Ismail, B. S. and L. S. Mah 1993. Effects of *Mikania micrathra* H. B. K. on germination and growth of weed species. Plant and Soil. 157 : 107-113.
- 61) Ito, I, K. Kobayashi and T. Yoneyama 1998. Fate of dehydromatricaria ester added to soil and its implications for the allelopathic effect of *Solidago altissima* L. Annals of Botany, 82 : 625-630.
- 62) Jaderund, A, O. Zackrisson and M. C. Nilsson 1996. Effects of bilberry (*Vaccinium myrtillus* L.) litter on seed germination

and early seeding growth of four boreal tree species. J. Chem. Ecol. 22(5) : 973-986.

- 63) Jimenez-Osornio, J. J. and S. R. Gliessman 1987. Allelopathic interference in a wild mustard (*Brassica campestris* L.) and broccoli (*Brassica oleracea* L. var. *italica*) intercrop agroecosystem. In "Allelochemicals : Role in agriculture and forestry" ed. by G. R. Waller, American Chemical Society, Washington, D.C., pp. 262-274.
- 64) Kainulainen, P., J. Oksanen, V. Palomaki, J. K. Hokopainen and T. Hokopainen 1992. Effect of drought and waterlogging stress on needle monoterpenes of *Picea abies*. Can. J. Bot. 70: 1613-1616.
- 65) Kiemnec, G. and L. Larson 1991. Germination and root growth of two noxious weeds as affected by water and salt stresses. Weed Technology, 5 : 612-615.
- 66) Kil, B. S. and K. W. Yun 1992. Allelopathic effects of water extracts of *Artemisia princeps* var. *orientalis* on selected plant species. J. Chem. Ecol. 18(1) : 39-51.
- 67) Kim, K. U. 1995. Possible utilization of plants and allelochemicals for weed control. In proceedings I (A) 15th Asian-Pacific Weed Science Society Conference, Tsukuba, Japan, pp. 292-299.
- 68) Kitou, M. and S. Yoshida 1993. Difference of phytotoxicity between undecomposed and microbially decomposed plant residues. Weed Res., Japan, 38(1) : 43-46.

- 69) Kobayashi, K. 1996. Performance of herbicide in soil and its phytotoxic activity. In Proceeding of Northeast Asian Area Weed Science Symposium of China, Korea and Japan. Northeast Agricultural University, China, pp. 119-122.
- 70) Kobayashi, K., M. Onoe and H. Sugiyama 1994. Thenylchlor concentration in soil water and its herbicidal activity. Weed Res., Japan, 39 (3) : 160-165.
- 71) Kobayashi, K., N. Nakamura, I. S. Shim and S. Nagatsuka 1996. Relationship of herbicidal activity of soil-applied mefenacet to its concentration in soil water and adsorption in soil. Weed Res., Japan, 41(2) : 98-102.
- 72) Kuo, Y. H. and C. H. Chen 1998. Sesquiterpenes from the leaves of *Tithonia diversifolia*. J. Natural Products, 61(6) : 827-828.
- 73) La Duke, J. C. 1982. Flavonoid chemistry and systematics of *Tithonia* (Compositae). Amer. J. Bot. 69(5) : 784-792.
- 74) Li, H. H., H. Nishimura, K. Hasegawa and J. Mizutani 1992a. Allelopathy of *Sasa cernua*. J. Chem. Ecol. 18(10) : 1785-1796.
- 75) Li, H. H., M. Urashima, M. Amano, L. Lajide, H. Nishimura, K. Hasegawa and J. Mizutani 1992b. Allelopathy of barnyardgrass (*Echinochloa crus-galli* L. Beauv. var. *crus-galli*). Weed Res., Japan, 37(2) :146-152.
- 76) Lisanework, N. and A. Michelsen 1993. Allelopathy in agroforestry systems: the effects of leaf extracts of *Cupressus lusitanica* and three *Eucalyptus* spp. on four Ethiopian crop.

Agroforestry Systems, 21 : 63-74.

- 77) Lovett, J. V. 1982. The effect of allelochemicals on crop growth and development. In "Chemical manipulation of crop growth and development" ed. by J. S. McLaren, Mackays of Chatham Ltd, UK, pp. 93-110.
- 78) Lovett J. V. and A. H. C. Hoult 1995. Allelopathy and self-defense in barley. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 184-192.
- 79) Lydon, J., J. R. Teasdale and P. K. Chen 1997. Allelopathic activity of annual warmwood (*Artemisia annua*) and the role of artemisinin. Weed Sci. 45 : 807-811.
- 80) Macias, F. A., R. M. Varela, A. Torres and J. M. G. Molinillo 1993. Potential allelopathic guaianolides from cultivar sunflower leaves, var. SH-222. Phytochemistry, 34(3) : 669-674.
- 81) Macias, F. A. 1995. Allelopathy in the search for natural herbicide models. In "Allelopathy : Organism, Processes and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 310-329.
- 82) Mandava, N. B. 1985. Chemistry and Biology of allelopathic agents. In "The chemistry of allelopathy biochemical interactions among plants" ed. by A. C. Thompson, American

Chemical Society, Washington, D.C., pp. 33-54.

- 83) Martin, L. D. and A. E. Smith 1994. Allelopathic potential of some warm-season grasses. *Crop Protection*, 13(5) : 388-392.
- 84) Mason-Sedun, W and R. S. Jessop 1988. Differential phytotoxicity among species and cultivars of the genus *Brassica* to wheat. *Plant and Soil*, 107 : 69-80.
- 85) May, F. E. and J. E. Ash 1990. An assessment of the allelopathic potential of *Eucalyptus*. *Aust. J. Bot.* 38 : 245-254.
- 86) Menelaou, M. A., J. D. Weidenhamer, G. B. Williamson, F. R. Fronczek, H. D. Fischer, L. Quijano and N. H. Fischer 1993. Diterpenes from *Chrysoma pauciflosculosa* : Effect on Florida sandhill species. *Phytochemistry*, 34 : 97-105.
- 87) Mersie, W. and M. Singh 1987. Allelopathic effect of partenium (*Partenium hysterophorus* L.) extract and residue on some agronomic crops and weeds. *J. Chem. Ecol.* 13(7) : 1739-1747.
- 88) Miller, D. A. 1996. Allelopathy in forage crop systems. *Agron. J.* 88 : 854-859.
- 89) Molisch, H. 1937. Der Einfluss einer Pflanze auf die andere-Allelopathie. Jena, Germany : Gustav Fischer.
- 90) Nair, M. G., C. J. Whiteneck and A. R. Putnam 1990. 2,2-Oxo-1,1-azobenzene, a microbially transformed allelochemical from 2,3-benzoxazolinone : I. *J. Chem. Ecol.* 16(2) : 353-364.
- 91) Nakamura, N., K. Kobayashi, I. S. Shim and S. Nagatsuka 1996. Influence of soil organic matter content on mefenacet

- concentration in soil water and the phytotoxic activity. Weed Res., Japan, 41(4) : 339-343.
- 92) Nakamura, N. and M. Nemoto 1993. Allelopathic potential of *Eupatorium odoratum* in abandoned shifting cultivation fields in the tropics. Weed Res., Japan, 38(2) : 103-108. (in Japanese).
- 93) Nanakorn, W. 1995. Plant in the Queen Sirikit Botanic Garden. Chiang Mai , Thailand. 122 p. (in Thai).
- 94) Nelson, C. J. 1996. Allelopathy in cropping systems. Agron. J. 88 : 991-996.
- 95) Nilsson, M. C., C. Gallet and A. Wallstedt 1998. Temporal variability of phenolics and batatasin-III in *Empetrum hermaphoditum* leaves over an eight-year period : interpretations of ecological function. Oikos, 81 : 6-16.
- 96) Ogawa, K, S. Phetchawee and O. Suriyapan 1980. Field observations and laboratory analyses of upland soils in Thailand. Tropical Agriculture Research Center, Tsukuba, Japan, 290 p.
- 97) Olofsdotter, M. and D. Navarez 1995. Approaches in rice allelopathy research. In proceedings I (A) 15th Asian-Pacific Weed Science Society Conference, Tsukuba, Japan, pp. 315-320.
- 98) Pandey, D. K., L. P. Kauraw and V. M. Bhan 1993. Inhibitory effect of pathenium (*Pathenium hysterophorus* L.) residue on growth of water hyacinth (*Eichhornia crassipes* Mart Solms)

- II. Relative effect of flower, leaf, stem and root residue. J. Chem. Ecol. 19 : 2663-2670.
- 99) Penuelas, J. and J. Llusia 1997. Effects of carbon dioxide, water supply, and seasonality on terpene content and emission by *Rosmarinus officinalis*. J. Chem. Ecol. 23(4) : 979-993.
- 100) Pereira, P. S., D. A. Dias, W. Vichnewski, A. M. T. T. Nasi and W. Herz 1997. Sesquiterpene lactones from Brazilian *Tithonia diversifolia*. Phytochemistry, 45(7) : 1445-1448.
- 101) Pool, C. F. and D. D. Toit 1995. Leaching depth of imazethabenz methyl and chlosulfuron+metsulfuron methyl in different soils. Applied Plant Science, 9(2) : 43-47.
- 102) Przepiorkowski, T. and S. F. Gorski 1994. Influence of rye (*Secale cereale*) plant residues on germination and growth of three triazine resistant and susceptible weeds. Weed Technology, 8(4) : 744-747.
- 103) Putnam, A. R. 1985. Allelopathic research in agriculture : Past highlights and potential. In "The chemistry of allelopathy biochemical interactions among plants" ed. by A. C. Thompson, American Chemical Society, Washington, D.C., pp.1-8.
- 104) Putnam, A. R. and C. S. Tang 1986. Allelopathy : State of the science. In "The science of allelopathy" ed. by A. R. Putnam and C. S. Tang, John Wiley, New York, pp.1-9.
- 105) Radanachales, T. and J. F. Maxwell 1992. List of weeds report in Thailand. Faculty of Agriculture, Chiang Mai University,

- Thailand. 138 p. (in Thai).
- 106) Rice, E. L. 1984. Allelopathy. Second edition, Academic Press, Inc., U.S.A. 422 p.
- 107) Rice, E. L. 1987. Allelopathy : An overview. In "Allelochemicals : Role in agriculture and forestry" ed. by G. R. Waller, American Chemical Society, Washington, D.C., pp. 8-22.
- 108) Rizvi, S. L. H. and V. Rizvi 1987. Improving crop productivity in India : Role of Allelochemicals. In "Allelochemicals : Role in agriculture and forestry" ed. by G. R. Waller, American Chemical Society, Washington, D.C., pp. 69-75.
- 109) Sahid, I. B. and J. B. Sugau 1993. Allelopathic effect of lantana (*Lantana camara*) and Siam weed (*Chromolaena odorata*) on selected crops. Weed Sci. 41 : 303-308.
- 110) Schmidt, S. K. 1988. Degradation of juglone by soil bacteria. J. Chem. Ecol. 14 : 1561-1571.
- 111) Schuster, A., S. Stokes, F. Papastergiou, V. Castro, L. Poveda and J. Jakupovic 1992. Sesquiterpene lactones from two *Tithonia* species. Phytochemistry. 31(9) : 3139-3141.
- 112) Shilling, D. G., Dusky J. A., M. A. Mossler and T. A. Bewick 1992. Allelopathic potential of celery residues on lettuce. J. Amer. Soc. Hort. Sci. 117(2) : 308-312.
- 113) Tang, C. S., W. F. Cai, K. Kohl and R. K. Nishimoto 1995. Plant stress and allelopathy. In " Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M.

- Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 142-157.
- 114) Teasdale, J. R. 1993. Interaction of light, soil moisture, and temperature with weed suppression by hairy vetch residue. *Weed Sci.* 41 : 46-51.
- 115) Vaughn, S. F. and R. A. Boyston 1997. Volatile allelochemicals released by crucifer green manures. *J. Chem. Ecol.* 23(9) : 2107-2116.
- 116) Waller, G. R., C. S. Cheng, C. H. Chou, D. Kim, C. F. Yang, S. C. Huang and Y. F. Lin 1995. Allelopathic activity of naturally occurring compounds from mung beans (*Vigna radiata*) and their surrounding soil. In "Allelopathy : Organisms, Processes, and Applications" ed. by Inderjit, K. M. M. Dakshini and F. A. Einhellig, American Chemical Society, Washington, D.C., pp. 242-257.
- 117) Wardle, D. A., K. S. Nicholson and M. Ahmed 1992. Comparison of osmotic and allelopathic effects of grass leaf extracts on grass seed germination and radicle elongation. *Plant and Soil*, 140 : 315-319.
- 118) Wardle, D. A., K. S. Nicholson, and A. Rahman 1996. Use of a comparative approach to identify allelopathic potential and relationship between allelopathy bioassays and "competition" experiments for ten grassland and plant species. *J. Chem. Ecol.* 22(5) : 933-948.
- 119) Weidenhamer, J. D. 1996. Distinguishing resource

- competition and chemical interference : overcoming the methodological impasse. Agron. J. 88 : 866-875.
- 120) Weston, L. A. 1996. Utilization of allelopathy for weed management in agroecosystems. Agron. J. 88 : 860-866.
- 121) Whittaker, R. H. and P. P. Feeny 1971. Allelochemicals : Chemical interactions between plant species. Science, 171: 757-770.
- 122) Williams, R. D. and R. E. Hoagland 1982. The effect of naturally phenolic compounds on seed germination. Weed Sci. 30 : 206-212.
- 123) Williamson, G. B. and J. D. Weidenhamer 1990. Bacterial degradation of juglone : Evidence against allelopathy. J. Chem. Ecol. 16(5) : 1739-1742.
- 124) Willis, R. J. 1994. Terminology and trends in allelopathy. Allelopathy J. 1(1) : 6-28.
- 125) Yun, K. W. and B. S. Kil 1992. Assessment of allelopathic potential in *Artemisia princeps* var. *orientalis* residues. J. Chem. Ecol. 18(11) : 1933-1940.
- 126) Zhang, Q. 1993. Potential role of allelopathy in soil and the decomposing root of Chinese-fir replant woodland. Plant and Soil, 151 : 205-210.
- 127) Zungsontiporn, S. 1995. Allelopathic effect of some tropical plants. Ph.D. dissertation, Tokyo University of Agriculture and Technology. 125 p.
- 128) Zungsontiporn, S and J. Harada 1995. Alien weeds invading

follow land of shifting cultivation in the highland of northern Thailand. Weed Res., Japan, 40(Supplement 1) : 174-175.