

Bibliography

- [1] O. Aberth, Iteration methods for finding all zeros of a polynomial simultaneously, *Math. Comp.*, 27 (1973), pp. 339–344.
- [2] A. V. Aho, J. E. Hopcroft and J. D. Ullman, *The Design and Analysis of Computer Algorithms*, Addison-Wesley, Reading, MA, 1974.
- [3] G. Alefeld and J. Herzberger, *Introduction to Interval Computations*, Academic Press, New York, 1983.
- [4] L. Atanassova, On the simultaneous determination of the zeros of an analytic function inside a simple smooth closed contour in the complex plane, *J. Comput. Appl. Math.*, 50 (1994), pp. 99–107.
- [5] H. E. Bell, Gershgorin's theorem and the zeros of polynomials, *Amer. Math. Monthly*, 72 (1965), pp. 292–295.
- [6] D. A. Bini, L. Gemignani and B. Meini, Factorization of analytic functions by means of Koenig's theorem and Toeplitz computations, *Numer. Math.*, 89 (2001), pp. 49–82.
- [7] L. Brugnano and D. Trigiante, Polynomial roots: the ultimate answer? *Linear Algebra Appl.*, 225 (1995), pp. 207–219.
- [8] M. P. Carpentier and A. F. D. Santos, Solution of equations involving analytic function, *J. Comput. Phys.*, 45 (1982), pp. 210–220.
- [9] C. Carstensen and T. Sakurai, Simultaneous factorization of a polynomial by rational approximation, *J. Comput. Appl. Math.*, 61 (1995), pp. 165–178.

Bibliography

- [10] C. Carstensen, Inclusion of the roots of a polynomial based on Gershgorin's theorem, *Numer. Math.*, 59 (1991), pp. 349–360.
- [11] L. M. Delves and J. N. Lyness, A numerical method for locating the zeros of an analytic function, *Math. Comput.*, 21 (1967), pp. 543–560.
- [12] P. Dörfler and G. Schmeisser, Construction of unitary and normal companion matrices, *Linear Algebra Appl.*, 202 (1994), pp. 193–220.
- [13] I. Eble and M. Neher, Computation of validated bounds for Taylor coefficients of analytic functions with ACETAF, Preprint Nr. 01/18 des IWRMM, Universität Karlsruhe, 2001.
- [14] A. Edelman and H. Murakumi, Polynomial roots from companion matrix eigenvalues, *Math. Comp.*, 64 (1995), pp. 763–776.
- [15] M. Fiedler, Expressing a polynomial as the characteristic polynomial of a symmetric matrix, *Linear Algebra Appl.*, 141 (1990), pp. 265–270.
- [16] S. Fortune, Polynomial root finding using iterated eigenvalue computation, *Proc. ISSAC 2001*, Ontario (2001), pp. 121–128.
- [17] I. Gargantini and P. Henrici, Circular arithmetic and the determination of polynomial zeros, *Numer. Math.*, 18 (1972), pp. 305–320.
- [18] G. H. Golub and G. F. Van Loan, *Matrix Computations*, 2nd edition, The Johns Hopkins University Press, Baltimore, 1990.
- [19] E. Hansen, *Global Optimization Using Interval Analysis*, Monographs and Textbooks in Pure and Applied Mathematics, 165, Marcel Dekker, Inc., New York, 1992.
- [20] P. Henrici, *Applied and Computational Complex Analysis, Vol. I*, Wiley, New York, 1974.
- [21] V. Hribernig and H. J. Stetter, Detection and validation of clusters of polynomial zeros, *J. Symbolic Comput.*, 24 (1997), pp. 667–681.

Bibliography

- [22] M. Igarashi and T. Ypma, Relationships between order and efficiency of a class of methods for multiple zeros of polynomials, *J. Comput. Appl. Math.*, 60 (1995), pp. 101–113.
- [23] N. I. Ioakimidis, Quadrature methods for numerical Integration: Recent Developments, Software and Applications, P.Keast and G. Fairweather, eds., Reidel, Dordrecht, The Netherlands, 1987, pp. 61–82.
- [24] M. A. Jenkins and J. F. Traub, A three-stage variable shift iteration for polynomial zeros and its relation to generalized Ritz iteration, *Numer. Math.*, 14 (1970), pp. 252–263.
- [25] L. N. Koh-Trefethen, Pseudozeros of polynomials and pseudospectra of companion matrices, *Numer. Math.*, 68 (1994), pp. 403–425.
- [26] P. Kravanja, T. Sakurai and M. Van Barel, On locating clusters of zeros of analytic functions, *BIT*, 39 (1999), pp. 646–682.
- [27] U. W. Kulisch, *Advanced Arithmetic for the Digital Computer. Design of Arithmetic Units*, Springer-Verlag, Vienna, 2002.
- [28] R. Kumar, A fast algorithm for solving a Toeplitz system of equations, *IEEE Trans. Acoust. Speech Signal Process.*, 33 (1985), pp. 254–267.
- [29] T.-Y. Li, On locating all zeros of an analytic function within a bounded domain by a revised Delves/Lyness method, *SIAM J. Numer. Anal.*, 20 (1982), pp. 865–871.
- [30] F. Malek and R. Vaillancourt, A composite polynomial zerofinding matrix algorithm, *Comput. Math. Appl.*, 30 (1995), pp. 37–47.
- [31] F. Malek and R. Vaillancourt, Polynomial zerofinding iterative matrix algorithms, *Comput. Math. Appl.*, 29 (1995), pp. 1–13.
- [32] M. Marden, *Geometry of Polynomials*, AMS Mathematical Surveys 3, 1996.
- [33] J. M. McNamee, A bibliography on roots of polynomials, *J. Comput. Appl. Math.*, 47 (1993), pp. 391–394.

Bibliography

- [34] T. Miyakoda, Iterative methods for multiple zeros of a polynomial by clustering, *J. Comput. Appl. Math.*, 28 (1989), pp. 315–326.
- [35] R. E. Moore, *Interval Analysis*, Prentice-Hall, Englewood Cliffs, New Jersey, 1966.
- [36] R. E. Moore, *Methods and Applications of Interval Analysis*, SIAM Studies in Applied Mathematics, Philadelphia, 1979.
- [37] R. G. Mosier, Root neighbourhoods of a polynomial, *Math. Comput.*, 47 (1986), pp. 265–273.
- [38] M. Nakao and N. Yamamoto, *Numerical Computation with Guaranteed Accuracy*(in Japanese), Nippon-Hyoron-Sha, Tokyo, 1998.
- [39] M. Neher, Validated bounds for Taylor coefficients of analytic functions, *Reliab. Comput.*, 7 (2001), pp. 307–319.
- [40] M. Neher, Improved validated bounds for Taylor coefficients and for Taylor remainder series, *J. Comput. Appl. Math.*, 152 (2003), pp. 393–404.
- [41] M. Neher, The mean value form for complex analytic functions, *Computing*, 67 (2001), pp. 255–268.
- [42] A. Neumaier, Enclosing clusters of zeros of polynomials, *J. Comput. Appl. Math.*, 156 (2003), pp. 389–401.
- [43] A. Neumaier, An existence test for root clusters and multiple roots, *Z. Angew. Math. Mech.*, 68 (1988), pp. 256–257.
- [44] A. Neumaier, *Interval Methods for Systems of Equations*, Cambridge University Press, Cambridge, 1990.
- [45] N. Ohsako, T. Sakurai, H. Sugiura and T. Torii, A stable method for generating complex coefficient polynomial remainder sequence(in Japanese), *Trans. of Japan Society for Industrial and Applied Mathematics*, 5 (1995), pp. 241–255.

Bibliography

- [46] N. Ohsako, H. Sugiura and T. Torii, A stable extended algorithm for generating polynomial remainder sequence(in Japanese), *Trans. of Japan Society for Industrial and Applied Mathematics*, 7 (1997), pp. 227–255.
- [47] N. Ohsako, H. Sugiura and T. Torii, An extended algorithm by Givens Rotations for the polynomial remainder sequence(in Japanese), *Trans. of Information Processing Society of Japan*, 38 (1997), pp. 158–160.
- [48] S. Oishi, *Numerical Computation with Guaranteed Accuracy*(in Japanese), Corona Press, Tokyo, 2000.
- [49] V.Y. Pan, Solving a polynomial equation: some history and recent progress, *SIAM Review*, 39 (1997), pp. 187–220.
- [50] M. Petković, *Iterative Methods for Simultaneous Inclusion of Polynomial Zeros*, Springer-Verlag, Berlin, 1989.
- [51] M. S. Petković, C. Carstensen and M. Trajković, Weierstrass' formula and zero-finding methods, *Numer. Math.*, 69 (1995), pp. 353–372.
- [52] M. S. Petković and L. D. Petković, *Complex Interval Arithmetic and Its Applications*, Wiley-VCH Verlag Berlin GmbH, Berlin, 1998.
- [53] M. S. Petković, Inclusion methods for the zeros of analytic functions, in Computer Arithmetic and Enclosure Method, L. Atanassova and J. Herzberger, eds., North-Holland, 1992, pp. 319–328.
- [54] M. S. Petković and D. Herceg, Higher-order iterative methods for approximating zeros of analytic functions, *J. Comput. Appl. Math.*, 39 (1992), pp. 243–258.
- [55] M. S. Petković and Z. M. Marjanović, A Class of simultaneous methods for the zeros of analytic functions, *Comput. Math. Appl.*, 22 (1991), pp. 79–78.
- [56] J. L. Phillips, The triangular decomposition of Hankel matrices, *Math. Comp.*, 25 (1971), pp. 559–602.

Bibliography

- [57] H. Ratschek and J. Rokne, *New Computer Methods for Global Optimization*, Ellis Horwood Ltd., Chichester, Halsted Press, New York, 1988.
- [58] H. Ratschek and J. Rokne, *Computer Methods for the Range of Functions*, Ellis Horwood Ltd., Chichester, Halsted Press, New York, 1984.
- [59] J. Rokne and P. Lancaster, Complex interval arithmetic. Algorithm 86, *Comput. J.*, 18 (1975), pp. 83–86.
- [60] S. M. Rump, Ten methods to bound multiple roots of polynomials, *J. Comput. Appl. Math.*, 156 (2003), pp. 403–432.
- [61] S. M. Rump, INTLAB-INTerval LABoratory, <http://www.ti3.tu-harburg.de/~rump/intlab/index.html>.
- [62] S. M. Rump, INTLAB - INTerval LABoratory, In *Developments in Reliable Computing*, T.Csendes (Ed.), Kluwer, Dordrecht, Netherlands, 77–105 (1999).
- [63] T. Sakurai, P. Kravanja, H. Sugiura and M. Van Barel, An error analysis of two related quadrature methods for computing zeros of analytic functions, *J. Comput. Appl. Math.*, 152 (2003), pp. 467–480.
- [64] T. Sakurai and H. Sugiura, On factorization of analytic functions and its verification, *Reliab. Comput.*, 6 (2000), pp. 459–470.
- [65] T. Sakurai and H. Sugiura, Improvement of convergence of an iterative method for finding polynomial factors of analytic functions, *J. Comput. Appl. Math.*, 140 (2002), pp. 713–725.
- [66] T. Sakurai, T. Torii, N. Ohsako and H. Sugiura, A method for finding cluster of zeros of analytic function, *Proc. ICIAM'95*, Hamburg, (1996), pp. 515–516.
- [67] T. Sasaki, *Symbolic and Algebraic Computation*(in Japanese), Ohmsha, Tokyo, 1981.
- [68] G. Schmeisser, A real symmetric tridiagonal matrix with a given characteristic polynomial, *Linear Algebra Appl.*, 193 (1993), pp. 11–18.

Bibliography

- [69] B. T. Smith, Error bounds for zeros of a polynomial based upon Gershgorin's theorems, *J. Assoc. Comput. Mach.*, 17 (1970), pp. 661–674.
- [70] G. W. Stewart, On a companion operator for analytic functions, *Numer. Math.*, 18 (1971), pp. 26–43.
- [71] J. Stoer and E. Bulirsch, *Introduction to Numerical Analysis*, Springer-Verlag, New York, 1980.
- [72] H. Sugiura and T. Katou, Cdomain: Circular complex arithmetic system for determining the analyticity of analytic functions(in Japanese), The 2002 annual conference of the Japan Society for Industrial and Applied Mathematics, (2002).
- [73] T. Torii and T. Sakurai, Global method for the poles of analytic function by rational interpolant on the unit circle, *World Sci. Ser. Appl. Anal.*, 2 (1993), pp. 389–398.
- [74] W.F. Trench, An algorithm for the inversion of finite Hankel matrices, *J. Soc. Indust. Appl. Math.*, 13 (1965), pp. 1102–1107.
- [75] J. H. Wilkinson, *Rounding Errors in Algebraic Process*, Prentice-Hall, Englewood Cliffs, NJ, 1963.
- [76] J. H. Wilkinson, *The Algebraic Eigenvalue Problem*, Oxford, Clarendon Press, 1965.
- [77] J. C. Yakoubsohn, Finding a cluster of zeros of univariate polynomials, *J. Complexity*, 16 (2000), pp. 603–638.
- [78] T. Yamamoto, S. Kanno and L. Atanassova, Validated computation of polynomial zeros by the Durand-Kerner method, Topics in Validated Computations (Oldenburg, 1993), pp. 27–53, Stud. Comput. Math., 5, North-Holland, Amsterdam, 1994.
- [79] T. J. Ypma, Finding a multiple zero by transformations and Newton-like method, *SIAM Review*, 25 (1983), pp. 365–378.
- [80] Z. Zeng, On ill-conditioned eigenvalues, multiple roots of polynomials, and their accurate computations, MSRI Preprint 1998-048, 1998.