

参考文献

- [1] Baule G.M., McFee R. Detection of the magnetic field of the heart. Am Heart J 1963;66:95-96.
- [2] Baule G.M., McFee R. Theory of magnetic detection of the heart's electrical activity. J Appl Phys 1965;36:2066-2073.
- [3] Cohen D. Magnetic field around the torso: Production by electrical activity of the human torso. Science 1967;156:652-654.
- [4] Cohen D. A shielded facility for low level magnetic measurements. J Appl Phys 1967;38:1295-1296.
- [5] Cohen D., Edelsack A., Zimmerman J.E. Magnetocardiograms taken inside a shielded room with a superconducting point-contact magnetometer. Appl Phys Letters 1970;16:278-80.
- [6] Cohen D., McCaughan D. Magnetocardiograms and their variation over the chest in normal subjects. Am J Cardiol 1972;29:678-85.
- [7] Rosen A., Inouye G.T., Morse A.L. Magnetic recording of the heart's electrical activity with a cryogenic magnetometer. J Appl Phys 1971;42: 3682-3684.
- [8] Zimmerman J.E., Frederic N.V. Miniature

ultrasensitive superconducting magnetic gradiometer and its use in cardiography and other application. *Appl Phys Lett* 1971;19:16-19.

- [9] Cohen D. Magnetoencephalography; Evidence of magnetic fields produced by alpha-rhythm currents. *Science* 1968;161:784-786.
- [10] Cohen D. Magnetoencephalography; Detection of the brain's electrical activity with a superconducting magnetometer. *Science* 1972;175: 664-666.
- [11] Teyler T. J., Cuffin B.N., Cohen D. The visually evoked magnetoencephalogram. *Life Sci* 1975;16: 683-692.
- [12] Brenner D., Williamson S. J., Kaufman L. Visually evoked magnetic field of the human brain. *Science* 1975;190:480-482.
- [13] Reite M., Edrich J., Zimmerman J. T., Zimmerman J. E. Human magnetic auditory evoked fields. *Electroenceph Clin Neurophysiol* 1978;45:114-117.
- [14] Brenner D., Lipton J., Kaufman L., Williamson S. J. Somatically evoked magnetic fields of the human brain. *Science* 1975;199:81-83.
- [15] Hari R., Kaukoranta E., Reinkainen K., Mauno J. Neuromagnetic responses to noxious stimulation. *Biomagnetism*. Pergamon Press, 1985:pp359-363.

- [16] Chapman R.M., Ilmoniemi R.J., Barbanera S., Romani G.L. Selective localization of alpha brain activity with neuromagnetic measurement. *Electroenceph Clin Neurophysiol* 1984;58:569-572.
- [17] Barth D.S., Sutherling J., Engel Jr. J., Beatty J. Neuromagnetic localization of epileptiform spike activity in the human brain. *Science* 1982; 218:891-894.
- [18] Erne S.N., Hoke M., Lutkenhoner B., Pantew C., Scheer H.J. Brainstem auditory magnetic fields. *Biomagnetism*. Tokyo Denki University Press, 1988:pp158-161.
- [19] Kado H., Nakanishi M., Koyanagi M., Kasai N., Tndo T., Nakasato N. DC-SQUID magnetometry. A trial to detect the auditory evoked magnetic field of neural activity from brain stem. *Tokyo Denki University Press*, 1988:pp162-165.
- [20] Mizutani Y., Kuriki S. Somatically evoked magnetic fields in the vicinity of the neck. *IEEE Trans Biomed Eng* 1986;BME-33:510-516.
- [21] Wikswo Jr. J.P., Barach J.P., Freeman J.A. The magnetic field of a nerve impulse; First measurement. *Science* 1980;280:53-55.
- [22] Armstrong R.A., Janday B. A brief review of magnetic fields from the human visual system.

Ophthalmic Physiol Opt 1989;9:299-301.

- [23] Kveder M., Zupancic I., Lahajnar G., Blinc R., Suput D., Ailion D.C., Ganesan K., Goodrich C. Water proton NMR relaxation mechanisms in lung tissue. Magn Reson Med 1998;7:432-441.
- [24] Weng X.T., Yao J.C., Wang Y.J., Hu T.X., Yu Z.G., Lu Z.Y., Guo J.S. A study of magnetic fields in the lung of workers in an asbestos factory. Ind Health 1989;27:9-15.
- [25] Stahlhofen W., Moller W. Behavior of magnetic micro-particles in the human lung. Radiat Environ Biophys 1993;32:221-238.
- [26] Stahlhofen W., Moller W. In vivo and in vitro studies of the cellular defence system of the human lung. Toxicol Lett 1994;72:127-136.
- [27] Safnov Y.D., Pravotorov V.M., Lube V.M., Yakimenkov L.J. Method of recording the magnetic field of the heart (magnetocardiography). Bull Exp Biol 1967;64:1022-1024.
- [28] Cohen D., Lepeschkin E., Hosaka H., Massell B.F., Myers G. Abnormal patterns and physiological variations in magnetocardiograms. J Electrocardiol 1976;9:398-409.
- [29] Zimmerman J.E., Fredrick N.V. Design and operation of stable R-F biased superconducting

point-contact quantum devices and a note on properties of perfectly clean metal contact. *Appl Phys Lett* 1970;41:1572-1580.

- [30] Opfer J.E., Yeo Y.K., Pierce J.M., Rorden L.H. A superconducting second-derivative gradiometer. *IEEE Trans Magn* 1974;10:536-540.
- [31] Hosaka H., Cohen D., Cuffin B.N., Horacek B.M. The effect of torso boundaries on the magnetocardiogram. *J Electrocardiol* 1976; 9: 418-425.
- [32] Hosaka H., Cohen D. Visual determination of generators of the magnetocardiogram. *J Electrocardiol* 1976;9:426-432.
- [33] Saarinen M., Karp P.J., Kalita T.E. The magnetocardiogram in cardiac disorders. *Cardiovasc Res* 1974;8:82-834.
- [34] Sumi M., Takeuchi A., Katayama M., Fukuda Y., Nomura M., Fujino K., Murakami M., Nakaya Y., Mori H. Magnetocardiographic P waves in normal subjects and patients with mitral stenosis. *Jpn Heart J* 1986;27(5):621-628.
- [35] 藤野和也, 中屋 豊. 左室負荷疾患の心磁図に関する研究. *日本臨床生理誌* 1983;13:202-210.
- [36] Fujino K., Sumi K., Saito M., Murakami M., Higuchi T., Nakaya Y., Mori H.. Magnetocardiograms of

patients with left ventricular overloading recorded with a second-derivative SQUID gradiometer. J Electrocardiol 1984;17(3): 219-226.

- [37] 須見昌輝, 中屋 豊, 村上 昌. 心磁図法による心磁図 QRS 波に関する検討-特徴的波形の定量的検討と出現部位. 臨床と研究 1987;97:482-489.
- [38] Stroink G., Lant J., Elliott P. Discrimination between myocardial infarct and ventricular tachycardia patients using magnetocardiographic trajectory plots and iso-integral maps. J Electrocardiol 1992;25:129-141.
- [39] Brockmeier K., Comani S., Erne S.N., Di Luzio S., Pasquarelli A., Romani GL. Magnetocardiography and exercise testing. J Electrocardiol 1994;27: 137-142.
- [40] Lant J., Stroink G., ten Voorde B., Horacek B. M., Montague T. J. Complementary nature of electrocardiographic and magnetocardiographic data in patients with ischemic heart disease. J Electrocardiol 1990;23:315-322.
- [41] Nomura N., Nakaya Y., Saito K., Kishi F., Watatsuki T., Miyoshi H., Nishikado A., Bando S., Ito S., Nishitani H., Wada M., Fujita S., Tamura I. Noninvasive localization of accessory pathways by magnetocardiographic imaging. Clin Cardiol 1994;17:239-244.

- [42] 葛西直子, 国分俊典, 木戸カヤノ, 粟野直行, 芽根一夫. 心磁図計測による副伝導路の位置推定-カテーテルを用いた侵襲的検査との比較. 医用電子と生体工学 1994;32:231-237.
- [43] Katila T, Maniewski R, Makijarvi M, . On the accuracy of source localization in cardiac measurements. Phys Med Biol 1987;32:125-131.
- [44] Weismuller P., Abraham-Fuchs K., Killmann R., Richter P., Harer W., Hoher M., Kochs M., Eggeling T., Hombach V. Magnetocardiography: three-dimensional localization of the origin of ventricular late fields in the signal averaged magnetocardiogram in patients with ventricular late potentials. Eur Heart J 1993;14:61-68.
- [45] Oeff M, Burghoff M. Magnetocardiographic localization of the origin of ventricular ectopic beats. Pace 1994;17:517-522.
- [46] 野村昌弘, 中屋 豊. 心磁界計測による不整脈発生源の推定. 病態生理 1991;10(12):963-970.
- [47] 野村昌弘, 中屋 豊, 若槻哲三, 三好由貴子, 岸 史子, 斎藤 憲, 伊藤 進, 中安紀美子, 西谷 弘, 松崎健司, 上野慎一, 田村逸朗, 和田昌夫, 藤田 智, 高江 勉. 心臓磁界測定による心室性期外収縮の心室最早期興奮部位の推定. 心臓 1993;25:907-913.
- [48] Farrell D.E., Tripp J.H., Norgren R. Magnetic

study of the His-Purkinje conduction system in man.
IEEE Trans Biomed Eng 1980;BME-27:1366-1372.

- [49] Fenici R.R., Romini G.L., Barvanera S., Zeppilli P., Carelli P., Modena I. High resolution magnetocardiography: non-invasive investigation of the His-Purkinje system activity in man. J Ital Cardiol 1980;10:1366-1370.
- [50] MacAulay C.E., Stroink G., Horacek B.M. Analysis of MCG maps during the PR-interval. Med Biol Eng Comput 1985;23 suppl2:1479-1480.
- [51] Lamothe R., Stroink G. Orthogonal expansions: their applicability to signal extraction in electrophysiological mapping data. Med Biol Eng Comput 1991;29:522-528.
- [52] ten Voorde B.J., Peters M.J., Stroink G., van der Wieken L.R. High-resolution magnetic mapping of PR-interval phenomena of normal subjects. Med Biol Eng Comput 1988;26:130-135.
- [53] 粟野亥佐武, 村本敦子, 粟野直行, 古賀清毅. 超電導の応用と心磁図. 低温工学 1978;4:89-94.
- [54] 粟野亥佐武, 保坂光弘, 古賀清毅, 村松敦子, 粟野直行. 2次勾配 SQUID 磁束計の心磁図測定への応用. 医用電子と生体工学 1979;17:696.
- [55] 井深 丹, 内藤誠一, 細松春夫. SQUID 磁束計による心磁測定. 電子通信学会 EMCJ 1980;79:65-70.

- [56] 大道 久, 渥美和彦. Cryoelectronics と Magnetocardiography. 医学のあゆみ 1976;92:521.
- [57] 大道 久, 井深 丹, 渥美和彦. SQUID 磁束計による心磁図の計測. 医用電子と生体工学 1980;18:56-57.
- [58] 小谷 誠, 内川義則, 千代谷慶三, 斎藤健一. 残留磁界測定による肺内蓄積粉塵量の推定. 医用電子と生体工学 1981;19:442-445.
- [59] Nakaya. Y, Sumi M, Saito. K, Fujino K., Murakami M, Mori H. Analysis of current source of the heart using isomagnetic and vector arrow maps. Jpn Heart J 1983;25:701-711.
- [60] 斎藤京子, 中屋 豊. 正常心磁図 QRS 波に関する研究. 日本臨床生理誌 1983;13:191-201.
- [61] 村上 昌, 中屋 豊. 心筋梗塞症の心磁図 QRS 波に関する研究. 日本臨床生理誌 1983;14:14-25.
- [62] 中屋 豊, 須見昌輝, 斎藤京子, 村上 昌, 藤野和也, 坂東重信, 森 博愛. 二次勾配 SQUID system による心磁図記録方法の検討. 日本臨床生理誌 1983; 13:4-12.
- [63] Nomura M., Nakayasu K., Nakaya Y., Miyoshi Y., Wakatsuki T., Saito K., Bando S., Ito S. Single moving dipole obtained from magnetic field of the heart in patient with left ventricular hypertrophy. Clin Cardiol 1992;15:752-758.

- [64] 片山まり子, 野村昌弘, 中屋 豊. 側面心磁図 QRS 波に関する研究. 日本臨床生理誌 1988;18:269-283.
- [65] Wakai R.T., Wang M., Pedron S.L., Reid D.L., Martin C.B.Jr. Spectral analysis of antepartum fetal heart rate variability from fetal magnetocardiogram recordings. Early Hum Dev 1993;35:15-24.
- [66] Wakai R.T., Wang M., Martin C.B.Jr. Spatiotemporal properties of the fetal magnetocardiogram. Am J Obstet Gynecol 1994;170: 770-776.
- [67] Baffa O., Wakai R.T., Sousa P.L., Verzola R.M. Fetal heart rate monitoring by magnetocardiograms. Braz J Med Biol Res. 1995;28:1333-1337.
- [68] Rassi D., Lewis M.J. Power spectral analysis of the foetal magnetocardiogram. Physiol Meas. 1995;16:111-120.
- [69] Wakai R.T., Wang M., Leuthold A.C., Martin C.B. Foetal magnetocardiogram amplitude oscillations associated with respiratory sinus arrhythmia. Physiol Meas. 1995;16:49-54.
- [70] Kariniemi V., Hukkanen K. Quantification of fetal heart rate variability by magnetocardiography and direct electrocardiography Am J Obstet Gynecol 1997;128:526-530.

- [71] Wakai R.T., Leuthold A.C., Wilson A.D., Martin C.B. Association of fetal junctional rhythm and respiratory arrhythmia detected by magnetocardiography. *Pediatr Cardiol* 1997;18: 201-203.
- [72] Wakai R.T., Leuthold A.C., Martin C.B. Atrial and ventricular fetal heart rate patterns in isolated congenital complete heart block detected by magnetocardiography. *Am J Obstet Gynecol* 1998; 179:258-260.
- [73] Leuthold A., Wakai R.T., Martin C.B. Noninvasive in utero assessment of PR and QRS intervals from the fetal magnetocardiogram. *Early Hum Dev*. 1999 Apr;54:235-243.
- [74] Hamada H., Horigome H., Asaka M., Shigemitsu S., Mitsui T., Kubo T., Kandori A., Tsukada K. Prenatal diagnosis of long QT syndrome using fetal magnetocardiography. *Prenat Diagn*. 1999;19:677-680.
- [75] Van Leeuwen P., Lange S., Bettermann H., Gronemeyer D., Hatzmann W. Fetal heart rate variability and complexity in the course of pregnancy. *Early Hum Dev* 1999 Apr;54:259-269.
- [76] 保坂栄弘. Magnetocardiogram の基礎的特性の解析. 医用電子と生体工学 1976;14:466-472.

- [77] Nenonen J., Purcell C., Horacek B.M., Stroink G., Katila T. Magnetocardiographic functional localization using a current dipole in a realistic torso. *IEEE Trans Biomed Eng* 1991;38:658-664.
- [78] Taccardi B. Distributions of heart potentials of the thoracic surface. *Circ Res* 1963;12:341-352.
- [79] Lawrence E.W., Liebman J., Thomas C., Fraenkel R., Rudy Y. Electrocardiographic body surface potential maps of the QRS and T of normal young men. Qualitative description and selected quantifications. *J Electrocardiol* 1988;21: 121-136.
- [80] Tsukada K., Haruta Y., Adachi A., Ogata H., Komuro T., Ito T., Takeda Y., Kandori A., Noda Y., Terada Y., Mitsui T. Multichannel SQUID system detecting tangential components of the cardiac magnetic field. *Rev Science Inst* 1995;66:5085-5091.
- [81] Kandori A., Tsukada K., Haruta Y., Noda Y., Terada Y., Mitsui T., and Sekihara K. Reconstruction of two-dimensional current distribution from tangential MCG measurement. *Phys Med Biol* 1996; 41:1705-1716.
- [82] Tsukada K., Kandori A., Miyashita T., Sasabuchi H., Suzuki H., Kondo S., Komiya Y., Teshigawara K. A simplified super-conducting quantum interference device system to analyze vector components of a cardiac magnetic field.

Proceedings of the 20th annual international conference of the IEEE engineering in medicine and biology society. HongKong, 1998, Oct 29–Nov 1:1524–527.

- [83] Miyashita T., Kandori A., Tsukada K., Sato M., Tearda Y., Horigome H., Mitsui T. Construction of tangential vectors from normal cardiac magnetic field components. Proceedings of the 20th annual international conference of the IEEE engineering in medicine and biology society. HongKong, 1998, Oct 29–Nov 1.:520–523.
- [84] Morikawa M., Saito K., Takahashi M., Nakaya Y., Bando S., Mori H. Magnetocardiogram in right ventricular hypertrophy. Jpn Heart J 1982; 23-Suppl.:725–727.
- [85] 須見昌輝, 中屋 豊, 村上 昌, 福田善晴, 野村昌弘, 竹内明美, 片山まり子, 森 博愛. 心磁図法による右室負荷の診断—僧帽弁狭窄における検討. 循環器科 1987;21:373.
- [86] 須見昌輝, 中屋 豊, 村上 昌, 福田善晴, 野村昌弘, 竹内明美, 片山まり子, 森 博愛. 右室負荷時の心磁図 QRS 波に関する検討. —特徴的波形の定量的検討と出現部位. 臨床と研究 1987;64-2:482–488.
- [87] 森 博愛, 中屋 豊, 村上 昌, 野村昌弘, 竹内明美. 医療における人体磁界測定の有効性. 一心磁図の効力. 低温工学 1987;22:233.

- [88] 森 博愛, 中屋 豊, 須見昌輝, 野村昌弘, 福田善晴. 心磁図 T 波の臨床. 循環器科 1987;21-3: 261-269.
- [89] Cohen D., Kaufman L. S. Magnetic determination of the relationship between the S-T segment shift and the injury current produced by coronary artery occlusion. Circ Res 1975;36-3:414-424.
- [90] Fenici R. R., Romani G. L., Leoni R., Masselli M., Modena I. Magnetocardiographic recording of the His-Purkinje system activity in man. Jpn Heart J 1982;23-Suppl. I:728-730.
- [91] Erne S. N., Fenici R. R., Hahlbohm H. D., Lehmann H. P., and Trontelz Z. Beat to beat surface recording and averaging of His-Purkinje activity in man. J Electrocardiol 1983;16-4:355-362.
- [92] Erne S. N., Fenici R. R., Hahlbohm H. D., Lehmann H. P., Trontelz Z. High resolution magnetocardiography. Recording of His-Purkinje activity in man. Jpn Heart J 1982; 23-Suppl. I:722-727.
- [93] Saarinen M., Siltanen P., Karp P., Katila T. The normal magnetocardiogram I Morphology. Ann Clin Res 1978;10-Suppl. 21:1-43.
- [94] 中安紀美子, 中屋 豊, 野村昌弘, 藤野和也, 伊東進. 左室負荷疾患における心磁図—Isomagnetic map

および Single moving dipole 法による診断. Jpn J
Electrocardiol 1992;12:269-276.

- [95] 泉田直己, 浅野 優, 白井俊純. 37 チャンネル生体
磁気計測装置を用いた心臓起電力の移動双極子の分
析. 新しい医療機器研究 1994;2:25-34.
- [96] 日吉康長, 桜田春水, 本宮武司. 37 チャンネル生体
磁気計測装置を用いた Magnetocardiography
による心起電力の Moving Dipole 分析. 新しい医療
機器研究 1994;2:35-42.
- [97] Durrer D., van Dam R.T., Freud G.E., Janse M.J.,
Meiller F.L., Arzbaecher R.C. Total excitation of
the isolated human heart. Circulation 1970;41:
899-912.
- [98] Scher A.M., Yung A.C. The pathway of ventricular
depolarization in the dog. Cic Res 1956;4:
461-469.
- [99] 早川弘一, 比江嶋一昌編 臨床心臓電気生理学第 2
版 1994:pp41-43.
- [100] Wilson F.N. The precordial electrocardiogram. Am
Heart J 1944;27:19.
- [101] Sokolow M., Lyon T.P. The ventricular complex in
left ventricular hypertrophy as obtained by
unipolar precordial and limb leads. Am Heart J
1949;37:161.

- [102] 林 博史. QRST isointegral map の有用性. 日本臨床 1995;53:34-42.
- [103] 林 博史, 渡部真司, 山田芳司, 石川富久, 水谷真規子, 竹内 純. 脚ブロックを合併した心筋梗塞に対する QRST isointegral map の診断的有用性. 日内会誌 1987;76:1390-1395.
- [104] Hayashi H., Watabe S., Yabe S., Takami K., Ohsugi S., Hirai M., Mizutani M., Saito H. Diagnostic value of QRST isointegral maps in detecting myocardial infarction complicated by bundle branch block.. Circulation 1989;80:542-550.
- [105] Montague T. J., Smith E. R., Cameron D. A., Rautaharju P. M., Klassen G. A., Felmington C. S., Horacek B. M. Isointegral analysis of body surface maps. Surface distribution and temporal variability in normal subjects. Circulation 1981;63:1166-1172.
- [106] Montague T. J., Smith E. R., Spencer C. A., Johnstone D. E., Lalonde L. D., Bessoudo R. M., Gardner M. J., Anderson R. N. Body surface electrocardiographic mapping in inferior myocardial infarction. Manifestation of left and right ventricular involvement. Circulation 1983;67:665-673.
- [107] 八巻通安. ST-T Isointegral map. 日本臨床 1995;53:43-47.

- [108] Nakajima T., Kawakubo K., Toda I., Mashima S., Otake T., Iio M., Sugimoto T. ST-T isointegral analysis of exercise stress body surface mapping for identifying ischemic area in patients with angina pectoris. Am Heart J 1988;115:1013-1021.
- [109] 松井幹之, 八巻通安, 池田こずえ, 安井昭二. トレッドミル運動負荷による心筋虚血領域の推定. T isointegral map を用いて. 心電図 1990;10:192-200.
- [110] 中島敏明, 川久保清, 真島三郎, 戸田為久, 三輪篤子, 村川裕二, 倉智嘉久, 杉本恒明. 心筋梗塞例における運動負荷時心筋虚血領域の ST-T isointegral map による評価. 心臓 1988;20:247-256.
- [111] Abildskov J.A. Distribution of body surface ECG potentials in familial QT interval prolongation. Am J Cardiol 1981;47:480.
- [112] Igarashi H., Kubota I., Ikeda K., Yamaki M., Tsuiki K. Body surface mapping for the assessment of left ventricular hypertrophy in patients with essential hypertension. Jpn Circ J 1987;51:284-292.
- [113] Hayabuchi Y., Matsuoka S., Kubo M., Mori K., Tatara K., Kuroda Y. Usefulness of QRST isointegral maps for the diagnosis of right ventricular pressure overload in patients with surgically repaired tetralogy of Fallot complicated by right bundle branch block.

J Electrocardiol 1996;29:111-117.

- [114] Nomura N., Nakaya Y., Fujino K., Ishihara S., Katayama M., Takeuchi A., Watanabe K., Hiasa Y., Aihara T., Mori H. Magnetocardiographic studies of ventricular repolarization in old inferior myocardial infarction. Eur Heart J 1989;10:8-15.
- [115] Marwick T. Is silent ischemia painless because it is mild? J Am Coll Cardiol 1995;25:1513-1515.
- [116] 三浦 傳. 無症候性心筋虚血の現状. 日内会誌 1996;85:1546-1553.
- [117] Hechet H., DeBord L. Truly silent ischemia and relationship of chest pain and ST segment changes to the amount of ischemic myocardium. J Am Coll Cardiol 1994;23:369-376.
- [118] Langer A., Freeman M. R., Josse R. G., Armstrong P. W. Metaiodobenzylguanidine imaging in diabetes mellitus: assessment of cardiac sympathetic denervation and its relation to automatic dysfunction and silent myocardial ischemia. J Am Coll Cardiol 1995;25:610-618.
- [119] Yanagiya N., Abe T. Cardiac automatic activity during exercise-induced ischemia. Comparison between symptomatic and asymptomatic ischemia. Jpn Heart J 1994;35:547-548.
- [120] Fleg J. L. Prevalence and prognostic significance

of exercise-induced silent myocardial ischemia in apparently healthy subjects. Am J Cardiol 1992; 69:14B-18B.

- [121] Tzivoni D., Gavish A., Zin D., Gottlieb S., Moriel M., Keren A., Banai S., Stern S. Prognostic significance of ischemic episodes in patients with previous myocardial infarction. Am J Cardiol 1988;62:661-664.
- [122] 三浦 傳, 安部豊彦, 木村 衛. 無症候性心筋虚血の臨床. 臨床的意義とその頻度. 循環科学 1991;11: 1174-1178.
- [123] Bugiardini R., Borghi A., Sassone B., Pozzati A., Puddu P. Prognostic significance of silent myocardial ischemia in variant angina pectoris. Am J Cardiol 1991;68:1581-1586.
- [124] Kemp H. G. Left ventricular function in patients with the anginal syndrome and normal coronary arteriograms. Am J Cardiol 1973;32:375-376.
- [125] Cannon III R. O., Epstein S. E. Microvascular angina as a cause of chest pain with angiographically normal coronary arteries. Am J Cardiol 1988;61: 1338-1343.
- [126] Epstein S. E., Cannon III R. O. Site of increased resistance to coronary flow in patients with angina pectoris and normal epicardial coronary

arteries. *J Am Coll Cardiol* 1986;8:459-461.

- [127] Maseri A., Crea F., Kaski J.C., Crake T. Mechanisms of angina pectoris in syndrome X. *J Am Coll Cardiol* 1991;17:499-506.
- [128] Egashira K., Inou T., Hirooka Y., Yamada A., Urabe Y., Takeshita A. Evidence of impaired endothelium-dependent coronary vasodilation in patients with angina pectoris and normal coronary angiograms. *N Engl J Med* 1993;328:1659-1664.
- [129] Quyyumi A.A., Cannon III R.O., Panza J.A., Diodati J.G., Epstein S.E. Endothelial dysfunction in patients with chest pain and normal coronary arteries. *Circulation* 1992;86:1864-1871.
- [130] Mossneri M., Yarom R., Gotsman M.S., Hasin Y. Histologic evidence for small-vessel coronary artery disease in patients with angina pectoris and patent large coronary arteries. *Circulation* 1986;74:964-972.
- [131] Opherk D., Zebe H., Weihe E., Mall G., Gravert B., Mehmel H.C., Schwarz F., Kubler W. Reduced coronary dilatory capacity and ultrastructural changes of the myocardium in patients with angina pectoris but normal coronary arteriograms. *Circulation* 1981;63:817-825.
- [132] Kaski J.C., Rosano G., Collins P., Nihoyannopoulos P., Maseri A., Poole-Wilson P.A.

Cardiac syndrome X: clinical characteristics and left ventricular function. J Am Coll Cardiol 1995; 25:807-814.

[133] Kaski J.C., Crea F., Nihoyannopoulos P., Hackett D., Maseri A., Watson J., O'sullivan C. Transient myocardial ischemia during daily life in patients with syndrome X. Am J Cardiol 1986;58:1242-1247.

[134] 山香英三, 太刀川恭治, 一ノ瀬昇. 高温超伝導入門. オーム社(東京), 1989.

[135] Autenrieth G., Surawicz B., Cuo C. S. Sequence of repolarization on the ventricular surface in the dog. Am Heart J 1975;89:463-469.

[136] Abildskov J. The sequence of normal recovery of excitability in the dog heart. Circulation 1975; 52:442-446.

[137] Aronson R. S. Characteristics of action potentials of hypertrophied myocardium from rats with renal hypertension. Circ Res 1980;47:443-454.

[138] Downar E., Janse M., Durrer D. The effect of acute coronary occlusion on subepicardial transmembrane potentials in the intact porcine heart. Circulation 1977;56:217-224.

[139] 児島弘直. 酸化物高温超伝導体の大型単結晶. 日本物理学雑誌 1999;54:173-177.

- [140] 松葉博則. 超電導工学. 東京電機大学出版局(東京) 1997.
- [141] Suddarth S. A method for matching high-temperature superconductor resonators used for NMR signal pickup at 400 MHz. IEEE Trans Biomed Eng 1998;45:1061-1066.
- [142] Hall A.S., Alford N.M., Button T.W., Gilderdale D.J., Gehring K.A., Young I.R. Use of high temperature superconductor in a receiver coil for magnetic resonance imaging. Magn Reson Med 1991; 20:340-343.