

参 考 文 献

1. Esler M, Jennings G, Korner P, Willett I, Dudley F, Hasking G, Anderson W, Lambert G. Assessment of human sympathetic nervous system activity from measurements of norepinephrine turnover. *Hypertension* 1988; 11: 3-20
2. Cohn J, Levine TB, Olivari MT, Garberg V, Lura D, Francis G, Simon AB, Rector T. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Eng J Med.* 1984; 311: 819-823
3. Mathes P, Gudbjarnason S. Changes in norepinephrine stores in the canine heart following experimental myocardial infarction. *Am Heart J.* 1971; 81: 211-219
4. Hori M, Koretsune Y, Kagiya T, Watanabe Y, Iwakura K, Iwai K, Kitabatake A, Yoshida H, Inoue M, Kamada T. An increase in myocardial beta-adrenoceptors to compensate for postischaemic dysfunction following coronary micro-embolisation in dogs. *Cardiovasc Res.* 1989; 23: 424-431
5. Algra A, Tijssen JG, Roelandt JR, Pool J, Lubsen J. Heart rate variability from 24-hour electrocardiography and the 2-year risk for sudden death. *Circulation* 1993; 88: 180-185
6. La Rovere MT, Bigger JT Jr, Marcus FI, Mortara A, Schwartz PJ. Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (Autonomic Tone and Reflexes After Myocardial Infarction) Investigators. *Lancet* 1998; 351: 478-484
7. Asanoi H, Wada O, Miyagi K, Ishizaka S, Kameyama T, Seto H, Sasayama S. New redistribution index of nutritive blood flow to skeletal muscle during dynamic exercise. *Circulation* 1992; 85: 1457-1463

8. Weber KT, Janicki JS. Anaerobic threshold and aerobic capacity in the evaluation of chronic cardiac or circulatory failure. *Adv Cardiol.* 1986; 35: 79-87
9. Rowell LB. Human cardiovascular adjustments to exercise and thermal stress. *Physiol Rev.* 1974; 54: 75-159
10. Saito M, Naito M, Mano T. Different responses in skin and muscle sympathetic nerve activity to static muscle contraction. *J Appl Physiol.* 1990; 69: 2085-2090
11. Rowell LB, O'Leary DS. Reflex control of the circulation during exercise: chemoreflexes and mechanoreflexes. *J Appl Physiol.* 1990; 69: 407-418
12. Rowell LB, O'Leary DS. Reflex control of the circulation during exercise: chemoreflexes and mechanoreflexes. *J Appl Physiol.* 1990; 69: 407-418
13. Ekblom B, Goldbarg AN, Kilbom A, Astrand PO. Effects of atropine and propranolol on the oxygen transport system during exercise in man. *Scand J Clin Lab Invest.* 1972; 30: 35-42
14. Asanoi H, Kameyama T, Ishizaka S, Miyagi K, Sasayama S. Ventriculoarterial coupling during exercise in normal human subjects. *Int J Cardiol.* 1992; 36: 177-186
15. Miller TD, Balady GJ, Fletcher GF. Exercise and its role in the prevention and rehabilitation of cardiovascular disease. *Ann Behav Med.* 1997; 19: 220-229
16. Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97: 553-561

17. Linxue L, Nohara R, Makita S, Hosokawa R, Hata T, Okuda K, Hamazaki H, Fujita M, Sasayama S. Effect of long-term exercise training on regional myocardial perfusion changes in patients with coronary artery disease. *Jpn Circ J.* 1999; 63: 73-78
18. Lopez A, Vial R, Balart L, Arroyave G. Effect of exercise and physical fitness on serum lipids and lipoproteins. *Atherosclerosis.* 1974; 20: 1-9
19. Klachko DM, Lie TH, Cunningham EJ, Chase GR, Burns TW. Blood glucose levels during walking in normal and diabetic subjects. *Diabetes.* 1972; 21: 89-100
20. Pope SE, Stinson EB, Daughters GT, Schroeder JS, Ingels NB Jr, Alderman EL. Exercise response of the denervated heart in long-term cardiac transplant recipients. *Am J Cardiol.* 1980; 46: 213-218
21. Esler, M, Jennings G, Korner P, Willett I, Dudley F, Hasking G, Anderson W, Lambert G. Assessment of human sympathetic nervous system activity from measurements of norepinephrine turnover. *Hypertension* 1988; 11: 3-20
22. Wieland DM, Brown LE, Rogers WL, Worthington KC, Wu JL, Clinthorne NH, Otto CA, Swanson DP, Beierwaltes WH. Myocardial imaging with a radioiodinated norepinephrine storage analog. *J Nucl Med.* 1981; 22: 22-31
23. Sisson JC, Wieland DM, Sherma P, Mangner TJ, Tobes MC, Jacques S Jr. Metaiodobenzylguanidine as an index of the adrenergic nervous system integrity and function. *J Nucl Med.* 1987; 28: 1620-1624
24. Schofer J, Spielmann R, Schuchert A, Weber K, Schluter M. Iodine-123 meta-iodobenzylguanidine scintigraphy: a noninvasive method to demonstrate myocardial adrenergic nervous system disintegrity in patients with idiopathic dilated cardiomyopathy. *J Am Coll Cardiol.* 1988; 12: 1252-1258

25. Henderson EB, Kahn JK, Corbett JR, Jansen DE, Pippin JJ, Kulkarni P, Ugolini V, Akers MS, Hansen C, Buja LM. Abnormal I-123 metaiodobenzylguanidine myocardial washout and distribution may reflect myocardial adrenergic derangement in patients with congestive cardiomyopathy. *Circulation*. 1988; 78: 1192-1199
26. Jöbsis FF. Noninvasive, infrared monitoring of cerebral and myocardial oxygen sufficiency and circulatory parameters. *Science*. 1977; 198: 1264-1267
27. La Rovere MT, Bigger JT Jr, Marcus FI, Mortara A, Schwartz PJ. Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. *Lancet*. 1998; 351: 478-484
28. Sullivan MJ, Knight JD, Higginbotham MB, Cobb FR. Relation between central and peripheral hemodynamics during exercise in patients with chronic heart failure. Muscle blood flow is reduced with maintenance of arterial perfusion pressure. *Circulation*. 1989; 80: 769-781
29. Wilson JR, Martin JL, Ferraro N. Impaired skeletal muscle nutritive flow during exercise in patients with congestive heart failure: role of cardiac pump dysfunction as determined by the effect of dobutamine. *Am J Cardiol*. 1989; 53: 1308-1315
30. 渡辺重行, 仁科秀崇, 石山実樹, 山口 巖, 鰯坂隆一, 稲葉 武, 江田一彦, 武安法之, 坂本和彦, 外山昌弘, 杉下靖郎, 斎藤 巧, 山内孝義. 健常例における定常運動負荷時の運動筋酸素化率動態の意義, 近赤外線分光法(NIRS)による検討. *心臓* 1999; 31: 37-39
31. Ferrari M, Binzoni T, Quaresima V. Oxidative metabolism in muscle. *Phil Trans R Soc Lond*. 1997; B352: 677-683

32. Koyama K, Inoue T, Hasegawa A, Oriuchi N, Okamoto E, Tomaru Y, Endo K. Alternating myocardial sympathetic neural function of athlete's heart in professional cycle racers examined with iodine-123-MIBG myocardial scintigraphy. Ann Nucl Med. 2001; 15: 307-312
33. Weber K, Kinsewitz G, Janicki J, Fishman A. Oxygen utilisation and ventilation during exercise in patients with chronic cardiac failure. Circulation. 1982; 65: 1213-1223
34. Cohen-Solal A, Esanu Y, Logeart D, Pessonne F, Dubois C, Dreyfus G, Gourgon R, Merlet P. Cardiac metaiodobenzylguanidine uptake in patients with moderate chronic heart failure: relationship with peak oxygen uptake and prognosis. J Am Coll Cardiol. 1999; 33: 759-766
35. Watkins L Jr, Burton JA, Haber E, Cant JR, Smith FW, Barger AC. The renin-angiotensin-aldosterone system in congestive failure in conscious dogs. J Clin Invest. 1976; 57: 1606-1617
36. Schelling P, Fischer H, Ganzen D. Angiotensin and cell growth: a link to cardiovascular hypertrophy? J Hypertens. 1991; 9: 3-15
37. Furchtgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. Nature 1980; 288: 373-376
38. Sisson JC, Shapiro B, Meyers L, Mallette S, Mangner TJ, Wieland DM, Glowniak JV, Sherman P, Beierwaltes WH. Metaiodobenzylguanidine to map scintigraphically the adrenergic nervous system in man. J Nucl Med. 1987; 28: 1625-1636
39. Montserrat E, Albert F, Ricard S, Carina M, Teresa P, Joan CM, Lluis B, Ana MC, Ana T, Ignasi C. Influence of exercise rehabilitation on myocardial perfusion and sympathetic heart innervation in ischaemic heart disease. Eur J Nucl Med. 2000; 27 : 333-339

40. Chiba M, Nakamura M, Kanaya Y, Kobayashi N, Ueshima K, Kawazoe K, Hiramori K. Improvement in lower limb vasodilatory reserve and exercise capacity in patients with chronic heart failure due to valvular heart disease. *Eur Heart J.* 1997; 18: 1931-1936
41. Szlachcic J, Massie BM, Kramer BL, Topic N, Tubau J. Correlation and prognostic implication of exercise capacity in chronic congestive heart failure. *Am J Cardiol.* 1985; 55: 1037-1042
42. Weber K, Kinasewitz G, Janicki J, Fishman A. Oxygen utilisation and ventilation during exercise in patients with chronic cardiac failure. *Circulation* 1982; 65: 1213-1223
43. Belardinelli R, Georgiou D, Cianci G, Purcaro A. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999; 99: 1173-1182
44. Steffen-Batey L, Nichaman MZ, Goff DC, Frankowski RF, Hanis CL, Ramsey DJ, Labarthe DR. Change in level of physical activity and risk of all-cause mortality or reinfarction. The corpus christi heart project. *Circulation* 2000; 102: 2204-2209
45. Dominguez H, Torp-Pedersen C, Koeber L, Rask-Madsen C. Prognostic value of exercise testing in a cohort of patients followed for 15 years after acute myocardial infarction. *Eur Heart J.* 2001; 22: 300-306
46. Victor RG, Pryor SL, Secher NH, Mitchell JH. Effects of partial neuromuscular blockade on sympathetic nerve responses to static exercise in humans. *Circ Res.* 1989; 65: 468-476
47. Victor RG, Secher NH, Lyson T, Mitchell JH. Central command increases muscle sympathetic nerve activity during intense intermittent isometric exercise in humans. *Circ Res.* 1995; 76: 127-131

48. Ettinger S, Gray K, Whisler S, Sinoway L. Dichloroacetate reduces sympathetic nerve responses to static exercise. *Am J Physiol.* 1991; 261: H1653-H1658
49. Saito M, Kanao Y, Tanaka H, Sakai T. Muscle sympathetic nerve responses during progressive cycling exercise. *Adv Exerc Sports Physiol.* 1999; 5: 19-25
50. Saito M, Mano T. Exercise mode affect muscle sympathetic nerve responsiveness. *Jpn J Physiol.* 1991; 41: 143-151
51. Victor RG, Bertocci LA, Pryor SL, Nunnally RL. Sympathetic nerve discharge is coupled to muscle cell pH during exercise in humans. *J Clin Invest.* 1988; 82: 1301-1305
52. Seals DR, Johnson DG, Fregosi RF. Hypoxia potentiates exercise-induced sympathetic neural activation in humans. *J Appl Physiol.* 1991; 71: 1032-1040
53. Scherrier U, Pryor SL, Bertocci LA, Victor RG. Arterial baroreflex buffering of sympathetic activation during exercise-induced elevations in arterial pressure. *J Clin Invest.* 1990; 86: 1855-1861
54. Wallin BG, Eckberg DL. Sympathetic transients caused by abrupt alterations of carotid baroreceptor activity in humans. *Am J Physiol.* 1982; 242: H185-H190
55. Eckberg DL, Wallin BG. Isometric exercise modifies autonomic baroreflex responses in humans. *J Appl Physiol.* 1987; 63: 2325-2330
56. Sanders JS, Ferguson DW. Cardiopulmonary baroreflexes fail to modulate sympathetic responses during isometric exercise in humans : direct evidence from microneurographic studies. *J Am Coll Cardiol.* 1988; 12: 1241-1251

57. Scherrer U, Vissing SF, Victor RG. Effects of lower-body negative pressure on sympathetic nerve responses to static exercise in humans. Microneurographic evidence against cardiac baroreflex modulation of the exercise pressor reflex. *Circulation* 1988; 78: 49-59
58. Bini G, Hagbarth KE, Hynnen P, Wallin BG. Thermoregulatory and rhythmic generating mechanisms governing the sudomotor outflow in human cutaneous nerve. *J Physiol.* 1980; 306: 537-552
59. Niimi Y, Matsukawa T, Sugiyama Y, Shamsuzzaman AS, Ito H, Sobue G, Mano T. Effect of heat stress on muscle sympathetic nerve activity in humans. *J Auto Nerv Syst.* 1997; 63: 61-67
60. Saito M, Sone R, Ikeda M, Mano T. Sympathetic outflow to the skeletal muscle in humans increases during prolonged light exercise. *J Appl Physiol.* 1997; 82: 1237-1243
61. Ray CA, Gracey KH. Augmentation of exercise-induced muscle sympathetic nerve activity during muscle heating. *J Appl Physiol.* 1997; 82: 1719-1725
62. Wieland DM, Brown LE, Rogers WL, Worthington KC, Wu JL, Clinthorne NH, Otto CA, Swanson DP, Beierwaltes WH. Myocardial imaging with a radioiodinated norepinephrine storage analog. *J Nucl Med.* 1981; 22: 22-31
63. Jaques S Jr, Tobes MC, Sisson JC, Baker JA, Wieland DM. Comparison of the sodium dependency of uptake of meta-iodobenzylguanidine and norepinephrine into cultured bovine adrenomedullary cells. *Molecul Pharmacol.* 1984; 26: 539-546
64. Kline R, Swanson DP, Wieland DM, Thrall JH, Gross MD, Pitt B, Beierwaltes WH. Myocardial imaging in man with I-123 metaiodobenzylguanidine. *J Nucl Med.* 1981; 22: 129-132

65. Meredith IT, Eisenhofer G, Lambert GW, Dewar EM, Jennings GL, Esler MD. Cardiac sympathetic nervous activity in congestive heart failure: Evidence for increased neuronal norepinephrine release and preserved neuronal uptake. *Circulation* 1993; 88: 136-145
66. Eisenhofer G, Friberg P, Rundqvist B, Quyyumi AA, Lambert G, Kaye DM, Kopin IJ, Goldstein DS, Esler MD. Cardiac sympathetic nerve function in congestive heart failure. *Circulation* 1996; 93: 1667-1676
67. Nakajo M, Shimabukuro K, Yoshimura H, Yonekura R, Nakabeppu Y, Tanoue P, Shinohara S. Iodine-131-metaiodobenzylguanidine intra- and extravesicular accumulation in the rat heart. *J Nucl Med.* 1986; 27: 84-89
68. Schofer J, Spielmann R, Schuchert A, Weber K, Schuler M. Iodine-131-metaiodobenzylguanidine scintigraphy : A noninvasive method to demonstrate myocardial adrenergic system disintegrity in patients with idiopathic dilated cardiomyopathy. *J Am Coll Cardiol.* 1988; 12: 1252-1258
69. Henderson EB, Kahn JK, Corbett JR, Jansen DE, Pippin JJ, Kulkarni P, Ugolini V, Akers MS, Hansen C, Buja LM. Abnormal I-123-metaiodobenzylguanidine myocardial washout and distribution may reflect myocardial adrenergic derangement in patients with congestive cardiomyopathy. *Circulation* 1988; 78: 1192-1199
70. Barker D, Saito M. Autonomic innervation of receptors and muscle fibers in skeletal muscle. *Proc R Soc Lond B Biol Sci.* 1981; 212: 317-332
71. Roatta S, Windhorst U, Ljubisavljevic M, Johansson H, Passatore M. Sympathetic modulation of muscle spindle afferent sensitivity to stretch in rabbit jaw closing muscles. *J Physiol.* 2002; 540: 237-248

72. Passatore M, Grassi C, Filippi GM. Sympathetically-induced development of tension in jaw muscles: the possible contraction of intrafusal muscle fibres. Pflugers Arch. 1985; 405: 297-304
73. Boyd IA. The response of fast and slow nuclear bag fibres and nuclear chain fibers in isolated cat muscle spindles to fusimotor stimulation, and the effect of intrafusal contraction on the sensory endings. Q J Exp Physiol. 1976; 61: 203-254
74. Sugihara H, Shiga K, Terada K, Kinoshita N, Taniguchi Y, Ito K, Adachi Y, Ushijima Y, Nakagawa M, Maeda T. Effect of exercise-induced activation of sympathetic nerve activity on clearance of ¹²³I-MIBG from the myocardium. Ann Nucl Med. 1998; 12: 175-178