

図表のタイトル一覧

第四章 損傷組織に対するクライオセラピーが足関節不安定性，神経筋機能および機能性パフォーマンスに及ぼす影響

4-1 損傷組織に対するクライオセラピーが足関節不安定性に及ぼす影響

Table 4-1-1. Characteristics of subjects

Table 4-1-2. Ankle stability evaluated by stress X-ray measurement

Fig. 4-1-1. Stress X-ray measurement of anterior talar translocation and talar tilt

Fig. 4-1-2. Effects of cryotherapy on anterior talar translocation of sprained ankle

Fig. 4-1-3. Effects of cryotherapy on talar tilt of sprained ankle

4-2 損傷組織に対するクライオセラピーが筋反応時間および機能性パフォーマンスに及ぼす影響

Table 4-2-1. Characteristics of subjects

Table 4-2-2. Mean muscle reaction times and functional performances before, after, 24 h after cryotherapy

Fig. 4-2-1. Sudden ankle inversion stress for measurement of the peroneal reaction time

Fig. 4-2-2. Recording of peroneal reaction time

Fig. 4-2-3. Peroneus longus reaction times (patient group)

Fig. 4-2-4. Peroneus brevis reaction times (patient group)

Fig. 4-2-5. Effect of cryotherapy on peroneus longus reaction times (patient group)

Fig. 4-2-6. Effect of cryotherapy on peroneus brevis reaction times (patient group)

Fig. 4-2-7. Effect of cryotherapy on peroneus longus reaction times (healthy group)

Fig. 4-2-8. Effect of cryotherapy on peroneus brevis reaction times (healthy group)

第五章 損傷組織に対するクライオセラピーが微小血管の血管透過性、血行動態および白血球動態に及ぼす影響

5-1 損傷組織に対するクライオセラピーが血管透過性に及ぼす影響

Fig. 5-1-1. The glass stage for observation of the rat cremaster muscle

Fig. 5-1-2. Contusion model

Fig. 5-1-3. Real-time confocal laser-scanning intra-vital microscope system and TV camera system

Fig. 5-1-4. Fluorescent images of the rat cremaster muscle microvasculature

Fig. 5-1-5. Average FITC-dextran 150s leakage area of the rat cremaster muscle microvasculature

5-2 損傷組織に対するクライオセラピーが血行動態および白血球動態に及ぼす影響

Fig. 5-2-1. Images of a rat cremaster muscle venular microvessel

Fig. 5-2-2. Vessel internal diameter under different treatments

Fig. 5-2-3. Percent change in blood flow rate under different treatments

Fig. 5-2-4. Percent change in erythrocyte velocity under different treatments

Fig. 5-2-5. Number (no.) of rolling leukocytes per length of venule (100 μ m) under different treatments

Fig. 5-2-6. Number (no.) of adhering leukocytes per length of venule (100 μm) under different treatments