

## 文献

- AAHPERD (1980) Test manual. Reston VA: American Alliance for Health, Physical Education, Recreation and Dance.
- Adner MM and Castelli PW (1980) Elevated high-density lipoprotein levels in marathon runners. *JAMA* 243: 534-535.
- American College of Sports Medicine (1995) Guidelines for Exercise Testing and Prescription (5th ed.). Williams & Wilkins, Philadelphia, pp. 49-190.
- American College of Sports Medicine (1986) Guidelines for Exercise Testing and Prescription (3rd ed.). Lea & Febiger, Philadelphia, pp. 31-44.
- 浅見俊雄, 足立長彦, 山本恵三, 北川薫, 広田公一 (1976) 主観による運動強度の選択について—10分間走の場合—. *体育科学* 4: 1-5.
- Åstrand PO and Ryhming I (1954) A nomogram for calculation of aerobic capacity (physical fitness) from pulse rate during submaximal work. *J Appl Physiol* 7: 218-220.
- Åstrand I (1960) Aerobic work capacity in men and women with special reference to age. *Acta Physiol Scand* 49: 1-92.
- Atomi Y and Miyashita M (1984) Maximal oxygen uptake of obese middle-aged women related to body composition and total body potassium. *J Sports Med* 24: 212-218.
- Bandura A (1977) Self-efficacy: Toward a unifying theory of behavioral change. *Psychol Rev* 84: 191-215.
- Beaver WL, Wasserman K and Whipp BJ (1985) Improved detection of lactate threshold during exercise using a log-log transformation. *J Appl Physiol* 59: 1936-1940.
- Beaver WL, Wasserman K, Whipp BJ (1986) A new method for detecting the anaerobic threshold by gas exchange. *J Appl Physiol* 60: 2020-2027.
- Beverly JW, Ruth GD, David CN and Diane EB (1993) Validation of 1-mile test in elderly women. *J Aging Phys Activity* 1: 13-21.

- Blair SN, Kohl HW, Paffenbarger RS, Clark DG, Cooper KH and Gibbons LW (1989) Physical fitness and all-cause mortality: a prospective study of healthy men and women. *JAMA* 262: 2395-2401.
- Borg GAV (1973) Perceived exertion: A note on "history" and methods. *Med Sci Sports* 5: 90-93.
- Brouha L (1943) The step test: A simple method of measuring physical fitness for hard muscular work in adult man. *Rev Canad Biol* 2: 86-91.
- Burke EJ (1979) Individualized fitness programme. *J Phys Educ Rec* 50: 35-37.
- Caiizzo VJ, Davis JA, Ellis JF., Azus JL and Vandagriff R (1982) Comparison of gas exchange indices used to detect the anaerobic threshold. *J Appl Physiol* 53: 1184-1189.
- Ceci R and Hassmen P (1991) Self-monitored exercise at three different RPE intensities in treadmill vs field running. *Med Sci Sports Exerc* 23: 732-738.
- Concu A and Marcello C (1993) Stroke volume response to progressive exercise in athletes engaged in different types of training. *Eur J Appl Physiol* 66: 11-17.
- Cooper KH (1968) A means of assessing maximal oxygen intake. *JAMA* 203: 135-138.
- Conley DL and Krahenbuhl GS (1980) Running economy and distance running performance of highly trained athletes. *Med Sci Sports Exerc* 12: 6-18.
- Cooper KH, Pollock ML, Martin RP, White SR, Linnerud AC and Jackson A (1976) Physical fitness levels vs selected coronary risk factors. A cross-sectional study. *JAMA* 236: 166-169.
- Costill DL, Miller JM and Fink WJ (1980) Energy metabolism in diabetic distance runners. *Physician Sportmed* 8: 64-71.
- Coyle EF, Martin WH and Ehsani AA (1983) Blood lactate threshold in some well-trained ischemic heart disease patients. *J Appl Physiol* 54: 18-23.
- Davies CTM (1968) Limitations to the prediction of maximum oxygen intake from cardiac frequency measurements. *J Appl Physiol* 24: 700-706.

- Demello JJ, Cureton KJ and Boineau RE (1987) Ratings of perceived exertion at lactate threshold in trained and untrained men and women. *Med Sci Sports Exerc* 19: 354-362.
- DeVries HA and Klafs CE (1965) Prediction of maximal O<sub>2</sub> intake from submaximal tests. *J Sports Med Phys Fit* 5: 207-214.
- Dishman RK, Graham RE, Holly RG and Tieman JG (1991) Estimates of Type A behavior do not predict perceived exertion during graded exercise. *Med Sci Sports Exerc* 23: 1276-1282.
- Dobelin WV, Astrand I and Bergstrom A (1967) An analysis of age and other factors related to maximal oxygen uptake. *J Appl Physiol* 22: 934-938.
- Doolittle TL and Bigbee R (1968) The twelve-minute run-walk: a test of cardio-respiratory fitness of adolescent boys. *Res Quart* 39: 491-495.
- Doston C (1988) Health fitness standards -aerobic endurance-. *J O P E R D* 57: 26-31.
- Dunbar CC, Robertson RJ, Baun R, Blandin MF, Metz K, Burdett R and Goss FL (1992) The validity of regulating exercise intensity by ratings of perceived exertion. *Med Sci Sports Exerc* 24: 94-99.
- Duncan JJ (1985) The effects of aerobic exercise on plasma catecholamines and blood pressure in patients with mild essential hypertension. *JAMA* 254: 2609-2613.
- Ebbeling CB, Ward A, Puleo EM, Widrick J and Rippe JM (1991) Development of a single-stage submaximal treadmill walking test. *Med Sci Sports Exerc* 23: 996-973.
- Ekholm B and Goldbarg AN (1971) The influence of physical training and other factors on the subjective rating of perceived exertion. *Acta Physiol Scand* 83: 399-406.
- Eston BG, Davie BL and Williams JG (1987) Use of perceived effort ratings to control exercise intensity in young healthy adults. *Eur J Appl Physiol* 56: 222-224.
- Fairshter RD, Walters J, Salness K, Fox M, Minh VD and Wilson AF (1983) A comparison of incremental exercise tests during cycle and treadmill ergometry. *Med Sci Sports Exerc* 15: 549-554.

- Farrell PA, Wilmore JH, Coyle EF, Billing JE and Costill DL (1979) Plasma lactate accumulation and distance running performance. *Med Sci Sports* 11: 338-344.
- Felts WM, Crouse S and Brunetz M (1988) Influence of aerobic fitness on ratings of perceived exertion during light to moderate exercise. *Percept Mot Skills* 67: 671-676.
- Fenstermaker KL, Plowman SA and Looney MA (1992) Validation of the rockport fitness walking test in females 65 years and older. *Res Quart* 63: 322-327.
- Fletcher GF (1990) Exercise standards: a statement for health professionals. *Circulation* 82: 2286-2322.
- Fox EL (1973) A simple, accurate technique for predicting maximal aerobic power. *J Appl Physiol* 35: 914-916.
- Friedman M and Rosenman RH (1959) Association of specific overt behavior pattern with blood and cardiovascular findings: Blood cholesterol level, blood clotting time, incidence of arcus senilis, and clinical coronary artery disease. *JAMA* 169: 1286-1296.
- Frisch RE, Wyshak G, Ibright TE, Albright NL and Schiff I (1986) Lower prevalence of diabetes in female former college athletes compared with nonathletes. *Diabetes* 35: 1101-1105.
- Glass SC, Knowlton RG and Becque MD (1992) Accuracy of RPE from graded exercise to establish exercise training intensity. *Med Sci Sports Exerc* 24: 1303-1307.
- Glassford RG, Baycroft GHY, Sedgwick AW and Macnab RBJ (1965) Comparison of maximal oxygen uptake values determined by predicted and actual methods. *J Appl Physiol* 20: 509-513.
- Hage P (1981) Perceived exertion: One measure of exercise intensity. *Physician Sportsmed* 19: 136-143.
- Haisman MF (1971) Assessment of the exercise capacity of young men. *Ergonomics* 14: 449-456.
- Heath GW, Hagberg JM, Ehsani AA (1981) A physiological comparison of young and older endurance athletes. *J Appl Physiol* 51: 634-640.

- Hermiston R and Faulkner JA (1971) Prediction of maximal oxygen uptake by a stepwise regression technique. *J Appl Physiol* 30: 833-837.
- Hetzler RK, Seip RL, Boutcher SH, Pierce E, Snead D and Weltman A (1991) Effect of exercise modality on ratings of perceived exertion at various lactate concentrations. *Med Sci Sports Exerc* 13: 377-383.
- Hollman W and Hettinger Th. (1990) *Sportmedizin arbeits-und trainingsgrundlagen*. Schattauer, Stuttgart, pp. 303-304.
- Issekutz B, Birkhead BNC and Rodahl K (1962) Use of respiratory quotients in assessment of aerobic work capacity. *J Appl Physiol* 17: 47-50.
- 猪飼道夫 (1967) 日本人の体力. 日経新書, 東京, pp. 181-184.
- Jackson A S and Coleman AE (1976) Validation of distance run test for elementary school children. *Res Quart* 47: 86-94.
- Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure (1993) The fifth report of the joint national committee on detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 153, 154-183.
- 金子公宥, 淵本隆文, 末井健作, 田路秀樹, 矢邊順子, 西田充 (1986) 簡便な屋内持久走テストの提案—シャトル・スタミナテスト (SST) の考案と検討—. *体育の科学*. 36: 809-815.
- 金子公宥, 淵本隆文, 森島博 (1988) 小学校 (7~12歳) における持久性能力の発達—シャトルスタミナテストによる調査結果から—. *体育科学* 16: 45-51.
- 金子公宥, 淵本隆文, 森島博 (1989) シャトルスタミナテストでみた持久性能力の発達—13~18歳男女を対象に—. *体育科学* 17: 48-53.
- 形本静夫 (1980) 最大酸素摂取量の間接測定. *体育の科学* 30: 823-827.
- 川村仁視, 新畑茂充, 山田岳志, 藤井勝紀 (1993) *現代人の健康と運動*. 杏林書院, 東京, pp. 184-186.
- Kim HS, Tanaka K and Maeda, K. (1992) Metabolic and cardiorespiratory responses during cycling exercise below, at, and above lactate threshold. *Jpn J Appl Physiol* 22: 307-315.

- Kittredge JM, Rimmer JH and Looney MA (1994) Validation of the rockport fitness walking test for adults with mental retardation. *Med Sci Sports Exerc* 26: 95-102.
- Kline GM, Porcari JP, Hintermeister R, Freedson PS, Ward A, Mccarron RF, Ross J and Rippe JM (1987) Estimation of  $\dot{V}O_{2max}$  from a one-mile track walk, gender, age, and body weight. *Med Sci Sports Exerc* 19: 253-259.
- 厚生省 (1998) 国民衛生の動向. 厚生省統計協会, 東京, pp. 101-107.
- 厚生省 (1989) 日本人の栄養所要量「健康づくりのための運動所要量」. 第一出版, 東京, pp. 151-156.
- 熊谷もりえ, 中垣内真樹, 西嶋尚彦, 田中喜代次 (1997) 個人情報と主観的運動強度を導入した全身持久性の簡易推定法—若年成人男性について—. *体力科学* 46: 179-188.
- Kravitz L, Robergs AR, Heyward HV, Wagner RD and Powers K (1997) Exercise mode and gender comparisons of energy expenditure at self-selected intensities. *Med Sci Sports Exerc* 29: 1028-1035.
- Laukkanen RMT, Oja P, Pasanen ME and Vuori IM (1992) Criterion validity of a 2-km walking test for estimating the maximal aerobic power in over-weight adults. *Int J Obes* 16: 263-268.
- Laukkanen RMT, Oja P, Pasanen ME and Vuori IM (1993) Criterion validity of a 2-km walking test for predicting the maximal oxygen uptake of moderately to highly active middle-aged adults. *Scand J Med Sci Sports* 3: 267-272.
- LeBlanc, J. (1979) Effect of physical training and adiposity on glucose metabolism and  $^{125}I$ -insulin binding. *J. Appl. Physiol.* 46: 235-239.
- Leger LA and Lambert J (1982) A maximal multistage 20m shuttle run test to predict  $\dot{V}O_{2max}$ . *Eur J Appl Physiol* 49: 1-12.
- Legge BJ and Banister EW (1986) The Åstrand-Ryhming nomogram revisited. *J Appl Physiol* 61: 1203-1209.
- Lohman TG (1981) Skinfolds and body density and relation to body fatness: A review. *Human Biol* 53: 181-225.
- Maksud MG and Coutts KD (1971) Application of Cooper 12-minute run-walk test to young males. *Res Quart* 42: 54-59.

- Margaria R, Aghemo P and Rovell E (1965) Indirect determination of maximal O<sub>2</sub> consumption in men. *J Appl Physiol* 20: 1070-1073.
- Maritz JS, Morrison JF, Peter J, Strydom NB and Wyndham CH (1961) A practical method of estimating an individual's maximal oxygen intake. *Ergonomics* 4: 97-122.
- Mastropaolo JA (1970) A prediction of maximal O<sub>2</sub> consumption in middle-aged men by multiple regression. *Med Sci Sports* 2: 124-127.
- McCauley E and Courneya K (1992) Self-efficacy relationships with affective and exertion responses to exercise. *J Appl Soc Psychol* 22: 321-326.
- McArdle WD, Katch FI and Katch VL (1981) Training for anaerobic and aerobic power. (In) *Exercise Physiology*. Lea & Febiger, Philadelphia, pp. 266-285.
- Metz KF and Alexander JF (1971) Estimation of maximal oxygen intake from submaximal work parameters. *Res Quart* 42: 54-59.
- 宮村実晴 (1986) 最大酸素摂取量の再検討. *体育の科学* 36: 358-364.
- 宮下充正, 武井義明, 福田裕之 (1984) PWC75%HR<sub>max</sub>の全身持久性の評価尺度としての妥当性の検討. *J J Sports Sci* 3: 559-562.
- 宮下充正, 山本義春, 田村真一, 篠原稔, 武藤芳照 (1989) 換気性作業閾値が無酸素性作業閾値を与える. *体育の科学* 39: 391-396.
- Morgan WP (1973) Psychological factors influencing perceived exertion. *Med Sci Sports* 5: 97-103.
- Morgan WP and Borg GA (1976) Perception of effort and the prescription of physical activity. (In) *Mental Health and Emotional Aspects of Sports*. American Medical Association, pp. 126-129.
- Morgan WP (1981) Psychophysiology of self-awareness during vigorous physical activity. *Res Quart* 52: 385-427.
- Morris JN, Everitt MG, Pollard R, Chave SPW and Semmence AM (1980) Vigorous exercise in leisure-time protection against coronary heart disease. *Lancet* 2: 1207-1210.
- 宗像恒次 (1996) 現代日本人社会の病原と予防策. (In) *行動科学からみた健康と病気* (初版). メヂカルフレンド社, 東京, pp. 45-47.

- 村上寿利, 進藤宗洋, 田中宏暁, 熊谷秋三, 生田純男, 佐々木淳 (1988a) 冠動脈硬化性心疾患危険因子の判定手法としての推定最大酸素摂取量の有効性. 動脈硬化 15: 1665-1672.
- 村上寿利, 進藤宗洋, 田中宏暁, 熊谷秋三, 今村英夫, 生田純男, 神宮純江, 高橋紀子, 佐々木淳 (1988b) 冠動脈硬化性心疾患危険因子の判定手法としての推定最大酸素摂取量の有効性 (第2報) -女性についての検討-. 動脈硬化 16: 495-500.
- Nagamine S and Suzuki S (1964) Anthropometry and body composition of Japanese young men and women. Hum. Biol 35: 8-15.
- Nakadomo F, Tanaka K, Hazama T and Maeda K (1990) Validation of body composition assessed by bioelectrical impedance analysis. Jpn J Appl Physiol 20: 321-330.
- 中川秀樹, 形本静夫 (1995) 12分間歩行テストによる全身持久性評価の可能性. 教育医学 40: 260-266.
- 中西とも子, 田中喜代次, 李美淑, 磯野香代子, 早川洋子, 佐久義昭, 中垣内真樹, 青山正恵, 張美蘭, 細川淳一 (1996) 中年肥満女性に対する院内監視型運動指導の有用性について-健康度指標“活力年齢”の改善という視点に着目して-. 肥満研究 2: 28-33.
- 中村好男, 山本義春 (1989) AT測定法のゆくえ. 体育の科学 39: 391-396.
- 西田保, 猪俣公宏, 岡沢祥訓 (1983) 運動負荷条件下におけるperceived exertionとパーソナリティ要因との関連性. 体育学研究 28: 129-139.
- Noble BJ (1982) Clinical applications of perceived exertion. Med Sci Sports Exerc 14: 406-411.
- 信田宜司, 長谷川陽三, 田中喜代次 (1991) 心拍変動フィードバック管理負荷制御方式に基づく正弦波的運動負荷トレーニングの肥満および高血圧に及ぼす影響. 臨床スポーツ医学 8: 65-73.
- 盧昊成, 田中喜代次, 竹田正樹, 海野英哉, 檜山輝男 (1996) 本態性高血圧症女性に対する運動療法の血圧および活力年齢への効果. 体力科学 45: 91-100.



- 涌井忠昭, 高橋繁浩, 北川薫, 石河利寛 (1987) 水泳および水中歩行における運動強度の指標としての心拍数, 運動スピード, 主観的運動強度の有効性. 東海保健体育科学 9: 1-9.
- 大蔵倫博, 田中喜代次, 中垣内真樹, 佐久義昭 (1996) 中高年有疾患者を対象としたAT水準の運動強度に関する提案. 日本生理人類学会誌 1: 27-31.
- 大蔵倫博, 上野Mリンダ, 田中喜代次 (1998) 最大下多段階漸増負荷法による自転車運動中の自覚的運動強度を利用した全身持久性体力の推定方法: 一般若年男性の検討. 体育学研究 43: 102-116.
- Oja P, Laukkanen RM, Pasanen ME, Tyry T and Vuori IM (1991) A 2-km walking test for assessing the cardiorespiratory fitness of healthy adults. *Int J Sports Med* 12: 356-362.
- 小野寺孝一, 宮下充正 (1976) 全身持久性運動における主観的運動強度と客観的運動強度の対応性 - Rating of perceived exertionの観点から -. 体育学研究 21: 191-203.
- Paffenbarger RS, Hyde RT, Wing AL and Rauramaa R (1986) Physical activity, all-cause mortality and longevity of college alumni. *New Engl J Med* 314: 605-613.
- Pandolf KB, Billings DS, Drolet LL, Pimental NA and Sawka MN (1984) Differentiated ratings of perceived exertion and various physiological responses during prolonged upper and lower body exercise. *Eur J Appl Physiol* 53: 5-11.
- Peters RK, Cady LD, Bischoff DP, Bernstein L and Pike MC (1983) Physical fitness and subsequent myocardial infarction in healthy workers. *JAMA* 249: 3052-3056.
- Pollock ML, Miller HS and Wilmore JH (1974) Physiological characteristics of champion American track athletes 40 to 75 years of age. *J Gerontol* 29: 645-649.
- Pollock ML (1988) Prescribing exercise for fitness and adherence. (In) Dishman RK (Ed.) *Exercise adherence: Its impact on public health*. Human Kinetics, Champaign, pp. 259-277.
- Powell KE, Thompson PD and Casperson CJ (1987) Physical activity and the incidence of coronary heart disease. *Ann Rev Public Health* 8: 253-287.

- Purvis JW, Cureton KJ (1981) Ratings of perceived exertion at the anaerobic threshold. *Ergonomics* 24: 295-300.
- Rejeski DR, Morley D and Miller HS (1983) Cardiac rehabilitation: coronary-prone behavior as a moderator of graded exercise test performance. *J Cardiac Rehabil* 3: 339-346.
- Rowell LB, Taylor HL and Wang Y (1964) Limitations to prediction of maximal oxygen intake. *J Appl Physiol* 19: 919-927.
- 坂野雄二, 東篠光彦 (1986) 一般性セルフ・エフィカシー尺度作成の試み. *行動療法研究* 12: 73-82.
- 佐藤祐造, 中井直也, 下村吉治 (1997) 糖尿病の人の運動. *保健の科学* 39: 4-9.
- Scharff-Olson M, Williford HN, and Smith FH. (1992) The heart rate  $\dot{V}O_2$  relationships of aerobic dance: a comparison target heart rate methods. *J Sports Med Phys fitness* 32: 372-377.
- Seals DR, Hagberg JM and Spina RJ (1994) Enhanced left ventricular performance in endurance trained older men. *Circulation* 89: 109-120.
- Seip RL, Snead D, Pierce EF, Stein P and Weltman A (1991) Perceptual responses and blood lactate concentration: effect of training state. *Med Sci Sports Exerc* 23: 80-87.
- Sheffield LT, Maloof JA, Sawyer JA and Koitman D (1978) Maximal heart rate and treadmill performance of healthy women in relation to age. *Circulation* 57: 79-84.
- Shephard RJ (1967) The prediction of maximal oxygen consumption using a new progressive step test. *Ergonomics* 10: 1-15.
- Siconolfi SE, Cullinane M, Carleton RA and Thompson PD (1982) Assessing  $\dot{V}O_{2max}$  in epidemiologic studies: modification of the Åstrand-Ryhming test. *Med Sci Sports Exerc* 14: 335-338.
- Skinner SJ, Hutsler R, Bergsteinova V and Buskirk ER (1973) Perception of effort during different types of exercise and under different environmental conditions. *Med Sci Sports* 5: 110-115.

- Spiriduso WW (1995) Physical dimensions of aging. *Human Kinetics*, Champaign, pp. 7-8.
- Storer TW, Davis JA and Caiozzo VJ (1989) Accurate prediction of  $\dot{V}O_{2max}$  in cycle ergometry. *Med Sci Sports Exerc* 22: 704-711.
- 田路秀樹, 末井健作, 金子公宥 (1992) シヤトルスタミナテスト (STT) における加齢の影響と評価基準. *体育の科学* 42: 367-372.
- 竹島伸生, 田中喜代次, 小林章雄 (1997) 高齢者の健康づくり—運動処方の実際と課題—. *メディカルレビュー社*, 東京.
- 竹島伸生, 田中喜代次, 小林章雄, 渡辺丈真, 鷺見勝博, 加藤孝之 (1992) 高齢者の全身持久性評価における種々の間接法の妥当性. *体力科学* 41: 295-303.
- 竹島伸生 (1990) 高齢者の乳酸性閾値ならびに最大酸素摂取量による有酸素性作業能の評価に関する研究. *愛知医科大学医学会誌* 18: 403-416.
- 竹田正樹, 田中喜代次, 浅野勝己, 渡辺寛, 檜山輝男 (1993) Lactate Thresholdを用いた冠動脈心疾患患者用の運動プログラムのあり方. *呼吸と循環* 41: 999-1003.
- Tanaka K (1985) Validity of endurance performance test as a possible substitute for cardiorespiratory fitness. *Osaka City Univ J Health Sci Phys Educ* 21: 19-28.
- Tanaka K, Watanabe H, Konishi Y, Mitsuzono R, Sumida S, Tanaka S, Fukuda T and Nakadomo F (1986a) Longitudinal associations between anaerobic threshold and distance running performance. *Eur J Appl Physiol* 55: 248-252.
- Tanaka K, Yoshimura T, Sumida S, Mitsuzono R, Tanaka S, Konishi Y, Watanabe H, Yamada T and Maeda K (1986b) Transient responses in cardiac function below, at, and above anaerobic threshold. *Eur J Appl Physiol* 55: 356-361.
- 田中喜代次, 吉村隆喜, 奥田豊子, 小西洋太郎, 角田聡, 出村慎一, 岡田邦夫 (1986a) AT水準以上の強度を基準とした完全監視型持久性運動療法および不完全監視型食事療法の併用が肥満者の健康・体力に及ぼす効果. *体力研究* 62: 26-40.
- 田中喜代次, 松本健, 榎原潤三, 村上晴彦, 中川敬 (1986b) 光電脈波方式による全身持久性の評価. *大阪市立大学保健体育学研究紀要* 22: 7-15.

- Tanaka K, Kim H S, Wakita M, Nakadomo F, Hazama T and Maeda K (1989) Effects of aerobic conditioning plus caloric restriction in obese middle-aged women. *Jpn J Appl physiol* 19: 495-504.
- 田中喜代次 (1989) 持久性競技者の競技成績とAT. *体育の科学* 39: 382-390.
- 田中喜代次, 山地繁信, 岡田邦夫 (1989) 踏台昇降テストによる全身持久性の年齢別評価基準値の作成—邦人男性について—. *体育の科学* 39: 725-729.
- Tanaka K, Takeshima N, Kato T, Niihata S and Ueda K (1990) Critical determinants of endurance performance in middle-aged and elderly endurance runners with heterogeneous training habits. *Eur J Appl Physiol* 59: 443-449.
- 田中喜代次 (1990) 高齢者に対する運動処方とその注意事項. *体育の科学* 40: 940-948.
- 田中喜代次, 信田宜司, 長谷川陽三, 金炫秀, 松浦義行 (1990) 心拍変動フィードバック管理負荷制御方式による本邦成人女性の全身持久性の評価. *筑波大学体育科学系紀要* 13: 215-223.
- Tanaka K (1991) Cardiorespiratory and lactate responses to a 1-hour submaximal running at the lactate threshold. *Ann Physiol Anthropol* 10: 155-162.
- Tanaka K, Nakadomo F, Watanabe K, Inagaki A, Kim HK and Matsuura Y (1992) Body composition prediction equations based on bioelectrical impedance and anthropometric variables for Japanese obese women. *Am J Hum Biol* 4: 739-745.
- 田中喜代次 (1995) 全身持久性能力の簡易評価法に関する提案 (第1報: 最大下での12分間トレッドミル歩行テストの妥当性). *臨床スポーツ医学* 12: 217-223.
- 田中喜代次, 金禧植, 李美淑, 佐藤喜久, 大浜三平, 上向井千佳子, 長谷川陽三, 檜山輝男 (1995a) 質問紙によるヒトの全身持久性体力の簡易評価法に関する提案—成人女性を対象として—. *臨床スポーツ医学* 12: 438-444.
- 田中喜代次, 水野康, 浅野勝己 (1995b) 有酸素性トレーニングの生理と効用 [7] 老年期の一般健康者に対する有酸素性トレーニング. *臨床スポーツ医学* 12: 1160-1366.

- 田中喜代次, 中垣内真樹, 大蔵倫博, 重松良祐, 盧昊成, 金憲経 (1998) 高齢者の全身持久性体力を簡便に求める新たな方法の提案. 臨床スポーツ医学 15: 843-848.
- Taylor HL, Buskirk E and Henschel A (1955) Maximal oxygen intake as an objective measure of cardiorespiratory performance. *J Appl Physiol* 8: 73-80.
- 谷口興一 (編) (1993) 心肺運動負荷テスト. 南江堂, 東京.
- Warren BJ, Doston RG and Butterworth DE (1985) Improved detection of lactate threshold during exercise using a log-log transformation. *J Appl Physiol* 59: 1936-1940.
- Wasserman K, Whipp BJ, Koyal SN and Beaver WL (1973) Anaerobic threshold and respiratory gas exchange during exercise. *J Appl Physiol* 35: 236-243.
- Widrick J, Ward A, Ebbeling C, Clemente E and Rippe MJ (1991) Treadmill validation of an over-ground walking test to predict peak oxygen consumption. *Eur J Appl Physiol* 64: 304-308.
- Wilmore JH and Costill DL (1994) Physical activity for health and fitness. (In) Wilmore JH, Costill DL (Eds.) *Physiology of sports and exercise. Human Kinetics, Champaign*, pp. 467-530.
- Wyndham CH, Strydom NB, Morrison JF, Peter J and Potgieter ZU (1959) Maximum oxygen intake and maximum heart rate during strenuous work. *J Appl Physiol* 18: 927-936.
- 山地啓司 (1992) 最大酸素摂取量の科学. 杏林書院, 東京.
- 山地啓司 (1997) 心拍数 (脈拍数) の測定意義・方法と主観的運動強度. *ランニング学研究* 8: 15-35.
- 吉田敬義, 石河利寛 (1978) 呼吸循環機能からみた幼児の持久走について. *体育学研究* 23: 59-65.