

表のタイトル一覧

第4章 運動習慣および大動脈伸展性が収縮期血圧に及ぼす影響

- Table 1. Subject characteristics (n=678). Mean, SD, and number of abnormality subjects (%).
- Table 2. Simple correlation coefficients matrix between variables (n=678).
- Table 3. Contribution rates of each factor for SBP.
- Table 4. Contribution rates of each factor for DBP.
- Table 5. Contribution rates of each factor for APWVI.
- Table 6. Contribution rates of each factor for TC.
- Table 7. Contribution rates of each factor for TG.
- Table 8. Contribution rates of each factor for UA.
- Table 9. Contribution rates of each factor for 2h-OGTT.
- Table 10. Contribution rates of each factor for %Fat.
- Table 11. Contribution rates of each factor for Cigarettes.
- Table 12. Contribution rates of each factor for Alcohol.
- Table 13. Contribution rates of each factor for PAI.

第5章 継続的運動が大動脈伸展性に及ぼす影響

- Table 14. Cross-sectional comparisons of physical characteristics in runner, active, and sedentary groups.
- Table 15. Anthropometric, blood pressure, and PAI values of the runner, active, and sedentary groups.

第6章 身体活動の著しい低下が大動脈伸展性に及ぼす影響

—視覚障害者における検討—

Table 16. The characteristics of blind, runner, and sedentary groups.

Table 17. Simple matrix of correlation coefficients between variables (n=40).

第7章 中高年期以降の運動が大動脈伸展性に及ぼす影響

7.1 運動の開始時期および継続期間が大動脈伸展性に及ぼす影響

Table 18. Physical characteristics of subjects.

7.2 短期間・低強度の運動トレーニングが大動脈伸展性に及ぼす影響

Table 19. Characteristics of subjects before and after 6 months of training.