

Acknowledgments

The author would like to express his sincere gratitude to Professor Mitsuhiro Ikura, Ontario Cancer Institute and Department of Medical Biophysics, University of Toronto, Canada, for his cordial guidance, continuing discussion, and warm encouragement. He wishes his sincere thanks to Dr. Toshiyuki Tanaka, Center for Tsukuba Advanced Research Alliance, and Dr. Mark B. Swindells, Helix Research Institute, for cordial guidance, helpful discussion, and hearty instruction. The author also thanks to Professor Tadashi Baba, Institute of Applied Biochemistry, University of Tsukuba, Professor Kazunari Taira, Institute of Applied Biochemistry, University of Tsukuba, and Professor Yoshimasa Tanaka, Institute of Biological Science, University of Tsukuba, for helpful advice and discussion for the preparation of the thesis.

It is great pleasure to express his sincere gratitude to Professor Eisuke Munekata, Institute of Applied Biochemistry, University of Tsukuba, for his cordial guidance and discussion for the preparation of the thesis.

The author is grateful to Dr. Claude Klee, National Institute of Health, USA, for kindly providing the *Xenopus* calmodulin expression system, Ms. Kit I. Tong and Ms. Kyoko L. Yap, University of Toronto, Canada, for the preparation of calmodulin, Mr. Frank Delaglio and Dr. Daniel S. Garrett, National Institute of Health, USA, for NMR data processing and analysis. He wishes his sincere thanks to Dr. Hiroshi Tokumitsu, Ms. Naomi Takahashi, Dr. Toshio Furuya, Helix Research Institute, for kindly providing CaMKK peptides, the mutagenesis studies, helpful discussions and hearty encouragement, Professor Norio Matsushima, Sapporo Medical College, and Professor Yoshinobu Izumi, University of Yamagata and the members of his laboratory for their instruction and helpful discussions on the small-angle X-ray scattering experiments, Dr. Hisayuki Yokokura, Dr. Tsutomu Inoue, and Professor Hiroyoshi Hidaka for the synthesis of the novel CaM antagonists,

biochemical tests of them, and helpful discussions, Dr. Masaya Orita and Mr. Ken-ichi Suzumura, Yamanouchi Pharmaceutical Co. Ltd., for NMR experiments and discussions.

Finally the author wishes to express his cordial appreciation to all the members of the laboratory of the chemistry for leads, Yamanouchi Pharmaceutical Co., Ltd. and Center for Tsukuba Advanced Research Alliance.

This doctoral program was supported by Yamanouchi Pharmaceutical Co., Ltd.