

## ACKNOWLEDGMENTS

I would like to express my sincere appreciation to Prof. Eisuke Munekata (Institute of Applied Biochemistry, University of Tsukuba) and Dr. Toshiyuki Tanaka (Center for Tsukuba Advanced Research Alliance, University of Tsukuba) for generous understanding and continuous encouragements to my research and helpful instructions during the course of this study. I also wishes my sincere thanks to Prof. Yoshimasa Tanaka (Institute of Biological Sciences, University of Tsukuba), Prof. Kazunari Taira (Institute of Applied Biochemistry, University of Tsukuba), and Prof. Tadashi Baba (Institute of Applied Biochemistry, University of Tsukuba) for helpful advice and discussion for the preparation of the thesis.

I am particularly indebted to Prof. Mitsuhiro Ikura (Ontario Cancer Institute, University of Toronto) and Prof. Masayori Inouye (Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey) for fruitful collaborations, stimulating discussion and warm encouragement. I sincerely appreciate Prof. Kimiko Umemoto (International Christian University), Dr. Takami Tomiyama (University of Osaka), and Ms. Kit I. Tong (Center for Tsukuba Advanced Research Alliance, University of Tsukuba) for helpful advice and encouragement for preparation of the thesis.

I thank to Prof. Masatsune Kainosho for providing the labeled ATP analogue sample. I am also indebted to Dr. Rieko Ishima, and Dr. Dingjiang Liu (Ontario Cancer Institute, University of Toronto) for technical support on NMR measurements and encouragement, and all the members of Laboratory of Structural Biology, Center for Tsukuba Advanced Research Alliance, University of Tsukuba for useful discussions and supports.

Finally, I thank ex-colleagues in Teijin Institute for Bio-Medical Research, Teijin Limited, and friends and my family for continuous encouragements.