

引用文献

Allison T, McCarthy G, Wood CC and Jones SJ: Potentials evoked in human and monkey cerebral cortex by stimulation of the median nerve. *Brain*, 114: 2465-2503, 1991.

荒木秀夫, 西平賀昭, 藤田紀盛: 反応動作の α -blockingと緩電位変動との関連について. 体力科学, 30: 148-156, 1981.

Angel R, Boylls CC and Weinrich M: Cerebral evoked potentials and somatosensory perception. *Neurology*, 34: 123-126, 1984.

Arezzo J, Vaughan HG Jr and Koss B: Relationship of neuronal activity to gross movement-related potentials in monkey pre- and postcentral cortex. *Brain Res.*, 132: 362-369, 1977.

Asami T, Ikai M and Shibayama H: The body reaction time in athletes. *Proc. Int. Congr. Sport Sci.*, 505-507, Tokyo, 1964.

浅見高明: 反射と反応に関する基礎的研究. 東京教育大学スポーツ研究所所報, 9: 31-42, 1971.

Berger H: Über das Elektrenkephalogramm des Menschen. *Arch. f. Psychiat.*, 87: 527-570, 1929.

Bocker KBE, Forget R and Brunia CH: The modulation of somatosensory evoked potentials during the foreperiod of a forewarned reaction time task. *Electroenceph. clin. Neurophysiol.*, 88: 105-117, 1993.

Boschert J and Deecke L: Handedness, footedness and finger and toe movement-related cerebral potentials. *Hum. Neurobiol.*, 5: 235-243, 1986.

Botwinick J and Thompson LW: Premotor and motor components of reaction time. *J. Exp. Psychol.*, 71: 9-15, 1966.

Brunia CH, Scheirs JG and Haagh SA: Changes of Achilles tendon reflex amplitudes during a fixed foreperiod of four seconds. *Psychophysiology*, 19: 63-70, 1982.

Campbell KB, Courchesne E, Picton TW and Squires KC: Evoked potential correlates of human information processing. *Biol. Psychol.*, 8: 45-68, 1979.

Chapman CE, Jiang W and Lamarre Y: Modulation of lemniscal input during conditioned arm movements in the monkey. *Exp. Brain Res.*, 72: 316-334, 1988.

Cheyne D, Endo H, Takeda T and Weinberg H: Sensory feedback contributes to early movement-evoked fields during voluntary finger movements in humans. *Brain Res.*, 771: 196-202, 1997.

Corbetta M, Miezin FM, Shulman GL and Petersen SE: A PET study of visuospatial attention. *J. Neurosci.*, 13: 1202-1226, 1993.

Cunnington R, Bradshaw JL and Iansek R: The role of the supplementary motor area in the control of voluntary movement. *Human Mov. Sci.*, 15: 627-647, 1996.

Cunnington R, Iansek R, Johnson KA and Bradshaw JL: Movement-related potentials in Parkinson's disease. Motor imagery and movement preparation. *Brain*, 120: 1339-1353, 1997.

Davis PA: Effects of acoustic stimuli on the waking human brain. *J. Neurophysiol.*, 2: 494-499, 1939.

Davis H, Davis PA, Loomis AL, Harvey EN and Hobart GI: Electrical reactions of the human brain to auditory stimulation during sleep. *J. Neurophysiol.*, 2: 500-514, 1939.

Dawson GD: Cerebral responses to electrical stimulation of peripheral nerve in man. *J. Neurol. Neurosurg. Psychiat.*, 10: 137-140, 1947.

Deecke L, Scheid P and Kornhuber H: Distribution of readiness potential, Pre-motion positivity, and motor potential of the human cerebral cortex preceding voluntary finger movements. *Exp. Brain Res.*, 7: 158-168, 1969.

Desmedt JE and Robertson D: Differential enhancement of early and late components of the cerebral somatosensory evoked potentials during forced-paced cognitive tasks in man. *J. Physiol.*, 271: 761-782, 1977.

Desmedt JE: P300 in serial tasks: An essential post-decision closure mechanism. *Progr. Brain Res.*, 54: 682-686, 1980.

Desmedt JE, Huy NT and Bourguet M: The cognitive P40, N60 and P100 components of somatosensory evoked potentials and the earliest electrical signs of sensory processing in man. *Electroenceph. clin. Neurophysiol.*, 56: 272-282, 1983.

Desmedt JE and Tomberg C: Mapping early somatosensory evoked potentials in selective attention: Critical evaluation of control conditions used for titrating by difference the cognitive P30, P40, P100 and N140. *Electroenceph. clin. Neurophysiol.*, 74: 321-346, 1989.

Elbert T, Hommel J and Lutzenberger W: The perception of neckercube reversal interacts with the Bereitschaftspotential. *Int. J. Psychophysiol.*, 3: 5-12, 1985.

Elbert T, Rockstroh B, Hampson S, Pantev C and Hoke M: The magnetic counterpart of the contingent negative variation. *Electroenceph. clin. Neurophysiol.*, 92: 262-272, 1994.

Fattapposta F, Amabile G, Cordischi MV, Venanzio Di, Foti A, Pierelli F, D'Alessio C, Pigozzi F, Parisi A and Morrocitti C: Long-term practice effects on a new skilled motor learning: An electrophysiological study. *Electroenceph. clin. Neurophysiol.*, 99: 495-507, 1996.

Fernandez T, Harmony T, Rodriguez M, Reyes A, Marosi E and Bernal J: Test-retest reliability of EEG spectral parameters during cognitive tasks: I. Absolute and relative power. *Int. J. Neurosci.*, 68: 255-261, 1993.

Ford JM, MacPherson L and Kopell BS: Differences in readiness potential associated with push-button construction. *Psychophysiology*, 9: 564-567, 1972.

Forss N, Hari R, Samelin R, Ahonen A, Hamalainen M, Kajola M, Knuutila J and Simola J: Activation of the human posterior parietal cortex by median nerve stimulation. *Exp. Brain Res.*, 99: 309-315, 1994.

麓 正樹, 西平賀昭, 下田政博, 宮 達夫, 八田有洋: 運動準備期の持続的な随意収縮がCNVとH反射に及ぼす影響. 脳波と筋電図, 27: 258-267, 1999.

Gaillard AWK: Effects of warning-signal modality on the contingent negative variation(CNV). *Biol. Psychol.*, 4: 139-154, 1976.

Garcia-Larrea L, Lukaszewicz AC and Mauguiere F: Somatosensory responses during selective spatial attention: The N120-to-N140 transition. *Psychophysiology*, 32: 526-537, 1995.

Gemba H and Sasaki K: Potential related to no-go reaction of go/no-go hand movement task with colour discrimination in human. *Neurosci. Lett.*, 101: 263-268, 1989.

Gemba H, Sasaki K and Tsujimoto T: Cortical field potentials associated with hand movements triggered by warning and imperative stimuli in the monkey. *Neurosci. Lett.*, 113: 275-280, 1990.

Goodin DS, Squires KC and Starr A: Variations in early and late event-related components of the auditory evoked potential with task difficulty. *Electroenceph. clin. Neurophysiol.*, 55: 680-686, 1983.

Grafton ST, Mazziotta JC, Presty S, Friston KJ, Frackowiak RS and Phelps ME: Functional anatomy of human procedural learning determined with regional cerebral blood flow and PET. *J. Neurosci.*, 12: 2542-2548, 1992.

Groves PM and Eason RG: Effects of attention and activation on the visual evoked cortical potential and reaction time. *Psychophysiology*, 5: 394-398, 1969.

Haagh SA and Brunia CH: Anticipatory response-relevant muscle activity, CNV amplitude and simple reaction time. *Electroenceph. clin. Neurophysiol.*, 61: 30-39, 1985.

萩原 仁, 調枝孝治: 人間の知覚一運動行動, 不昧堂, 東京, 1975.

Hamalainen H, Kekoni J, Sams M, Reinikainen K and Naatanen R: Human somatosensory evoked potentials to mechanical pulses and vibration: contributions of SI and SII somatosensory cortices to P50 and P100 components. *Electroenceph. clin. Neurophysiol.*, 75: 13-21, 1990.

Hamano T, Luders HO, Ikeda A, Collura TF, Comair YG and Shibasaki H: The cortical generators of the contingent negative variation in humans: A study with subdural electrodes. *Electroenceph. clin. Neurophysiol.*, 104: 257-268, 1997.

Hikosaka O, Sakai K, Miyauchi S, Takino R, Sasaki Y and Putz B: Activation of human presupplementary motor area in learning of sequential procedures: A functional MRI study. *J. Neurophysiol.*, 76: 617-621, 1996.

Hink RF, Deecke L and Kornhuber HH: Force uncertainty of voluntary movement and human movement-related potentials. *Biol. Psychol.*, 16: 197-210, 1983.

平松謙一, 秋本 優, 丹羽真一, 亀山知道, 斎藤 治, 山崎清之, 伊藤憲治: 選択反応課題における分裂病患者の反応時間とP300潜時. 精神医学, 27: 1055-1063, 1985.

平松謙一, 秋本 優, 福田正人, 斎藤 治, 亀山知道: 課題無関連刺激のP300成分—効果的課題遂行戦略のための冗長な情報の利用—. 脳波と筋電図, 14: 264-272, 1986.

Hoshiyama M and Sheean G: Changes of somatosensory evoked potentials preceding rapid voluntary movement in Go/No-go choice reaction time task. *Cognit. Brain Res.*, 7: 137-142, 1998.

Hultin L, Rossini P, Romani GL, Hogstedt P, Tecchio F and Pizzella V: Neuromagnetic localization of the late component of the contingent negative variation. *Electroenceph. clin. Neurophysiol.*, 98: 435-448, 1996.

Hyman R: Stimulus information as a determinant of reaction time. *J. Exp. Psychol.*, 45: 188-196, 1953.

猪飼道夫, 浅見高明, 芝山秀太郎: 全身反応時間の研究とその応用. *Olympia*, 7: 210-219, 1961.

Ikeda A, Luders HO, Burgess RC and Shibasaki H: Movement-related potentials associated with single and repetitive movements recorded from human supplementary motor area. *Electroenceph. clin. Neurophysiol.*, 89: 269-277, 1993.

池田昭夫, 柴崎 浩: 運動皮質機構の非侵襲的研究—小脳と基底核の関与—. *臨床神経学*, 35: 1522-1524, 1995.

池田昭夫, 柴崎 浩: ヒト補足運動野機能についての非侵襲的検討. *臨床脳波*, 38: 152-157, 1996.

Ishihara T and Yoshii N: Multi-variate analytic study of EEG and mental activity in juvenile delinquents. *Electroenceph. clin. Neurophysiol.*, 33: 71-80, 1972.

石井みゆき, 宮内利郎, 萩元 浩, 遠藤青磁, 梶原 智, 山沢 浩, 岡嶋次郎: 脳波基礎活動とP300成分. *臨床脳波*, 37: 754-758, 1995.

岩瀬勝彦: 準備電位と意欲. *臨床脳波*, 15: 702-706, 1973.

Jasiukaitis P and Hakerem G: The effect of prestimulus alpha activity on the P300. *Psychophysiology*, 25: 157-165, 1988.

Jasper HH and Penfield W: Electrocorticograms in man: Effects of voluntary movement upon the electrical activity of the precentral gyrus. *Arch. Psychiat. Neurol.*, 183: 163-174, 1949.

Jenkins IH, Brooks DJ, Nixon PD, Frackowiak RS and Passingham RE: Motor sequence learning: A study with positron emission tomography. *J. Neurosci.*, 14: 3775-3790, 1994.

Joshiassen RC, Shagass C, Roemer RA, Slepner SS and Czartorysky B: Early cognitive components of somatosensory event-related potentials. *Int. J. Psychophysiol.*, 9: 139-149, 1990.

Kakigi R, Koyama S, Hoshiyama M, Watanabe S, Shimojo M and Kitamura Y: Gating of somatosensory evoked responses during active finger movements magnetoencephalographic studies. *J. Neurol. Sci.*, 128: 195-204, 1995.

Kakizaki T: Evaluation of mental task strain based on occipital beta activity and subjective rating of task difficulty. *Eur. J. Appl. Physiol.*, 54: 466-470, 1985.

河辺章子, 大築立志: フェイント刺激による誤反応の修正. 体育学研究, 27: 217-227, 1982.

Kawashima R, Matsumura M, Sadato N, Naito E, Waki A, Nakamura S, Matsunami K, Fukuda H and Yonekura Y: Regional cerebral blood flow changes in human hand brain related to ipsilateral and contralateral complex movements. --a PET study--. *Eur. J. Neurosci.*, 10: 2254-2260, 1998.

Keller I and Heckhausen H: Readiness potentials preceding spontaneous motor acts: Voluntary vs. involuntary control. *Electroenceph. clin. Neurophysiol.*, 76: 351-361, 1990.

Komiyama T and Tanaka R: The differences in human spinal motoneuron excitability during the foreperiod of a motor task. *Exp. Brain Res.*, 79: 357-364, 1990.

Kornhuber HH and Deecke L: Hirnpotentialanderungen bei Willkürbewegungen und passiven Bewegungen des Menschen. Bereitschaftspotential und reafferente potentielle. *Pflugers Arch. ges. Physiol.*, 284: 1-17, 1965.

Kristeva R, Cheyne W, Lang G, Lindinger G and Deecke L: Movement-related potentials accompanying unilateral and bilateral finger movements with different inertial loads. *Electroenceph. clin. Neurophysiol.*, 75: 410-418, 1990.

Kutas M, McCarthy G and Donchin E: Augmenting mental chronometry: The P300 as a measure of stimulus evaluation time. *Science*, 197: 792-795, 1977.

Kutas M and Donchin E: Preparation to respond as manifested by movement-related brain potentials. *Brain Res.*, 202: 95-115, 1980.

- Loveless NE and Sanford AJ: Slow potential correlates of preparatory set. *Biol. Psychol.*, 1: 303-314, 1974.
- Luft AR, Skalej M, Stefanou A, Klose U and Voigt K: Comparing motion- and imagery-related in the human cerebellum: A functional MRI study. *Human Brain Mapping*. 6: 105-113, 1998.
- Luppino G, Matelli M, Camarda RM, Gallese V and Rizzolatti G: Multiple representations of body movements in mesial area 6 and the adjacent cingulate cortex: an intra-cortical microstimulation study in the macaque monkey. *J. Comp. Neurol.*, 311: 463-482, 1991.
- Magliero A, Bashore TR, Coles MGH and Donchin E: On the dependence of P300 latency on stimulus evaluation processes. *Psychophysiology*, 21: 171-186, 1984.
- McAdam DW and Seales DM: Bereitschaftspotential enhancement with increased level of motivation. *Electroenceph. clin. Neurophysiol.*, 27:73-75, 1969.
- McCarthy G and Donchin E: A metric for thought: A comparison of P300 latency and reaction time. *Science*, 211: 77-80, 1981.
- McCarthy G, Wood CC, Williamson PD and Spencer DD: Task-dependent field potentials in human hippocampal formation. *J. Neurosci.*, 9: 4253-4268, 1989.
- Mecklinger A, Kramer AF and Strayer DL: Event-related potentials and EEG components in a semantic memory search task. *Psychophysiology*, 29: 104-119, 1992.
- 中村隆一：筋電図反応時間による運動発現の検討. 臨床脳波, 30: 566-572, 1988.
- Nashmi R, Mendonca AJ and MacKay WA: EEG rhythms of the sensorimotor region during hand movements. *Electroenceph. clin. Neurophysiol.*, 91: 456-467, 1994.
- 西平賀昭, 藤田紀盛, 荒木秀夫:隨意動作に先行する脳運動関連電位. 体力科学, 27: 140-148, 1978.
- Nishihira Y, Araki H and Ishihara A: Comparison among various movement task of cerebral motor potential preceding voluntary movement. *Electromyogr. clin. Neurophysiol.*, 29: 179-184, 1989.

Nishihira Y, Araki H, Funase K, Imanaka K, Suzuki J and Takemiya T: Somatosensory evoked potentials following voluntary movement during upper arm compression. *Electromyogr. clin. Neurophysiol.*, 36:21-28, 1996.

西平賀昭：神経系の運動調節適応能。竹宮 隆，石河利寛（編），運動適応の科学－トレーニングの科学的アプローチー，杏林書院，東京，pp.135-152, 1998.

Nishihira Y, Ohno T, Hatta A, Fumoto M, Kaneda T, Tokitou S, Imanaka K and Funase K: P300 before and after transient hard exercise. *Adv. Exerc. Sports Physiol.*, 5: 49-54, 1999.

Oddie SD and Bland BH: Hippocampal formation theta activity and movement selection. *Neurosci. Biobehavior Reviews*, 22: 221-231, 1998.

Ohtsuki T: Decrease in grip strength induced by simultaneous bilateral exertion with reference to finger strength. *Ergonomics*, 24: 37-48, 1981.

大塚立志：「たくみ」の科学。朝倉書店，東京，1988。

Orgogozo JM and Larsen B: Activation of the supplementary motor area during voluntary movements in man suggest it works as supramotor area. *Science*, 206: 847-850, 1979.

Patterson JV, Michalewski HJ and Starr A: Latency variability of the components of auditory event-related potentials to infrequent stimuli in ageing, Alzheimer-type dementia, and depression. *Electroenceph. clin. Neurophysiol.*, 71: 450-460, 1988.

Pfefferbaum A, Ford J, Johnson R Jr, Wenegrat B and Kopell B: Manipulation of P3 latency: Speed vs. accuracy instructions. *Electroenceph. clin. Neurophysiol.*, 55: 188-197, 1983.

Pfefferbaum A, Ford J, Weller BJ and Kopell B: ERPs to response production and inhibition. *Electroenceph. clin. Neurophysiol.*, 60: 423-434, 1985.

Pfurtscheller G and Aranibar A: Event-related desynchronization detected by power measurements of scalp EEG. *Electroenceph. clin. Neurophysiol.*, 42: 817-826, 1977.

Pfurtscheller G: Central beta rhythm during sensorimotor activities in man. *Electroenceph. clin. Neurophysiol.*, 51: 253-264, 1981.

- Pfurtscheller G, Stancak A Jr and Neuper C: Post-movement beta synchronization. A correlate of an idling motor area? *Electroenceph. clin. Neurophysiol.*, 98: 281-293, 1996.
- Picton TW: The P300 wave of the human event-related potential. *J. Clin. Neurophysiol.*, 9: 456-479, 1992.
- Polich J and Lardon MT: P300 and long-term physical exercise. *Electroenceph. clin. Neurophysiol.*, 103: 493-498, 1997.
- Polich J: On the relationship between EEG and P300: Individual differences, ageing, and ultradian rhythms. *Int. J. Psychophysiol.*, 26: 299-317, 1997.
- Pribram KH and McGuinness D: Arousal, activation, and effort in the control of attention. *Psychol. Rev.*, 82: 116-149, 1975.
- Price GW: The effect of pre-stimulus alpha activity on the auditory P300 paradigm: A prospective study. *Brain Topogr.*, 9: 169-176, 1997.
- Ritter W, Ford J, Gaillard AWK, Harter MR, Kutas M, Naatanen R, Polich J, Renault B and Rohrbaugh J: Cognition and event-related potentials. I. The relation of negative potentials and cognitive processes. *Ann. N. Y. Acad. Sci.*, 425: 24-38, 1984.
- 齊藤 治, 豊嶋良一:歴史と概論. 丹羽真一, 鶴 紀子(編), 事象関連電位, 新興医学出版社, 東京, pp.3-21, 1997.
- Salmelin R and Hari R: Spatiotemporal characteristics of sensorimotor neuromagnetic rhythms related to thumb movement. *Neurosci.*, 60: 537-550, 1994.
- 佐々木和夫, 玄番央恵, 南部 篤, 松崎竜一:No-Go電位の電気生理学的および磁気生理学的研究. 日本生体磁気学会誌, 5: 36-39, 1992.
- Scheich H and Simon O: The reaction of alpha activity during exercise. *Electroenceph. clin. Neurophysiol.*, 29: 217-218, 1970.
- Seitz RJ and Roland PE: Learning of sequential finger movements in man: A combined kinematic and positron emission tomography(PET) study. *Eur. J. Neurosci.*, 4:154-465, 1992.
- Shaw JC: Intention as a component of the alpha-rhythm response to mental activity. *Int. J. Psychophysiol.*, 24: 7-23, 1996.

- Shibasaki H, Barrett G, Halliday E and Halliday AM: Components of the movement-related cortical potential and their scalp topography. *Electroenceph. clin. Neurophysiol.*, 49: 213-226, 1980.
- Shibasaki H, Sadato N, Lyshkow H, Yonekura Y, Honda M, Nagamine T, Suwazono S, Magata Y, Ikeda A, Miyazaki M, Fukuyama H, Asato R and Konishi J: Both primary motor cortex and supplementary motor area play an important role in complex finger movement. *Brain*, 116: 1387-1398, 1993.
- Simonetta M, Clanet M and Rascol O: Bereitschaftspotential in a simple movement or in a motor sequence starting with the same simple movement. *Electroenceph. clin. Neurophysiol.*, 81: 129-134, 1991.
- Simson R, Vaughan HG Jr and Ritter W: The scalp topography of potentials in auditory and visual Go/NoGo tasks. *Electroenceph. clin. Neurophysiol.*, 43: 864-875, 1977.
- Squires NK, Squires KC and Hillyard SA: Two varieties of long-latency positive waves evoked by unpredictable auditory stimuli in man. *Electroenceph. clin. Neurophysiol.*, 38: 387-401, 1975.
- Squires NK, Squires KC and Hillyard SA: Decision-related cortical potentials during an auditory signal detection task with cued observation intervals. *J. Exp. Psychol.: Human Perception & Performance*, 1: 268-279, 1975.
- Staines WR, Brooke JD, Angerilli PA and McIlroy WE: Phasic modulation of somatosensory potentials during passive movement. *Neuroreport*, 7: 2971-2974, 1996.
- Stancak A Jr and Pfurtscheller G: Desynchronization and recovery of beta rhythms during brisk and slow self-paced finger movements in man. *Neurosci. Lett.*, 196: 21-24, 1995.
- Starr A and Cohen LG: Gating of somatosensory evoked potentials begins before the onset of voluntary movement in man. *Brain Res.*, 348: 183-186, 1985.
- Starr A, Sandroni P and Michalewski HJ: Readiness to respond in a target detection task: Pre- and post-stimulus event-related potentials in normal subjects. *Electroenceph. clin. Neurophysiol.*, 96: 76-92, 1995.

- Stippich C, Freitag P, Kassubek J, Soros P, Kamada K, Kober H, Scheffler K, Hopfengartner R, Bilecen D, Radu EW and Vieth JB: Motor, Somatosensory and auditory cortex localization by fMRI and MEG. *Neuroreport*, 9: 1953-1957, 1998.
- Stoffel EJ: Uncertainty and processing routes in the selection of a response: An S-R compatibility study. *Acta Psychologica*, 94: 227-252, 1996.
- Surwillo WW: The relation of simple response time to brain wave frequency and the effect of age. *Electroenceph. clin. Neurophysiol.*, 15: 105-114, 1963.
- Sutton S, Braren M and Zubin J: Evoked potential correlates of stimulus uncertainty. *Science*, 150: 1187-1188, 1965.
- 忠井俊明, 福井義久, 上月清司, 井爪 尚, 加藤光彦, 村田伸文, 中嶋照夫: 分裂病の情報処理障害—事象関連電位による情報処理モデルからの検討—. 精神医学, 29: 817-828, 1987.
- 竹内 賢, 丹羽真一: 課題遂行と事象関連電位—反応時間を含めて—. 丹羽真一, 鶴 紀子(編), 事象関連電位, 新興医学出版社, 東京, pp.65-75, 1997.
- Tanaka R: Reciprocal Ia inhibition during voluntary movements in man. *Exp. Brain Res.*, 21: 529-540, 1974.
- Tanji J: The supplementary motor area in cerebral cortex. *Neurosci. Res.*, 19: 231-268, 1994.
- 丹治 順: 運動と行動の神経機構. 神經進歩, 40: 997-999, 1996.
- Tarkka IM and Hallett M: Cortical topography of premotor and motor potentials preceding self-paced voluntary movement of dominant and non-dominant hands. *Electroenceph. clin. Neurophysiol.*, 75: 36-43, 1990.
- Taylor MJ: Bereitschaftspotential during the acquisition of a skilled motor task. *Electroenceph. clin. Neurophysiol.*, 45: 568-567, 1978.
- Tecce JJ, Savignano-Bowman J and Meinbresse D: Contingent negative variation and the distraction arousal hypothesis. *Electroenceph. clin. Neurophysiol.*, 41: 277-286, 1976.

Tomporowski PDS and Ellis NR: Effects of exercise on cognitive process: A review. *Psychol. Bull.*, 99: 338-346, 1986.

van Boxtel GJ and Brunia CH: Motor and non-motor aspects of slow brain potentials. *Biol. Psychol.*, 38: 37-51, 1994.

Vanderwolf CH: The electrocorticogram in relation to physiology and behaviour: A new analysis. *Electroenceph. clin. Neurophysiol.*, 82: 165-175, 1992.

Vidal F, Bonnet M and Macar F: Programming the duration of a motor sequence: Role of the primary and supplementary motor areas in man. *Exp. Brain Res.*, 106: 339-350, 1995.

Walter WG, Cooper R, Aldridge VJ, McCallum WC and Winer AL: Contingent negative variation: An electric sign of sensorimotor association and expectancy in the human brain. *Nature*, 203: 380-384, 1964.

Wascher E, Verleger R, Jaskowski P and Wauschkuhn B: Preparation for action: An ERP study about two tasks provoking variability in response speed. *Psychophysiology*, 33: 262-272, 1996.

Weiss T, Sust M, Beyer L, Hansen E, Rost R and Schmalz T: Theta power decreases in preparation for voluntary isometric contractions performed with maximal subjective effort. *Neurosci. Lett.*, 193: 153-156, 1995.

Wildgruber D, Erb M, Klose U and Grodd W: Sequential activation of supplementary motor area and primary motor cortex during self-paced finger movement in human evaluated by functional MRI. *Neurosci. Lett.*, 227: 161-164, 1997.

山口修平：選択的視覚注意と事象関連電位. 脳波と筋電図, 26: 371-380, 1998.

Yingling CD: Cognition, action and mechanisms of EEG asymmetry. In: Pfurtscheller G, Buser FH, Lopes da Silva and Petsche(Eds.), *Rhythmic EEG activities and cortical functioning*. Elsevier/North Holland, Amsterdam, 79-90, 1980.